

DT # 44984

QA: NA

7/11/05

MOL.20060404.0026

STABILITY CONSTANTS OF NP(V) COMPLEXES WITH FLUORIDE
AND SULFATE AT VARIABLE TEMPERATURES

Yuanxian Xia^{1*}, Judah I. Friese¹, Dean A. Moore¹, Linfeng. Rao²

1. Pacific Northwest National Laboratory

PO Box 999

Richland, WA 99352, USA

2. Lawrence Berkeley National Laboratory

1 Cyclotron Road

Berkeley, CA 94720, USA

* Corresponding Author: Phone (509) 373-5767; Fax (509) 372-1632

E-mail: yuanxian.xia@pnl.gov.

ABSTRACT

A solvent extraction method was used to determine the stability constants of Np(V) complexes with fluoride and sulfate in 1.0 M NaClO₄ from 25°C to 60°C. The distribution ratio of Np(V) between the organic and aqueous phases was found to decrease as the concentrations of fluoride and sulfate were increased. Stability constants of the 1:1 Np(V)-fluoride complexes and the 1:1 Np(V)-sulfate and 1:2 Np(V)-sulfate complexes, dominant in the aqueous phase under the experimental conditions, were calculated from the effect of [F⁻] and [SO₄²⁻] on the distribution ratio. The enthalpy and entropy of complexation were calculated from the stability constants at different temperatures by using the Van't Hoff equation.

Key words: Neptunium (V) / Fluoride / Sulfate/ Complexation / Stability constants / Solvent extraction

INTRODUCTION


Neptunium is an important radioactive component of nuclear wastes disposed of in low-level and high-level waste repositories because of its high toxicity and very long half life (^{237}Np , $t_{1/2} = 2.14 \times 10^6$ years) as well as its high abundance in nuclear waste. The fractional dose attributed to ^{237}Np will reach 67% at 75,000 years.¹ As a result, the migration of neptunium in the postclosure chemical environment of waste repositories is of great concern to long-term repository performance. Depending on the chemical environment in the repositories and the surrounding geologic media, neptunium could exist in multiple oxidation states in nature (e.g., III, IV, V and VI). Np(V) is the most stable oxidation state under environmental and waste conditions. The solubility of neptunium after disposal will depend largely on its reactions (e.g., solubility and complexation) with various components present in the repository. The interactions of actinides with some inorganic ligands at elevated temperatures may be a significant factor in determining the solubility and migration of the actinides in the nuclear waste repository. For example, fluoride and sulfate are expected to be present in the water in contact with the nuclear waste packages and affect the speciation of actinides by forming complexes. The Np(V) complexes with fluoride and sulfate at elevated temperatures may play an important role in the migration

behavior of Np(V). Studies of complexation of Np(V) with various inorganic ligands have recently been reviewed.² Fluoride complexation has been studied in near-neutral solutions by several authors using spectrophotometric,³ solvent extraction,⁴⁻⁶ and potentiometric⁷ methods. But all these data on the stability constants of Np(V) complexes with fluoride were obtained from room temperature. Five experimental studies describing the Np(V) sulfate complexes^{4, 6, 8-10} have been published. Much data have been obtained from studies at room temperature, but only Halperin and Olliver¹⁰ reported their data on the stability constants obtained at temperatures of 5, 25, and 45°C. It is well known that the equilibrium constants of complexation could vary by orders of magnitude in the temperature range (25–80°C) relevant to waste repositories. To minimize uncertainty in the prediction of Np(V) migration, it is necessary to know sufficient and reliable thermodynamic data on its complexes at elevated temperatures. Unfortunately, experimental data on the effects of temperatures in these systems are very scarce at present.

In this work, a solvent extraction method¹¹⁻¹⁴ was used to determine the stability constants of Np(V) complexes with fluoride and sulfate anions in 1.0 M NaClO₄ solutions at the various temperatures ranging from 25°C to 60°C. Di-(2-ethylhexyl) phosphoric acid (HDEHP) was used as a solvent extractant and heptane as a solvent.

The complex reaction of Np(V) with fluoride and sulfate under

experimental conditions used in this study can be expressed as:



with the overall stability constant β_q defined as:

$$\beta_q = (\text{NpO}_2\text{F}_q^{1-q}) / [(\text{NpO}_2^+) (\text{F}^-)^q] \quad (3)$$

$$\beta_q = [\text{NpO}_2(\text{SO}_4)_q^{1-2q}] / [(\text{NpO}_2^+) (\text{SO}_4^{2-})^q] \quad (4)$$

In the solvent extraction system, the distribution ratio (D) is defined as:

$$D = \Sigma (M)_o / \Sigma (M)_a \quad (5)$$

Assuming only a single organic (o) phase species, MA_n , and the various species in the aqueous phase (a), Equation (5) can be written with the stability constants as:¹¹⁻¹²

$$D_o/D = 1 + \Sigma \beta_i(L)^i \quad (6)$$

For the Np(V)-fluoride and Np(V)-sulfate complex systems at a fixed pH (see Equation 7), Np(V) stands for M, fluoride and sulfate anions are L, A is the HDEHP anion, and D_o is the distribution ratio in the absence of complexation.

EXPERIMENTAL

Reagents and Solutions

All reagents used in this work were analytical grade or higher. The stock solutions of 1.0 M and 0.1 M sodium fluoride (Aldrich) and sodium sulfate (Sigma) were prepared by dissolving appropriate amounts of these reagents in deionized water, respectively. The stock solution of fluoride was analyzed by a fluoride combination ion selective electrode (Accumet). Sodium perchlorate (Aldrich) was used for ionic strength adjustment without further purification. Perchloric acid (Aldrich) and NaOH (1.0 N standard, Aldrich) were used to adjust pH. Sodium bromate (Aldrich) and NaNO_2 (Aldrich) were used for oxidizing and reducing neptunium, respectively, during the preparation of the stock solution of NpO_2^+ . Di-(2-ethylhexyl) phosphoric acid, HDEHP (Alfa Aesar), was not further purified, and the stock solutions of 0.1 M and 0.01 M HDEHP in heptane (anhydrous, Sigma-Aldrich) were prepared. The organic solutions were pre-equilibrated with the aqueous stock solution of 1.0 M NaClO_4 and stored in a dark glass bottle. All solutions except Np(V) tracer solution used in the experiments were filtered with a 0.2- μm microfiltration system to minimize the sorption of Np(V) by suspended particles in solution.

An Np(V) solution was prepared using the procedure described in the literature.¹⁵ A small amount of solid NaBrO_3 was added to a solution of neptunium in 1.0 M HClO_4 , and the solution was warmed to facilitate the oxidation of neptunium to Np(VI) . Then a small amount of NaNO_2 was added to this solution, causing a change in color from pink to green

indicating the reduction of NpO_2^{2+} to NpO_2^+ . Finally, the NpO_2^+ species was precipitated by adding 1.0 M NaOH to the solution. The $\text{NpO}_2(\text{OH})$ precipitate was washed several times with deionized water and dissolved with 0.1 M HClO_4 . The oxidation state was confirmed spectrophotometrically by a Cary 500 Scan ultraviolet visible near infrared spectrophotometer and showed that Np was present as NpO_2^+ . A stock solution of Np(V) was prepared from this in a solution of 0.06 N HClO_4 such that 10 μl of the solution gives ca. 15,600 cpm. A Wallac (Model 1414) liquid scintillation counter (LSC) was used to measure ^{237}Np .

Temperature-Controlled Extraction Equipment

To shake the samples at the desired temperature, a reaction block with the heat tape or a block heater was used. The reaction block was strapped onto a Lab-line orbital shaker at an approximately 45° angle to facilitate the mixing of the two phases. The temperature was controlled with the block heater's temperature control or through a variable transformer output unit when using the heat tape.

The reaction blocks are aluminum blocks that have been dimensioned to fit pre-purchased dry block heaters. Each of the blocks has machined holes for the selected sample tubes and a specific hole for a bayonet insert.

Two J-KEM digital temperature monitors for Type T thermocouples

with a range from -200 to 250°C were used with two Type T reaction block thermocouples to monitor the temperature of the reaction blocks and centrifuge during the experiment. The thermocouples have armored cable to withstand constant shaking and a quick-release spring-loaded bayonet style connector. Both of the thermocouples were cross calibrated with each of the monitors.

A Speedfuge, HSX10K centrifuge with a water jacket enclosed around the rotor was used to centrifuge the samples. Water from a Brinkmann RM6 Lauda temperature-controlled water circulator controlled the temperature of the samples inside the centrifuge.

Temperature adjusted reaction blocks were used to maintain the temperature of experimental solutions, samples, and pH buffers.

Determination of H^+ Concentrations

A Corning 355 pH meter was used with a combination of glass electrodes (Ross Orion) to measure the change in pH. Because pH was measured in sodium perchlorate solution, the inner solution (3 M KCl) of the glass electrode was replaced with saturated NaCl solution to avoid erratic electrode performance. The electrode was calibrated with 4.00 ± 0.01 and 7.00 ± 0.01 pH buffer standard solutions at the same temperature as the reaction block. The pH meter readings (pH_m) were converted to a

hydrogen concentration ($\text{pcH} = -\log [\text{H}^+]$) using the following calibration equation proposed by Irving et al.:¹⁶

$$\text{pcH} = \text{pH}_m + b \quad (7)$$

where b is the correction factor of an electrode, depending on the solution composition, such as $b = 0.27$ in 1.0 M NaClO_4 solution.¹¹

Solvent Extraction

The distribution experiments were conducted following the procedure described in the literature.¹¹⁻¹² For each extraction experiment, 2.0 ml of aqueous solution was adjusted to the appropriate pcH having the desired amounts of fluoride/sulfate, and an equal volume of 0.01 M HDEHP in heptane was placed in a borosilicate glass vial. After adding 10 μl $^{237}\text{Np(V)}$ tracer, the vials were inserted into the holes of the temperature-controlled reaction block and shaken for 3 hours at the desired temperature. Kinetic studies confirmed that the extraction equilibrium was attained in less than 1 hour.¹¹ The vials were centrifuged at the same temperature as the reaction block on the shaker, and the aliquots of 0.5 ml were taken from both phases to measure the α -activity in Opti Fluor cocktail by LSC. The remaining aqueous phase was used for pH measurement at the same temperature as the reaction block on the shaker.

RESULTS AND DISCUSSION

Np(V)-Fluoride Complexation

A series of experiments with constant pcH were conducted at an ionic strength of 1.0 M NaClO₄, and distribution ratios (D) of Np(V) in organic and aqueous phases were found to decrease with increased fluoride concentration (for a complete set of the extraction data, see Appendix A Table A1-A4). The decrease in D can be attributed to the increase in concentration of Np(V)-fluoride complexes in the aqueous phase, which are not extracted by HDEHP. Stability constants of the 1:1 Np(V)-fluoride complexes, dominant in the aqueous phase under the experimental conditions, were calculated from the effect of $[F^-]$ on the distribution ratio.

A typical fit curve showing the function of D_o/D versus fluoride anion concentration is shown in Figure 1 in which the points are experimental data and the solid line represents 1:1 complexes that provide the first stability constant. In the pcH range used in this work, the free fluoride anion concentration ($[F^-]_f$) can be considered as the total concentration of fluoride ($[F^-]_T$). The first stability constant (β_1) can be calculated as a slope of the linear regression between D_o/D and $[F^-]_T$ (see Figure 1).

In the extraction experiments, the pH/pcH values measured after extraction sometimes varied slightly. The minor corrections to the D

values for this pcH fluctuation were made by the following relationship:

$$\log D = \log D_{\text{exp}} + n (\text{pcH}_{\text{avg}} - \text{pcH}) \quad (8)$$

where n is the experimental slope of the $\log D_{\text{exp}}$ dependence on pcH in the absence of ligand for the certain NaClO_4 ionic strength solution (1.0 M) at the required temperature (see Appendix A Tables A1-A7). The pcH_{avg} is an average value of pcH measured for each sample.

Literature data indicate that the hydrolysis of Np(V) is insignificant when $\text{pcH} < 7$ in the temperature range of 10–85°C.¹⁷ The experiments in this study were conducted at $\text{pcH} \sim 5$. As a result, the hydrolysis reactions of Np(V) are not included in the calculation.

The apparent stability constants are listed in Table 1 along with the literature values for comparison. The first stability constant of the Np(V) -fluoride complex at room temperature that was obtained from our experiments is consistent with the literature values. Our results show that the stability constants of the 1:1 Np(V) -fluoride complexes were enhanced with increasing temperature.

Np(V) -Sulfate Complexation

Similar to the Np(V) -fluoride system, a series of experiments with constant pcH and temperature were conducted at ionic strength of 1.0 M ($\text{NaClO}_4 + \text{Na}_2\text{SO}_4$). The distribution ratio of Np(V) between the organic

and aqueous phases was found to decrease as the concentration of sulfate was increased (for a complete set of the extraction data, see Appendix A Table A5-A7). Stability constants of the 1:1 Np(V)-sulfate and 1:2 Np(V)-sulfate complexes, dominant in the aqueous phase under the experimental conditions, were calculated from the effect of $[\text{SO}_4^{2-}]$ on the distribution ratio.

A typical fit curve showing function of D_o/D versus free sulfate anion concentration ($[\text{SO}_4^{2-}]_f$) is shown in Figure 2 in which the points are experimental data. The curve is a polynomial fit including both the first stability constant (β_1) and the second stability constant (β_2), and the solid line only represents 1:1 complexes that provide β_1 . The stability constants, β_1 and β_2 , were obtained in the ionic strength of 1.0 M ($\text{NaClO}_4 + \text{Na}_2\text{SO}_4$) by analyzing such curves. The minor corrections to the D values for this pH fluctuation were made using the same method as the system of Np(V)-fluoride. It should be noted that the sulfate was completely dissociated to SO_4^{2-} at $\text{pH} > 3$; thus, in the pH range used in this work, the free sulfate anion concentration $[\text{SO}_4^{2-}]_f$ can be replaced by the total concentration of sulfate (see Figure 2).

The first apparent stability constants of Np(V) complexes with sulfate are listed in Table 2, including the literature values for comparison. Our measured stability constants for the 1:1 Np(V)-sulfate complex do not compare well with any literature value, but our value does fall within the

overall spread of these literature values. Halperin and Oliver's results¹⁰ indicated that in sulfate concentrations of up to 0.3 M, there is no evidence for Np(V) complexes higher than the monosulfate. The results from our experiments show that there is a formation of 1: 2 complexes of Np(V)-sulfate complex at sulfate concentrations lower than 0.3 M (see Figure 2). The stability constant for 1:2 Np(V)-sulfate complex was calculated to be 1.60 ± 0.11 at 40° C and 1.82 ± 0.07 at 50° C. The sulfate complexation of Np(V) is relatively weak, but increases with increasing temperature.

Effect of Temperature on Stability Constants

The effect of temperature on the stability constants was evaluated at the range of temperature from 25°C to 60°C for Np(V)-F⁻ complexes and from 25°C to 50°C for Np(V)-SO₄²⁻ complexes, respectively. Halperin and Oliver¹⁰ have pointed out that although the thermodynamic values of free energy, enthalpy, and entropy cannot be evaluated from the concentration dependent stability constants, the effective values for these thermodynamic quantities can be calculated by using Van't Hoff equation. To distinguish these calculated values from the true thermodynamic values, they will be hereafter referred to as prime free energy ($\Delta G^{\circ'}$), prime enthalpy ($\Delta H^{\circ'}$), and prime entropy ($\Delta S^{\circ'}$), respectively. From the determined stability constants listed in Tables 1 and 2, the corresponding

values of $\Delta G^{o'}$ defined according to Equation (9) for each value at a fixed temperature were calculated:

$$\Delta G^{o'} = -RT \ln \beta \quad (9)$$

The values of $\Delta H^{o'}$ and $\Delta S^{o'}$ at 25°C were calculated by using the Van't Hoff equation:

$$\partial \ln \beta / \partial (1/T) = -\Delta H^{o'} / R ; \quad \Delta S^{o'} = (\Delta H^{o'} - \Delta G^{o'}) / T \quad (10)$$

These calculated values are listed in Table 3. The small values of $\Delta G^{o'}$ reflect the low values of the stability constant. The Np(V)-fluoride complexes are stronger than those with Np(V)-sulfate complexes. The results indicate that the values of $\Delta H^{o'}$ and $\Delta S^{o'}$ are positive and relatively large. NpO_2^+ should be classified as a hard acceptor¹⁰ using the nomenclature and criteria of Person and Ahrland¹⁸⁻²⁰ whereas F^- and SO_4^{2-} have been classified as hard donors.²⁰ Ahrland²⁰ has pointed out that inner complexes are generally entropy controlled. The gain in enthalpy supports the interpretation of inner sphere binding for the Np(V) complexes with fluoride and sulfate.¹⁰

CONCLUSION

A solvent extraction method was used to determine the stability constants of the Np(V)-fluoride and -sulfate complexes in 1.0 M ionic strength at various temperatures. The 1:1 Np(V)-fluoride complex was

found to be dominant in the aqueous phase under the experimental conditions, and the values of the stability constants increased with increasing temperature. The sulfate complexes of NpO_2^+ are relatively weak, and their respective stability constants are relatively small, but they show an enhancement with increasing temperature. The values of enthalpy and entropy of complexation calculated by Equation (10) are relatively large positive values that are consistent with the inner sphere complexing.^{10, 20} The 1:2 Np(V)-sulfate complex was found to form under the experimental conditions (sulfate concentrations from 0.03 to 0.21 M). The values obtained in these experiments show that both Np(V)-fluoride and sulfate complexes may be important in the migration behavior of Np(V) in nuclear waste given elevated surface temperatures and high fluoride/sulfate concentrations.

ACKNOWLEDGEMENTS

This work was supported by the Office of Civilian Radioactive Waste Management (OCRWM), U.S. Department of Energy. Pacific Northwest National Laboratory is operated for the U.S. Department of Energy by Battelle under Contract DE-AC05-76RL01830. We thank Andrea D. Choiniere for reviewing this document.

REFERENCES

1. Office of Civilian Radioactive Waste Management (OCRWM).
"Yucca Mountain Science and Engineering Report Rev. I,"
DOE/RW-0539-1, North Las Vegas, NV (2002).
2. J. Fuger, H. Nitsche, P. Potter, M. H. Rand, J. Rydberg, K. Spahiu,
J. C. Sullivan, W. J. Ullman, P. Vitorge, and H. Wanner. Chemical
Thermodynamics of Neptunium and Plutonium. Edited by The
Organization for Economic Cooperation and Development, Nuclear
Energy Agency, Data Bank, Chemical Thermodynamics 4, Elsevier,
NY (2001).
3. P. R. V. Rao and S. K. Patil. *Radiochim. Radioanal. Lett.*, 36 (1978)
169.
4. P. R. V. Rao, N. M. Gudi, S. V. Bagawde, and S. K. Patil. *J. Inorg.
Nucl. Chem.*, 41 (1979) 235.
5. G. R. Choppin and L. F. Rao. *Acta*, 37 (1984) 143.
6. Y. Inoue and O. Tochiyama. Studies of the formation of complexes of
Np(V) with inorganic ligands by means of solvent extraction with 2-
thenoyltrifluoroacetone and 1-10-phenanthroline. II. fluoro, sulfato,
and phosphate complexes, *Bull. Chem. Soc. Jpn.* 58, 2228-2233
(1985).
7. R. M. Sawant, G. H. Rizvi, N. K. Chaudhuri, and S. K. Patil. *J.*

- Radioanal. Nucl. Chem., 89 (1985) 373.
8. A. I. Moskvina and A. N. Poznyakov. Russ. J. Inorg. Chem., 24 (1979) 1357.
 9. S. K. Patil, V. V. Ramakrishna, and N. M. Gudi. Studies on complexing of some actinide ions in aqueous solutions. In: "Proc. Nucl. Chem. Radiochem. Symp. 1980," Indian Department of Atomic Energy, pp 388-390 (1980).
 10. J. Halperin and J. H. Oliver. Radiochem. Acta, 33 (1983) 29.
 11. G. R. Choppin, H. N. Erten, and Y. Xia. Radiochim. Acta, 74 (1996) 123.
 12. Y. Xia, L. Rao, D. Rai, and A. R. Felmy. Radiochim. Acta, 86 (1999) 33.
 13. M. Borkowski, R. C. Moore, M. G. Bronikowski, J. Chen, O. S. Pokrovsky, Y. Xia, and G. R. Choppin. J. Radioanalytical and Nuclear Chemistry, 248 (2001) 467.
 14. G. R. Choppin, A. H. Bond, M. Borkowski, M. G. Bronikowski, J. Chen, S. Lis, J. Mizera, O. Pokrovsky, N. A. Wall, Y. Xia, and R. C. Moore. Waste Isolation Pilot Plant Actinide Source Term Test Program: Solubility Studies and Development of Modeling Parameters. Sandia National Laboratories Technical Report, SAND99-0943, Albuquerque, NM (2001).
 15. L. Rao and G.R. Choppin. Inorg. Chem., 23 (1984) 2351.

16. H. M. Irving, M. G. Miles, and L. D. Pettit. *Anal. Chem. Acta*, 38 (1967) 475.
17. L. Rao, T. G. Srinivasan, A. Yu. Garnov, P. Zanonato, P. Di Bernardo, and A. Bismondo, *Geochim. Cosmochim. Acta*, 68 (2004) 4821.
18. R. G. Pearson. *J. Chem. Ed.*, 45 (1968) 581.
19. R. G. Pearson. *J. Chem. Ed.*, 45 (1968) 643.
20. S. Ahrland. *Coord. Chem. Rev.*, 8 (1972) 21.

APPENDIX A: SOLVENT EXTRACTION DATA TABLES

Table A1. Solvent extraction data for the system of
Np(V)-F⁻-1.0 M NaClO₄-0.01 M HDEHP/
heptane at 25±1°C

pcH	[F ⁻] _T (M)	Activity (cpm)		D ^(a)
		Org. Phase	Aq. Phase	
4.78	0	2868	929	6.59
4.90	5.00e-03	2976	860	5.23
4.94	7.50e-03	3005	839	4.84
4.98	1.25e-02	3014	786	4.57
5.07	2.00e-02	3044	737	3.81
5.10	2.25e-02	3080	694	3.74
5.13	2.50e-02	3070	722	3.30
5.17	2.75e-02	3138	690	3.17
5.17	3.00e-02	3083	706	3.03
5.18	3.25e-02	3083	743	2.79

(a) The minor corrections to the D values for the pcH fluctuation were made by the relationship:

$$\log D = \log D_{\text{exp}} + n (\text{pcH}_{\text{avg}} - \text{pcH})$$

where n (1.25) is the experimental slope of the log D_{exp} dependence on pcH in the absence of ligand for the certain NaClO₄ ionic strength solution.

The activity of background: 1 cpm.

Table A2. Solvent extraction data for the system of
 $\text{Np(V)}\text{-F}^-$ -1.0 M NaClO_4 -0.01 M HDEHP/
 heptane at $35\pm 1^\circ\text{C}$

pcH	$[\text{F}^-]_{\text{T}}$ (M)	Activity (cpm) Org. Phase	Activity (cpm) Aq. Phase	$D^{(a)}$
4.90	0	3223	577	10.96
5.05	5.00e-03	3239	507	7.99
5.08	7.50e-03	3177	486	7.67
5.12	1.25e-02	3245	515	6.43
5.13	2.00e-02	3037	535	5.62
5.24	2.75e-02	3185	497	4.72
5.28	3.00e-02	3205	429	4.88
5.29	3.25e-02	3181	462	4.33

(a) The minor corrections to the D values for the pcH fluctuation were made by the relationship:

$$\log D = \log D_{\text{exp}} + n(\text{pcH}_{\text{avg}} - \text{pcH})$$

where n (1.25) is the experimental slope of the log D_{exp} dependence on pcH in the absence of ligand for the certain NaClO_4 ionic strength solution.

The activity of background: 1 cpm.

Table A3. Solvent extraction data for the system of
Np(V)- F⁻ -1.0 M NaClO₄ -0.01 M HDEHP/
heptane at 50±1°C

pcH	[F ⁻] _T (M)	Activity (cpm)		D ^(a)
		Org. Phase	Aq. Phase	
4.96	0	3025	595	8.48
5.03	5.00e-03	2946	541	7.21
5.08	7.50e-03	3026	546	6.41
5.11	1.00e-02	3004	566	5.47
5.17	1.50e-02	3024	510	5.12
5.22	2.00e-02	3046	516	4.24
5.27	2.50e-02	3092	573	

(a) The minor corrections to the D values for the pcH fluctuation were made by the relationship:

$$\log D = \log D_{\text{exp}} + n (\text{pcH}_{\text{avg}} - \text{pcH})$$

where n (1.41) is the experimental slope of the log D_{exp} dependence on pcH in the absence of ligand for the certain NaClO₄ ionic strength solution.

The activity of background: 1 cpm.

Table A4. Solvent extraction data for the system of
Np(V)- F⁻ -1.0 M NaClO₄ -0.01 M HDEHP/
heptane at 60±1°C

pcH	[F ⁻] _T (M)	Activity (cpm) Org. Phase	Activity (cpm) Aq. Phase	D ^(a)
5.16	0	3077	348	19.92
5.25	5.00e-03	3164	300	17.26
5.28	7.50e-03	3160	304	15.63
5.32	1.25e-02	3125	309	13.37
5.37	1.50e-02	3100	319	10.85
5.49	2.00e-02	3144	245	9.77
5.63	2.50e-02	3080	179	8.34
5.74	2.75e-02	3086	149	7.14

(a) The minor corrections to the D values for the pcH fluctuation were made by the relationship:

$$\log D = \log D_{\text{exp}} + n (\text{pcH}_{\text{avg}} - \text{pcH})$$

where n (1.41) is the experimental slope of the log D_{exp} dependence on pcH in the absence of ligand for the certain NaClO₄ ionic strength solution.

The activity of background: 1 cpm.

Table A5. Solvent extraction data for the system of
 $\text{Np(V)}\text{-SO}_4^{2-}$ -1.0 M NaClO_4 -0.01 M HDEHP/
 heptane at $25\pm 1^\circ\text{C}$

pcH	$[\text{SO}_4^{2-}]_{\text{T}}$ (M)	Activity (cpm)	Activity (cpm)	$D^{(a)}$
		Org. Phase	Aq. Phase	
5.02	0	2675	1193	3.07
5.02	0.03	2537	1342	2.64
5.07	0.06	2408	1476	1.95
5.08	0.09	2234	1635	1.59
5.09	0.12	2167	1734	1.42
5.21	0.15	2053	1822	0.93
5.22	0.18	1860	1987	0.74
5.24	0.21	1767	2137	0.63

(a) The minor corrections to the D values for the pcH fluctuation were made by the relationship:

$$\log D = \log D_{\text{exp}} + n (\text{pcH}_{\text{avg}} - \text{pcH})$$

where n (1.20) is the experimental slope of the log D_{exp} dependence on pcH in the absence of ligand for the certain NaClO_4 ionic strength solution.

The activity of background: 1 cpm.

Table A6. Solvent extraction data for the system of
 Np(V)-SO₄²⁻-1.0 M NaClO₄-0.01 M HDEHP/
 heptane at 40±1°C

pcH	[SO ₄ ²⁻] _T (M)	Activity (cpm)		D ^(a)
		Org. Phase	Aq. Phase	
5.21	0	3103	671	6.21
5.23	0.03	3027	689	5.53
5.26	0.06	2995	744	4.74
5.31	0.09	2917	826	3.73
5.36	0.12	2845	892	2.95
5.39	0.15	2806	952	2.55
5.42	0.18	2666	1152	1.88
5.46	0.21	2620	1101	1.76

(a) The minor corrections to the D values for the pcH fluctuation were made by the relationship:

$$\log D = \log D_{\text{exp}} + n(\text{pcH}_{\text{avg}} - \text{pcH})$$

where n (1.04) is the experimental slope of the log D_{exp} dependence on pcH in the absence of ligand for the certain NaClO₄ ionic strength solution.

The activity of background: 1 cpm.

Table A7. Solvent extraction data for the system of
Np(V)-SO₄²⁻-1.0 M NaClO₄ -0.01 M HDEHP/
heptane at 50±1°C

pcH	[SO ₄ ²⁻] _T (M)	Activity (cpm)		D ^(a)
		Org. Phase	Aq. Phase	
5.13	0	2946	637	7.47
5.27	0.03	2875	674	5.38
5.36	0.06	2871	690	4.48
5.37	0.09	2715	808	3.56
5.37	0.12	2589	903	3.05
5.53	0.15	2571	964	2.14
5.57	0.18	2388	994	1.79
5.62	0.21	2450	1139	1.47

(a) The minor corrections to the D values for the pcH fluctuation were made by the relationship:

$$\log D = \log D_{\text{exp}} + n (\text{pcH}_{\text{avg}} - \text{pcH})$$

where n (0.77) is the experimental slope of the log D_{exp} dependence on pcH in the absence of ligand for the certain NaClO₄ ionic strength solution.

The activity of background: 1 cpm.

Figure captions

Figure 1. Plot of D_o/D vs. concentration of fluoride in the extraction system of Np(V)-F-1.0 M NaClO_4 -0.01 M HDEHP/heptane at 35°C, points are experimental data, solid line is calculated.

Figure 2. Plot of D_o/D vs. concentration of sulfate in the extraction system of Np(V)-sulfate-1.0 M NaClO_4 -0.01 M HDEHP/heptane at 40°C.




Figure.1




Figure 2

Table 1. Apparent stability constants of Np(V) complexes
with fluoride ($\text{NpO}_2^+ + \text{F}^- = \text{NpO}_2\text{F}$)

Method	Ionic medium	t (°C)	$\log\beta_{101}$	Reference
sp	2 M NaClO ₄	25	1.35 ± 0.30	[3]
dis	2 M NaClO ₄	25	0.99 ± 0.10	[4]
dis	1 M NaClO ₄	23	1.26 ± 0.30	[5]
dis	1 M NaClO ₄	25	1.39 ± 0.30	[6]
ise-F	0.1 M NaClO ₄	21	1.51 ± 0.50	[7]
dis	1 M NaClO ₄	25±1	1.42 ± 0.10	This work
dis	1 M NaClO ₄	35±1	1.63 ± 0.03	This work
dis	1 M NaClO ₄	50±1	1.77 ± 0.04	This work
dis	1 M NaClO ₄	60±1	1.80 ± 0.03	This work

Note: sp—spectrophotometric
dis—solvent extraction
ise-F—potentionmetric

Table 2. Apparent stability constants of Np(V) complexes
with sulfate ($\text{NpO}_2^+ + \text{SO}_4^{2-} = \text{NpO}_2\text{SO}_4^-$)

Method	Ionic medium	t (°C)	$\log \beta_{101}$	Reference
cop	1.5 M NH_4Cl	20 ± 2	1.04 ± 0.4	[8]
dis	2 M NaClO_4	25	0.44 ± 0.09	[4]
dis	8.5 M NaClO_4	25	0.45	[9]
dis	0.52 m NaClO_4	5	$-(0.43 \pm 0.35)$	[10]
dis	1.05 m NaClO_4	5	$-(0.18 \pm 0.09)$	[10]
dis	2.2 m NaClO_4	5	$-(0.04 \pm 0.11)$	[10]
dis	0.52 m NaClO_4	25	$-(0.10 \pm 0.08)$	[10]
dis	1.05 m NaClO_4	25	0.06 ± 0.04	[10]
dis	2.2 m NaClO_4	25	0.19 ± 0.04	[10]
dis	0.52 m NaClO_4	45	0.25 ± 0.03	[10]
dis	1.05 m NaClO_4	45	0.34 ± 0.03	[10]
dis	2.2 m NaClO_4	45	0.40 ± 0.05	[10]
dis	1 M NaClO_4	25	0.76 ± 0.04	[6]
sv*	0 m	25	0.44 ± 0.27	[2]
dis	1 M NaClO_4	25 ± 1	0.49 ± 0.31	This work
dis	1 M NaClO_4	40 ± 1	0.65 ± 0.19	This work
dis	1 M NaClO_4	50 ± 1	0.73 ± 0.16	This work

Note: cop—coprecipitation; dis—solvent extraction

* Selective value—the recommended value from editors of
Chemical Thermodynamics of Neptunium and Plutonium
(2001) [2].

Table 3. Stability constants and values of ΔG° , ΔH°
and ΔS° for NpO_2F^- and $\text{NpO}_2\text{SO}_4^-$

	I		ΔG°	ΔH°	ΔS°	
C°	(M)	$\log \beta_1$	(kJ /mol)	(kJ /mol)	(J/[k mol])	Reference
For the complex formation of NpO_2F^-						
25	1.0	1.42 ± 0.10	-8.11	20.8	96.9	This work
For the complex formation of $\text{NpO}_2\text{SO}_4^-$						
25	1.0	0.49 ± 0.31	-2.80	17.8	69.2	This work
25	1.0	0.06 ± 0.04	-0.37	22.2	76.0	[10]