



## ERNEST ORLANDO LAWRENCE BERKELEY NATIONAL LABORATORY


Title: Endosymbiosis In Statu Nascendi:  
Close Phylogenetic Relationship  
Between Obligately Endosymbiotic and  
Obligately Free-Living *Polynucleobacter*  
*Strains (Betaproteobacteria)*

Author(s): Claudia Vannini, Matthias Pockl, Guilio  
Petroni, Qinglong L. Wu, Elke Lang, Erko  
Stackebrandt, Martina Schrallhammer, Paul M.  
Richardson, and Martin W. Hahn

Division: Genomics

Submitted to: Environmental Microbiology

Month Year: 7/06



**Endosymbiosis In Statu Nascendi: Close Phylogenetic Relationship**

**Between Obligately Endosymbiotic and Obligately Free-Living**

***Polynucleobacter* Strains (*Betaproteobacteria*)**

Claudia Vannini<sup>1</sup>, Matthias Pöckl<sup>2</sup>, Giulio Petroni<sup>1</sup>, Qinglong L. Wu<sup>2,3,\*</sup>, Elke Lang<sup>4</sup>, Erko  
Stackebrandt<sup>4</sup>, Martina Schrällhammer<sup>5</sup>, Paul M. Richardson<sup>6</sup>, and Martin W. Hahn<sup>2, §</sup>

<sup>1</sup> Department of Biology – Protistology and Zoology Unit, University of Pisa, Via A. Volta  
4/6, I-56126 Pisa, Italy

<sup>2</sup> Institute for Limnology, Austrian Academy of Sciences, Mondseestrasse 9, A-5310  
Mondsee, Austria

<sup>3</sup> Nanjing Institute of Geography and Limnology, Chinese Academy of Sciences, 73 East  
Beijing Road, Nanjing 210008, P.R.China

<sup>4</sup> DSMZ GmbH - German Collection of Microorganisms and Cell Cultures, Mascheroder  
Weg 1b, D-38124 Braunschweig, Germany

<sup>5</sup> Department of Zoology, University of Stuttgart, D-70550 Stuttgart, Germany

<sup>6</sup> Department of Energy, Joint Genome Institute, 2800 Mitchell Drive, Walnut Creek, CA,  
USA

\* Present address: Division of Environmental Science & Engineering, National University of  
Singapore, 9 Engineering Drive 1, EA-07-23, Singapore 117576

§ Corresponding author, Phone: +43 6232 3125-29, Fax: +43 6232 3578, e-mail:

martin.hahn@oeaw.ac.at

Running title: Complete niche separation in *Polynucleobacter* strains

26 **Summary**

28 Bacterial strains affiliated to the phylogenetically shallow subcluster C (PnecC) of the  
30 *Polynucleobacter* cluster, which is characterized by a minimal 16S rRNA gene sequence  
32 similarity of approx. 98.5%, have been reported to occur as obligate endosymbionts of  
34 ciliates (*Euplotes* spp.), as well as to occur as free-living cells in the pelagic zone of  
40 freshwater habitats. We investigated if these two groups of closely related bacteria represent  
36 strains fundamentally differing in lifestyle, or if they simply represent different stages of a  
42 facultative endosymbiotic lifestyle. The phylogenetic analysis of 16S rRNA gene and 16S-  
44 23S ITS sequences of five endosymbiont strains from two different *Euplotes* species and 40  
pure culture strains demonstrated host-species-specific clustering of the endosymbiont  
sequences within the PnecC subcluster. The sequences of the endosymbionts showed  
characteristics indicating an obligate endosymbiotic lifestyle. Cultivation experiments  
revealed fundamental differences in physiological adaptations, and determination of the  
genome sizes indicated a slight size reduction in endosymbiotic strains. We conclude that the  
two groups of PnecC bacteria represent obligately free-living and obligately endosymbiotic  
strains, respectively, and do not represent different stages of the same complex lifecycle.  
These closely related strains occupy completely separated ecological niches. To our best  
knowledge, this is the closest phylogenetic relationship between obligate endosymbionts and  
obligately free-living bacteria ever revealed.

**Introduction.** In 1975, Heckmann described a bacterial endosymbiont of the freshwater  
46 ciliate *Euplotes aediculatus* (Hypotrichia) and named it “omikron” (Heckmann, 1975). These  
bacterial endosymbionts were found to be individually contained in cytoplasmic vesicles, to  
48 which ribosomes are often attached. The omikron endosymbiont is characterized by the  
presence of usually 3-9 nucleoid-like structures (Heckmann and Schmidt, 1987). So far,  
50 omikron-like endosymbionts were reported from nine *Euplotes* species, all but one of these  
species inhabit freshwater systems (Görtz, 2002; Petroni et al, 2002).

52 Heckmann demonstrated that *Euplotes aediculatus* cured from their endosymbionts by  
antibiotic treatment were able to perform 1-2 more cell divisions but died 15-20 days after the  
54 last division (Heckmann, 1975; Heckmann and Schmidt, 1987). Treatment of aposymbiotic  
ciliates with ciliate homogenate containing intact omikron-like endosymbionts or by direct  
56 injection of symbiont-containing cytoplasm resulted in both, the reestablishment of the  
endosymbiont population, and the recovery of the temporarily aposymbiotic ciliates  
58 (Heckmann, 1975; Fujishima and Heckmann, 1984). Repeated attempts to cultivate the  
endosymbionts in pure culture failed completely (Heckmann, 1975). Based on these  
60 observations it was concluded that omikron is an obligately endosymbiotic bacterium, which  
is essential for its ciliate host (Heckmann, 1975; Heckmann et al., 1983; Heckmann and  
62 Schmidt, 1987). Meanwhile, the dependence of at least eight *Euplotes* species on omikron-  
like endosymbionts was demonstrated (Heckmann et al., 1983; Vannini et al., 2005).

64 In 1987, Heckmann & Schmidt described the omikron endosymbiont of *Euplotes*  
*aediculatus* as *Polynucleobacter necessarius* (Heckmann and Schmidt, 1987). This species is  
66 one of a few validly described bacterial species lacking a pure culture of its type strain. In  
1996, Springer and colleagues determined the 16S rRNA gene sequence of the type strain,  
68 and verified the affiliation of the sequence to the endosymbiont by fluorescence in situ  
hybridization (FISH) with a *P. necessarius*-specific probe (Springer et al., 1996).

70 From 1996 on, cloning and sequencing of PCR-amplified fragments of bacterial 16S  
rRNA genes from surface freshwater habitats yielded more than 100 sequences related to the  
72 sequence of the endosymbiont *P. necessarius* (e.g., Bahr et al., 1996; Hiorns et al., 1997;  
Semenova and Kuznedelov, 1998, Simek et al., 2001). The monophyletic cluster formed by  
74 these environmental sequences and the *P. necessarius* sequence was first designated ACK-1  
cluster (Hiorns et al., 1997), and later, after the discovery of the affiliation of a valid species  
76 with this cluster, it was renamed as *Polynucleobacter necessarius* cluster (Zwart et al., 2002).  
With one exception (Pedersen et al., 1996) the more than 100 retrieved environmental  
78 sequences affiliated with the *P. necessarius* cluster were obtained from surface freshwater  
habitats. The *Polynucleobacter* cluster contains four monophyletic, species-like (minimal 16S  
80 rRNA gene sequence similarity >97%) subclusters (Hahn, 2003). The 16S rRNA sequences  
of five previously investigated endosymbionts of the two ciliate species *Euplotes aediculatus*  
82 (Springer et al., 1996; Vannini, et al., 2005) and *E. harpa* (Vannini, et al., 2005) are affiliated  
with subcluster C (PnecC) of the *Polynucleobacter* cluster. This subcluster also contains  
84 cultivated strains isolated by the filtration acclimatization method (Hahn et al., 2004) from  
the pelagic zone of diverse freshwater habitats (Hahn, 2003). The highest 16S rRNA gene  
86 sequence similarity between a PnecC strain isolated from freshwater and the *P. necessarius*  
type strain is 99.5%.

88 Recently, it was demonstrated by fluorescence in situ hybridization (FISH) with  
subcluster-specific probes that strains affiliated to subcluster PnecC or other  
90 *Polynucleobacter* subclusters occur with high abundances (up to  $10^6$  cells ml<sup>-1</sup>) as free-living  
cells in the plankton of freshwater habitats (Hahn et al., 2005; Wu and Hahn, 2006a; Wu and  
92 Hahn, 2006b). Furthermore, it was demonstrated that every ribotype (16S-23S ITS genotype)  
detected by a cultivation-independent method in a particular habitat could be cultivated by  
94 the acclimatization method (Hahn et al., 2005). On the other hand, all previous attempts to

obtain pure cultures of the *P. necessarius* endosymbionts of *E. aediculatus* by standard  
96 microbiological methods failed (Heckmann, 1975).

At first glance, it seems that the strains affiliated to the species-like subcluster PnecC fall  
98 into two groups, i.e., a group of free-living strains, and a group of endosymbiotic strains.  
However, it is not known if these two groups differ fundamentally in lifestyle, i.e., if the one  
100 group represents obligately free-living and the other group represents obligately  
endosymbiotic strains, or if both groups simply represent different stages of a facultative  
102 endosymbiotic lifestyle. Previously, it was suggested that the isolated strains and the  
endosymbionts differ completely in lifestyle (Hahn, 2003). This suggestion was based on  
104 morphological differences between the two groups of bacteria (differences in cell shape and  
size, and lack of the 3-9 nucleoid-like structures in strains isolated from freshwater).  
106 Meanwhile, pure cultures of new PnecC strains were established, which share with the  
endosymbiotic stages some of their characteristic morphological features (Hahn, unpublished  
108 data), however, the presence of the common traits (i.e., elongated cells and nucleoid-like  
structures) was always restricted to minorities of the cells in the pure cultures. This observed  
110 morphological plasticity of some cultivated PnecC strains does not support the previous  
conclusions on differences in lifestyles of free-living and endosymbiotic stages.

112 In the study presented here, we investigated if the endosymbiotic and the free-living stages  
of PnecC bacteria really differ fundamentally in lifestyle. We examined if (i) free-living and  
114 endosymbiotic stages differ in culturability, (ii) compared the 16S rRNA gene and ITS  
sequence of 40 cultivated PnecC strains and 5 endosymbiotic strains, (iii) searched for  
116 endosymbiotic genotypes in the pelagic of freshwater habitats with specific fluorescence in  
situ hybridization (FISH) probes, (iv) searched for potential genome reduction in  
118 endosymbiotic strains by comparison of genome sizes of endosymbiotic and free-living  
stages, and (v) investigated if the free-living and the endosymbiotic stages differ in the

120 presence of deleterious mutations in their ribosomal sequences. The obtained results support  
the hypothesis of fundamental differences in the lifestyles of free-living and endosymbiotic  
122 PnecC bacteria. Thus, the species-like subcluster PnecC contains closely related obligate  
endosymbionts and obligate free-living strains. Consequently, the subcluster PnecC contains  
124 at least three groups of strains (two groups of host-specific endosymbionts, and the free-  
living strains) occupying completely separated ecological niches. To our best knowledge, the  
126 PnecC bacteria represent the closest phylogenetic relationship between obligate  
endosymbionts and obligately free-living bacteria ever revealed, and the *Polynucleobacter-  
128 Euplotes* system most likely represents the evolutionary youngest obligate endosymbiosis  
discovered so far.

130

## Results

132 **Analysis of ribosomal sequence.** The 16S rRNA gene sequences and 16S-23S ITS  
sequences of 45 strains affiliated with the species-like subcluster PnecC of the  
134 *Polynucleobacter* cluster were analyzed. The 45 strains included 40 pure culture strains  
isolated from freshwater habitats (subsequently termed free-living strains), and five  
136 endosymbiotic strains hosted by the ciliates *Euplotes aediculatus* (two strains) and *E. harpa*  
(three strains) (Table 1). Phylogenetic analysis as well as analysis by BLAST indicated that  
138 the set of 16S rRNA sequences of the 40 pure culture strains included those *Polynucleobacter*  
sequences closest related to the endosymbiont sequences, i.e. public data bases did not  
140 contain any environmental *Polynucleobacter* sequence more similar to the endosymbiont  
sequences as the analyzed sequences. The aligned sequence set consisted of 2022 positions  
142 (1505 positions belonged to the partial 16S rRNA gene sequences, and 517 positions to the  
ITS sequences). The 16S rRNA gene sequence newly obtained from the type strain of  
144 *Polynucleobacter necessarius* (harbored by *E. aediculatus* strain ATCC 30859) differed in 4

positions from the previously determined sequence (acc. number X93019, (Springer et al.,  
146 1996)). None of the 4 positions in the new sequence were in conflict with the secondary  
structure model of the 16S rRNA provided by the ARB software package, therefore, further  
148 analysis was based on the new sequence.

The 16S rRNA gene and ITS sequences of the two endosymbionts of *E. aediculatus*  
150 strains ATCC 30859 and STIR1, which were obtained from sites located more than 400 km  
apart from each other (Fig. 1), were identical over the entire length of 2002 nucleotides (0  
152 polymorphic sites). The three sequences obtained from the endosymbionts of the three *E.*  
*harpa* strains (isolated from habitats located up to 3000 km apart) possessed one polymorphic  
154 position in the 16S rRNA gene, and three polymorphic positions in the ITS. In contrast,  
sequences of the 40 free-living *PnecC* strains contained 106 polymorphic sites, and sequences  
156 of strains obtained from the same habitat possessed 2 to 60 polymorphic sites (9 habitats; 2 to  
6 isolates per habitat; in average 25.2 polymorphic sites in sequences of strains from the same  
158 habitat). The highest number of polymorphic sites in sympatric strains was found in six  
isolates obtained from Lake Mekkojärvi (Finland).

160 Phylogenetic trees based on 16S rRNA gene or ITS sequences, respectively, of the 45  
*PnecC* strains and several strains affiliated with other *Polynucleobacter* subclusters were  
162 calculated by three methods. In all trees the endosymbiont sequences clustered within the  
radiation of subcluster *PnecC* (Fig. 2 and 3). The endosymbiont sequences obtained from *E.*  
164 *aediculatus* and *E. harpa* hosts never appeared in a common cluster separated from all  
sequences of free-living strains, but appeared as two separate host-specific clusters within the  
166 *PnecC* subcluster. The sequences of the two host-specific groups of endosymbionts showed  
for both phylogenetic markers the highest divergence from other members of this subcluster  
168 (Fig. 2 and 3). Besides the higher number of mutations in all five ribosomal sequences of  
endosymbionts, the three *E. harpa* endosymbionts possessed in comparison to other members



170 of the *Polynucleobacter* cluster a significantly higher number of base pair mismatches in  
helices (according to the secondary structure model provided by the software package ARB)  
172 (Fig. 4). Interestingly, this trait is not shared by the two endosymbionts of *E. aediculatus*.

The 16S rRNA gene sequences of the endosymbiotic strains possessed host species-  
174 specific signatures. For instance, both endosymbionts of *E. aediculatus* contain an insertion  
of a guanine nucleotide between positions 193 and 194 (*E. coli* numbering), and all three  
176 endosymbionts of *E. harpa* possessed the same signature of seven single-nucleotide  
substitutions in stem regions, resulting in mismatches in base-pairing. These characteristics  
178 were neither found in any of the >100 *Polynucleobacter* strains so far isolated from  
freshwater habitats, nor in one of the >100 environmental *Polynucleobacter* sequences  
180 deposited in the public databases.

The 16S rRNA gene sequences of the endosymbiotic *Polynucleobacter* bacteria and the  
182 free-living *Polynucleobacter* strains differ only in a few positions. Such small sequence  
dissimilarities have so far not been reported for obligate endosymbionts and their closest free-  
184 living relatives (Table 2).

186 **Genome size and G+C content of *Polynucleobacter* bacteria.** The genome size of three  
endosymbiotic and eleven free-living *Polynucleobacter* strains were determined by pulsed  
188 field gel electrophoresis (PFGE). The endosymbiotic strains possessed genome sizes of 1.7  
and 1.8 Mb, while the eight free-living strains affiliated with subcluster PnecC possessed  
190 significantly larger genome sizes (t-test,  $P < 0.001$ ) ranging from 2.1 to 2.5 Mb (Fig. 5).

Genome sequencing of the freshwater isolate QLW-P1DMWA-1 confirmed the genome size  
192 determined for this strain by PFGE (2.2 Mb in both cases). On the other hand, the three  
investigated free-living strains affiliated with subcluster PnecD possessed genome sizes,  
194 which did not differ significantly from the genome size of the endosymbiotic PnecC strains

(t-test,  $P > 0.1$ ). Though the genus *Polynucleobacter* is affiliated with the family  
196 *Burkholderiaceae* (*Burkholderiales*, *Betaproteobacteria*), *Polynucleobacter* strains studied  
here differ significantly in genome size (Fig. 5B) and G+C content (Fig. 6) from all other  
198 *Burkholderiaceae* strains investigated so far.

200 **Cultivation experiments with two endosymbiotic strains.** Attempts to cultivate two  
endosymbiotic *Polynucleobacter* strains by using the filtration-acclimatization and the  
202 dilution-acclimatization methods did not result in pure cultures of the strains. In the case of  
the filtration-acclimatization method, the failure of the experiment was most likely a result of  
204 the inability of the endosymbiotic strains to pass through 0.2  $\mu\text{m}$  filters. Microscopical  
investigations of 0.2  $\mu\text{m}$ -filtrates of the endosymbiont-containing ciliate homogenate did not  
206 result in the detection of endosymbionts. Nevertheless, samples were further processed by the  
acclimatization method, but no *Polynucleobacter* cultures were finally obtained.

208 In the experiments performed with the dilution-acclimatization method, the presence of  
intact endosymbiont cells in the homogenates used as inoculum was verified microscopically.  
210 Diluted and undiluted homogenates were acclimatized to higher substrate concentrations, and  
finally screened for the presence of PnecC bacteria by FISH. In order to increase the chances  
212 of successful cultivation, different initial media were used in the acclimatization processes. In  
some experiments, acclimatization to higher substrate concentrations was substituted by  
214 cultivation in sterile ciliate homogenate instead. None of these cultivation experiments did  
result in any cultures of *Polynucleobacter* strains. Several established cultures, however,  
216 showed growth of non-*Polynucleobacter* strains, which obviously originated from free-living  
strains, which were present in large numbers in the ciliate cultures and as minor fractions in  
218 the ciliate homogenate used for the cultivation experiments. The growth of these bacteria

demonstrated, however, that the homogenates used for the cultivation experiments contained  
220 viable bacteria.

222 **Investigation of ciliate cultures by FISH.** The sequence of probe Omi10 published by  
Springer et al. (Springer et al., 1996) did not match with the 16S rRNA sequence of the *P.*  
224 *necessarius* type strain (ATCC30859) published by the same authors, but matched perfectly  
with the sequence obtained from the type strain in our study, as well as the sequence of *P.*  
226 *necessarius* strain STIR1 (Vannini et al., 2005). Therefore, we used in our study probe  
Omi10, as well as the new probe Omi10-mod, which was adjusted to the 16S rRNA sequence  
228 of *P. necessarius* ATCC30859 published by Springer et al. Both probes did not match (one or  
two mismatches) with any of the more than one hundred 16S rRNA sequences of all free-  
230 living strains, or with the sequences of the three endosymbiotic *Polynucleobacter* strains  
found in *E. harpa* (Vannini et al., 2005). Hybridization conditions appropriate for the  
232 discrimination of the *P. necessarius* endosymbionts of *E. aediculatus* from all other members  
of subcluster PnecC were established for both probes. The appropriate formamide  
234 concentration for the original probe Omi10 was found to be much higher than the  
concentration for probe Omi10-mod (65% versus 35%). This may indicate that the original  
236 probe fits perfectly to the sequence of the 16S rRNA gene, however, probe Omi10-mod also  
allowed the discrimination of *P. necessarius* endosymbionts from the other PnecC strains.  
238 FISH performed on *Euplotes* spp. cells picked one by one by a micropipette and fixed on a  
glass slide resulted in the detection of many *Polynucleobacter* symbionts inside of the ciliate  
240 cells, but never resulted in the detection of *Polynucleobacter*-like bacteria outside the ciliate  
cells. Different results, i.e. the detection of cells targeted by the endosymbiont-specific probe  
242 outside of ciliates were observed when ciliates were filtered onto 0.2µm pore size membrane  
filters prior to hybridization. The majority of probe-positive cells were located nearby to

244 ciliate cells (< 100 µm), which indicates that this detection of endosymbiont genotypes  
outside of ciliate cells was an artifact resulting from the damaging of ciliate cells during the  
246 filtration process.

Since the two Omi probes did not match the sequences of the *E. harpa* endosymbionts, probe  
248 Poly\_862 was used for the investigation of the *E. harpa* cultures. Again, probe-positive  
bacteria were detected exclusively inside of the host cells when ciliate cells have been  
250 handled carefully.

252 **Investigation of environmental samples by FISH.** Samples from the pelagic zone of five  
lakes and one pond were investigated by FISH (Table 3). The PnecC-specific probe PnecC-  
254 16S-445 detected free-living bacteria in samples from all six investigated freshwater habitats.  
All cells detected possessed small cell sizes typical for bacterioplankton cells. Probe-positive  
256 cells attached to particles or organisms, as well as cells located in any protist cell were not  
detected. Abundances of probe-positive cells ranged from  $3.7 \times 10^3$  to  $6.8 \times 10^5$  cells ml<sup>-1</sup>. By  
258 contrast, probe Omi10-mod, which is specific for endosymbionts of *E. aediculatus*, did not  
detect any cells in the investigated environmental samples.

260

**Experiments on enrichment and cultivation of *Euplotes* species from habitats containing**  
262 **free-living *Polynucleobacter* bacteria.** Enrichments were established from samples of three  
lakes and one pond. These three lakes are known to contain populations of free-living  
264 *Polynucleobacter* bacteria (Table 3). *Euplotes* species were not observed in any of the  
established enrichment cultures, however, ciliates belonging to other genera could be  
266 enriched and subsequently isolated (data not shown).

268 **Discussion**

The independent phylogenetic analysis of 16S rRNA genes and 16S-23S ITS sequences placed *Polynucleobacter* strains isolated from the pelagic of freshwater habitats, the endosymbionts of the freshwater ciliate *E. aediculatus*, and the endosymbionts of the brackish water ciliate *E. harpa* in the species-like subcluster PnecC of the *Polynucleobacter* cluster (Fig.2 and 3). This cluster represents in terms of genome size (Fig. 5) and G+C content of ITS sequences (Fig. 6B) a homogenous group well separated from other members of the family *Burkholderiaceae*. The minimal divergence of 16S rRNA genes (0.5%) of strains isolated from freshwater habitats and endosymbionts of *Euplotes* spp. suggested a close phylogenetic relationship of these organisms. Fundamental differences in lifestyle of such closely related bacteria have not been reported so far.

For two reasons, it is highly likely that the isolated strains existed as free-living, planktonic bacteria prior to isolation. First, the investigation of a free-living population of PnecC strains in the water column of a freshwater habitat by cultivation-independent and cultivation methods resulted in the detection and cultivation of the same ribosomal genotypes (Hahn et al., 2005). This demonstrated that the environmental sequences and the isolates obtained originated from a free-living population. Second, it is known, that the potential hosts of endosymbiotic *Polynucleobacter* strains, i.e. *Euplotes* spp., are usually absent from the pelagic of freshwater habitats (Foissner et al., 1991; Foissner et al., 1999). These ciliates usually inhabit interstitial, benthic sites rich in organic material (i.e., mud and detritus), and only in exceptional cases *Euplotes* spp. were reported from the pelagic of aquatic systems (Foissner et al., 1999). The few observations of planktonic *Euplotes* spp. were always restricted to anoxic parts of the pelagic zone, but all isolated *Polynucleobacter* strains were obtained from oxic water bodies. By contrast, the five strains termed above “endosymbiotic strains” undoubtedly represent endosymbionts of the two *Euplotes* species. This was

repeatedly demonstrated by FISH (Springer et al., 1996; Vannini et al., 2005), and electron  
294 microscopy (Vannini et al., 2005).

The occurrence of strains affiliated with the phylogenetically narrow subcluster PnecC as  
296 free-living and endosymbiotic stages could either be explained by a facultative  
endosymbiotic lifestyle of all members of the subcluster, or the members of the subcluster  
298 could fundamentally differ in lifestyle, i.e. could represent obligate endosymbionts and  
obligately free-living strains. The results presented here strongly argue against a facultative  
300 endosymbiotic lifestyle of the PnecC bacteria. The endosymbiotic strains demonstrated  
several characteristics of obligate symbionts. The analysis of the ribosomal sequences of the  
302 endosymbionts revealed a tight relationship between host species and endosymbiont  
genotypes (Fig. 2 and 3). This host-specific linkage stands in strong contrast to the large  
304 geographical distance between the sampling sites of the endosymbionts (Fig. 1), and to the  
significant ribosomal gene diversity within subcluster PnecC (Fig. 3). Both clearly indicate an  
306 obligate symbiotic relationship between the hosts and the endosymbionts. The cultivation  
experiments indicated pronounced physiological differences between the free-living stages  
308 and the endosymbiotic stages. The inability of the endosymbiotic strains to grow in media  
supporting growth of free-living strains could be a result of a genetic adaptation to the  
310 environmental conditions in the cytoplasm of the host cells. This may include the loss of  
metabolic capabilities necessary for the survival as free-living strains, or for growth in the  
312 provided media. The lack of culturability is one indicator for obligate endosymbiosis known  
from numerous bacterial endosymbionts, e.g., *Buchnera* (Tamas et al., 2002), *Wigglesworthia*  
314 (Aksoy, 1995), *Blochmannia* (Sauer et al., 2000), and is interpreted as a loss of unneeded  
metabolic pathways (Moran, 2002). The lack of culturability of the *Polynucleobacter*  
316 endosymbionts, and the previously reported dependence of host cells on their endosymbionts  
(Heckmann, 1975; Vannini et al., 2005) indicate an obligate interdependence of

318 endosymbionts and hosts, which is a typical characteristic of obligate endosymbiosis (Zientz  
et al., 2004). The loss of genes encoding dispensable metabolic function also results in  
320 reduced genome sizes of obligate endosymbiotic bacteria (Moran, 2002). In several cases,  
their genomes are reduced to ~20% of the genome size of the closest related free-living  
322 relatives (Moran et al., 2003). The majority of obligate endosymbionts investigated so far  
possess genome sizes of < 1.5 Mb, while the majority of free-living bacteria possess genome  
324 sizes of 2 to 10 Mb (Moran, 2002). In contrast to all obligate endosymbionts investigated so  
far (e.g. Shigenobu et al., 2000, Tamas et al., 2002) the genome size of the endosymbiotic  
326 *Polynucleobacter* differed much less significantly from that of the closest free-living relatives  
(Fig. 5). Another characteristic of the genomes of obligate endosymbionts is the accumulation  
328 of mildly deleterious mutations (Lambert and Moran, 1998). Our analysis indicated  
destabilizing base substitutions in the 16S rRNA genes of *E. harpa* endosymbionts but not in  
330 the *E. aediculatus* and the free-living strains (Fig. 4). Furthermore, the two groups of  
endosymbionts showed each for both investigated phylogenetic markers, i.e. the 16S rRNA  
332 genes and the ITS sequences, the highest divergence from other members of subcluster  
PnecC. This may indicate elevated rates of sequence evolution compared to free-living  
334 relatives, which was repeatedly demonstrated for other obligate endosymbionts (Aksoy,  
1995; Herbeck et al, 2003).

336 The lack of pronounced genome reduction in the *Polynucleobacter* endosymbionts, as well  
as the lack of an increased number of mismatches in helix regions of the 16S rRNA of *E.*  
338 *aediculatus* endosymbionts is in agreement with the proposed close phylogenetic relationship  
between the endosymbionts and their closest free-living relatives. This close relationship  
340 suggests that the time point of divergence of the endosymbiotic and free-living  
*Polynucleobacter* bacteria was much more recent than in the case of other obligate

342 endosymbionts. In the case of *Buchnera* and their closest free-living relatives a divergence  
period of >100 million years was estimated (Moran et al., 2003).

344 In contrast to the endosymbiotic strains, an endosymbiotic lifestyle was not evident for  
the free-living strains. Large free-living *Polynucleobacter* populations were observed in the  
346 pelagic zone of habitats from which potential hosts, i.e. *Euplotes* spp., could not be enriched.  
Even if *Euplotes* spp. have been present in these habitats, their population sizes must have  
348 been several orders of magnitude smaller than the size of the free-living *Polynucleobacter*  
populations. This has to be expected due to the known restriction of these ciliates to the  
350 benthic zone, and due to further restrictions caused by a partially low habitat suitability  
(Foissner et al., 1991; Foissner et al., 1999) of the benthic zone in these lakes. The resulting  
352 imbalance between the population size of free-living *Polynucleobacter* bacteria and potential  
hosts makes a symbiotic relationship between these organisms unlikely. Furthermore, the  
354 free-living and the endosymbiotic *Polynucleobacter* strains differ from known facultative  
symbionts (e.g. *Vibrio fischeri*) and known facultative parasites of protists (e.g. *Legionella*  
356 spp.) in the lack of motility. Due to this lack *Polynucleobacter* strains should be in principal  
unable to chemotactically find potential hosts.

358 **Conclusions.** The presented results indicate that the phylogenetically shallow subcluster  
PnecC harbors both, obligately free-living strains and obligate endosymbionts of at least two  
360 ciliate species. Consequently, this subcluster harbors at least three groups of organisms  
occupying completely separated ecological niches. Niche differentiation is well documented  
362 for more distantly related bacteria. For instance, Gray et al. (2004) demonstrated niche  
differentiation in sympatric *Achromatium* spp., which are separated by 16S rRNA gene  
364 sequence similarities of 91.5%, and Schauer et al. (2005) demonstrated complete niche  
separation in two groups of filamentous bacteria both affiliated with the SOL cluster but  
366 separated by 16S rRNA gene sequence similarities of 92.1%. The finding of niche separation



in *Polynucleobacter* bacteria, as well as the ecological differentiation in *Brevundimonas alba*  
368 strains sharing identical 16S rRNA gene sequences (Jaspers & Overmann, 2004) highlights  
the ecological significance of microdiversity.

370 To our best knowledge, the phylogenetic relationship between the obligately  
endosymbiotic and the obligately free-living *Polynucleobacter* bacteria is the closest  
372 currently known. The closest free-living relatives of all other obligate endosymbionts  
investigated so far are more distantly related (Table 2). We did not find any example of an  
374 obligate endosymbiont, which also shares a 16S rRNA gene sequence similarity >99% with  
its closest free-living relative. Further investigations on the evolutionary adaptations of  
376 *Polynucleobacter* endosymbionts and their hosts may provide exciting insights into processes  
taking place during the early evolutionary phase of endosymbiosis.

378

### **Experimental procedures**

380 **Microbial strains, culture conditions, and media.** The *Polynucleobacter* endosymbionts of  
five ciliate strains of the genus *Euplotes* (Hypotrichia) were investigated (Table 1). Three of  
382 the strains belong to the species *Euplotes harpa*, and two strains belong to *Euplotes*  
*aediculatus*. The strain *E. aediculatus* ATCC 30859 harbors the type strain of  
384 *Polynucleobacter necessarius*. The three *E. harpa* strains were cultured in artificial seawater  
at low salinity (5‰) at 20°C. The *E. aediculatus* strain STIR1 was cultured in sterile spring  
386 water enriched with boiled rice grains at 20°C, and *E. aediculatus* strain ATCC 30859 was  
cultured in sterile WC medium (Guillard, 1975) at 22°C. *Euplotes aediculatus* strain ATCC  
388 30859 was fed with the algae *Cryptomonas* sp. SAG26.80, and the other strains were fed with  
the green algae *Dunaliella tertiolecta*. All ciliate cultures were non-axenic. The cultures were  
390 kept in the light, either under permanent illumination (*E. aediculatus* ATCC 30859), or under  
a light dark cycle of 12:12 hours (other strains).

392 **Determination of 16S rRNA gene sequences and 16S-23S internal transcribed spacer**  
**(ITS1) sequences.** For the determination of ribosomal sequences of endosymbiotic  
394 *Polynucleobacter* strains, the host ciliate cells were separated from free-living bacteria also  
present in the ciliate cultures by filtration onto a 15µm pore size membrane filter, washed  
396 with sterile medium, and finally resuspended in sterile medium. Ciliates were pelleted by  
centrifugation at 200g for ten minutes. DNA was extracted from the obtained biomass. The  
398 forward *Polynucleobacter*-specific primer 16S-Poly-F467 (Vannini et al., 2005) and a reverse  
primer binding at position 457 (*E. coli* numbering) of the 23S rRNA gene (Ludwig et al.,  
400 1992) were used for the amplification of ITS sequences of the endosymbionts from three *E.*  
*harpa* strains and from *E. aediculatus* STIR1. For the amplification of the 16S rRNA gene  
402 and the 16S-23S internal transcribed spacer (ITS) of the endosymbiont of *E. aediculatus*  
strain E24 (ATCC 30859) the forward primer 19F (Hahn et al., 2005) and the PnecC-specific  
404 reverse primer PnecCr-5 (Hahn et al., 2005) were used. Amplification of 16S rRNA genes  
and ITS sequences of free-living strains were performed as described previously (Hahn,  
406 2003; Hahn et al., 2005).

**Comparative sequence analysis.** Sequences were aligned with the autoalignment tool  
408 provided by the software package ARB (Ludwig et al., 2004), and alignment errors were  
corrected manually. Sequence polymorphisms were analyzed with the software DnaSP 4.1  
410 (Rozas et al., 2003). Phylogenetic trees were calculated by using several methods. The  
DNAPARS program was used for maximum parsimony analysis (Felsenstein, 1989), with a  
412 bootstrap analysis of 1,000 pseudoreplicates. A maximum likelihood analysis (Felsenstein,  
1981; Olsen et al., 1994) was performed by the FastDNAm1 tool, and a neighbor joining  
414 analysis (Saitou and Nei, 1987) was performed by the DNADIST tool with Kimura correction  
(Kimura, 1980), as well as by the software Mega2 (Kumar et al., 2001).

416 **Determination of genome size.** The genome size of three endosymbiotic strains, eight free-  
living PnecC, and three free-living PnecD strains was determined by pulsed field gel  
418 electrophoresis (PFGE). *Polynucleobacter*-harboring ciliates and free-living  
*Polynucleobacter* cells were harvested and washed in sterilized artificial sea-water and  
420 phosphate-buffered saline, respectively. Pellets were immobilized in agarose plugs and  
treated as described by Strous et al. (2006). PFGE was then performed in a contour-clamped  
422 homogeneous electric field system (CHEF-DR III; Bio-Rad) at 11°C, using 1x TAE pH 8 and  
1% Seakem Gold Agarose (Cambrex). Gels were run by three resolution blocks, differing for  
424 pulse, angle and time. Fragment length was estimated by comparison with standard size  
ladders (*Saccharomices cerevisiae* and *Hansenula wingei* PFGE Markers; Bio-Rad).

426 **Cultivation experiments with endosymbionts.** Cultivation experiments with  
endosymbionts of *E. aediculatus* strain STIR1 and of *E. harpa* strain FSP1.4 were performed  
428 by using the acclimatization method (Hahn et al., 2004), which was successfully used for the  
cultivation of environmental isolates (Hahn, 2003; Hahn et al., 2005). For the separation of  
430 endosymbionts from free-living bacteria present in the non-axenic ciliate cultures, the ciliate  
cells were filtered through a 100 µm pore-size filter, harvested onto 15 µm pore-size filters,  
432 and resuspended in sterile IBM medium (Hahn et al., 2004). Forty ml of these suspensions  
were harvested by centrifugation at 200g (10 min) and concentrated in 4 ml IBM medium.

434 The endosymbionts were released from the hosts by destroying the ciliates by centrifugation  
at 6,000g (15 min) and subsequent vortexing. Presence of released endosymbionts in the  
436 homogenate was verified by staining with DAPI (4',6'-diamino-2-phenylindole) and  
epifluorescence microscopical inspection. Furthermore, the presence of free-living bacteria in  
438 the homogenate was tested by DAPI-staining and hybridization with the probe PnecC-16S-  
445 (see below), which is specific for the *Polynucleobacter* cluster but did not hybridize with  
440 free-living bacteria present in the ciliate cultures. The homogenates containing the highly

enriched endosymbionts were used for two different cultivation experiments. (i) Samples  
442 were processed according to the standard protocol of the filtration-acclimatization method  
(FAM (Hahn et al., 2004)) (Table 4). This method employs filtration through 0.2µm filters  
444 for the separation of target organisms from organisms able to overgrow the target organisms.  
(ii) Samples were processed by the dilution acclimatization method (DAM (Hahn et al.,  
446 2005)). This method employs a dilution step instead of the filtration step. Briefly, samples of  
100 µl of the homogenate were diluted into six different concentrations (1:1 to 1:100000).  
448 One hundred µl of the dilutions were used for the inoculation of wells of sterile cell culture  
plates (24-wells) containing 500 µl of the starting medium. Four different approaches were  
450 tried in the DAM experiments: (i) stepwise acclimatization to NSY 3 gl<sup>-1</sup> (Hahn et al., 2004)  
using IBM as a starting medium; (ii) acclimatization to NSY 3 gl<sup>-1</sup> using ciliate homogenate  
452 filtered through a 0.2 µm pore-size membrane as a starting medium, and stepwise addition of  
NSY 3 gl<sup>-1</sup>; (iii) cultivation attempts without any acclimatization (i.e. inoculation of NSY 3gl<sup>-1</sup>  
454 <sup>1</sup> with dilutions of the enriched endosymbionts), and (iv) cultivation in ciliate homogenate  
filtered through 0.2 µm pore-size filters without increasing the substrate concentration (i.e. no  
456 addition of medium). For experiments (i) and (ii) the standard acclimatization protocol  
described by Hahn et al. (2004) was followed (i.e. stepwise addition of increasing amounts of  
458 dissolved organic material). In experiments (ii) and (iv) controls receiving no inoculum were  
used for testing sterility of the 0.2µm-filtered ciliate homogenate. Screening of the wells for  
460 the growth of bacteria was done by dropping 10 µl samples from each well onto NSY agar  
plates, or in the case of experiment iv onto 0.2 µm-filtered ciliate homogenate solidified with  
462 agar (15 g/l). Samples of colonies grown on plates were tested for the presence of  
*Polynucleobacter* bacteria by FISH using the probes Omi10-mod and PnecC-16S-445 (see  
464 below).

**Experiments on the isolation of *Euplotes* strains from freshwater habitats.** The benthic  
466 zone of Lake Krottensee, Lake Wolfgangsee, Lake Egelsee (M2), and an unnamed pond were  
sampled at sites appropriate for freshwater *Euplotes* species. Samples were microscopically  
468 screened for the presence of *Euplotes* spp. immediately after sampling, as well as after  
several days of incubation. Aliquots of the samples were also enriched with prey organisms  
470 appropriate for *Euplotes* (i.e. *Dunaliella tertiolecta*, *Phaeodactylum* sp., *Enterobacter*  
*aerogenes*) and daily screened for the presence of *Euplotes* spp. over a twenty-days period.

**Analysis of ciliate cultures and environmental samples by FISH.** Samples taken from  
472 ciliate cultures and from the pelagic of six freshwater habitats located in the Austrian  
Salzkammergut area (near Salzburg) were investigated by fluorescence in situ hybridisation  
474 (FISH). Twenty ml samples were fixed with paraformaldehyde solution (2% final  
concentration) and filtered onto 0.2µm polycarbonate membrane filters as described  
476 elsewhere (Hahn et al., 2005). Whole-cell *in situ* hybridizations of sections of the filters were  
performed as described previously (Hahn et al., 2005). In parallel, ciliate cells were also  
478 processed in a different way: they were picked one by one, together with their culture  
medium, by a glass micropipet and then fixed on microscope slides. Slides were then  
480 processed for FISH as described elsewhere (Petroni et al., 2003). Four different Cy3-labelled  
oligonucleotide probes were used for the hybridization of samples. (i) Probe PnecC-16S-445  
482 (Hahn et al., 2005), which is specific for the entire subcluster C (PnecC) of the  
*Polynucleobacter* cluster. (ii) Probe Omi10 (Springer et al., 1996), which is specific for the  
endosymbiotic *Polynucleobacter necessarius*, (iii) probe Omi10-mod (5'-TTT CCC CCC  
486 TTC AGG GCG T-3', the underlined nucleotide is not contained in the probe Omi10), which  
is a slightly modified version of Omi10, and (iv) probe Poly\_862 (Vannini et al., 2005),  
488 which is specific for the entire *Polynucleobacter* cluster. Probe Omi10 is matching the 16S  
rRNA sequence of *P. necessarius* strain ATCC30859 (sequence obtained in this study), and

490 the sequence of *P. necessarius* STIR1 (Vannini et al., 2005), while probe Omi10-mod is  
matching the previously published sequence of *P. necessarius* ATCC30859 (Springer et al.,  
492 1996). Probe Omi10-mod was applied with 35% formamid concentration in the hybridization  
buffer, probe Omi10 with 65%, and probe Poly\_862 with 0% formamid (Vannini et al.,  
494 2005).

**Nucleotide sequence accession numbers.** The obtained 16S-23S ITS sequences of  
496 endosymbiotic and free-living PnecC strains were submitted to the DDBJ/EMBL/GenBank  
databases under accession numbers XXXXXXXXX to XXXXXXXXX. The 16S rRNA gene  
498 sequences of these strains have been deposited previously.

#### 500 **Acknowledgments**

We are grateful to Lauri Arvola and Marko Järvinen for providing samples from Finnish  
502 lakes, Karel Šimek and Jan Jezbera for providing samples from two Brazilian habitats, and  
Silke Langenheder for providing samples from Swedish lakes, as well as Jannik Pitt for  
504 assistance in sampling of habitats at Corsica. Gabriel Zwart and Paul Bodelier are  
acknowledged for their contributions in the isolation of a Polynucleobacter strain from a  
506 Dutch habitat. This study was supported by the bilateral exchange program between the  
Austrian Academy of Sciences and the Italian Accademia Nazionale dei Lincei, as well as by  
508 the Italian Ministry of University and Research, and the Austrian Science Fund (project  
P15655).

510

510 **References**

- 512 Aksoy, S. 1995. *Wigglesworthia* gen. nov. and *Wigglesworthia glossinidia* sp. nov., taxa  
consisting of the mycetocyte-associated, primary endosymbionts of tsetse flies. Int. J.  
Syst. Bacteriol. **45**: 848-851.
- 514 Bahr, M., Hobbie, J.E. and Sogin, M.L. 1996. Bacterial diversity in an arctic lake: a  
freshwater SAR11 cluster. Aquat. Microb. Ecol. **11**: 271-277.
- 516 Distel, D.L., Lane, D.J., Olsen, G.J., Giovannoni, S.J., Pace, B., Pace, N.R., Stahl, D.A.  
and Felbeck, H. 1988. Sulfur-oxidizing bacterial endosymbionts: analysis of phylogeny  
518 and specificity by 16S rRNA sequences. J. Bacteriol. **170**: 2506-2510.
- Distel, D.L., Lee, H.K. and Cavanaugh, C.M. 1995. Intracellular coexistence of methano-  
520 and thioautotrophic bacteria in a hydrothermal vent mussel. Proc. Natl. Acad. Sci.  
U.S.A. **92**: 9598-9602.
- 522 Everett, K.D., Bush, R.M., and Andersen, A.A. 1999. Emended description of the order  
*Chlamydiales*, proposal of *Parachlamydiaceae* fam. nov. and *Simkaniaceae* fam. nov.,  
524 each containing one monotypic genus, revised taxonomy of the family *Chlamydiaceae*,  
including a new genus and five new species, and standards for the identification of  
526 organisms. Int. J. Syst. Bacteriol. **49**: 415-440.
- Felsenstein, J. 1981. Evolutionary trees from DNA sequences a maximum likelihood  
528 approach. J. Mol. Evol., **17**: 368-376.
- Felsenstein, J. 1989. PHYLIP - Phylogeny inference package (Version 3.2). Cladistics,  
530 **5**:164-166.
- Foissner, W., Berger, H., and Schaumburg, J. 1999. Identification and ecology of limnetic  
532 plankton ciliates. - Informationsberichte des Bayerischen Landesamtes für  
Wasserwirtschaft, 3/99: 1-793.
- 534 Foissner, W., Blatterer, H., Berger, H., and Kohmann, F. 1991. Taxonomische und  
ökologische Revision der Ciliaten des Saprobiensystems. – Band I: Cyrtophorida,

- 536 Oligotrichida, Hypotrichida, Colpodea. – Informationsberichte des Bayer. Landesamtes  
für Wasserwirtschaft, Heft 1/91, 1-478
- 538 Fritsche, T.R., Horn, M., Seyedirashti, S., Gautom, R.K., Schleifer, K.H., and Wagner, M.  
1999. In situ detection of novel bacterial endosymbionts of *Acanthamoeba* spp.  
540 phylogenetically related to members of the order Rickettsiales. Appl. Environ.  
Microbiol. **65**: 206-212.
- 542 Fujishima, M., and Heckmann, K. 1984. Intra- and interspecies transfer of endosymbionts  
in *Euplotes*. J. Exp. Zool. **230**:3 39-345.
- 544 Görtz, H.D. 2002. Symbiotic association between ciliates and prokaryotes. The  
prokaryotes, electronic edition. Release 3.11.  
546 [http://141.150.157.117:8080/prokPUB/chaprender/jsp/showchap.jsp?chapnum=355&in  
itsec=04\\_01](http://141.150.157.117:8080/prokPUB/chaprender/jsp/showchap.jsp?chapnum=355&in<br/>itsec=04_01)
- 548 Gray, N.D., Comaskey, D., Miskin, I.P., Pickup, R.W., Suzuki, K., and Head, I.M. 2004.  
Adaptation of sympatric *Achromatium* spp. to different redox conditions as a  
550 mechanism for coexistence of functionally similar sulphur bacteria. Environ Microbiol.  
**6**: 669-677.
- 552 Guillard, R.R. 1975. Cultures of phytoplankton for feeding of marine invertebrates, p. 29–  
60. In W. L. Smith and M. H. Chanley [eds.], Culture of marine invertebrate animals.  
554 Plenum.
- Hahn, M.W. 2003. Isolation of strains belonging to the cosmopolitan *Polynucleobacter*  
556 *necessarius* cluster from freshwater habitats located in three climatic zones. Appl.  
Environ. Microbiol. **69**: 5248-5254.
- 558 Hahn, M.W., M. Pöckl, and Wu, Q.L. 2005. Low intraspecific diversity in a  
*Polynucleobacter* subcluster population numerically dominating bacterioplankton of a  
560 freshwater pond. Appl. Environ. Microbiol. **71**: 4539–4547.



- 562 Hahn, M.W., Stadler, P., Wu, .Q.L., and Pöckl, M. 2004. The filtration-acclimatization-  
method for isolation of an important fraction of the not readily cultivable bacteria. J.  
Microb. Meth. **57**: 379-390.
- 564 Heckmann, K. 1975. Omikron, ein essentieller Endosymbiont von *Euplotes aediculatus*. J.  
Protozool. **22**: 97-104.
- 566 Heckmann, K., and Schmidt, H.J. 1987. *Polynucleobacter necessarius* gen. nov., sp. nov.,  
an obligately endosymbiotic bacterium living in the cytoplasm of *Euplotes aediculatus*.  
568 Int. J. Syst. Bacteriol. **37**: 456-457.
- Heckmann, K., Ten Hagen, R. and Görtz, H.-D. 1983. Freshwater *Euplotes* species with 9  
570 type I cirrus pattern depend upon endosymbionts. J. Protozool. **30**: 284-289.
- Herbeck, J.T., Funk, D.J., Degnan, P.H., and Wernegreen, J.J. 2003. A conservative test of  
572 genetic drift in the endosymbiotic bacterium Buchnera: slightly deleterious mutations in  
the chaperonin groEL. Genetics **165**: 1651-1660.
- 574 Hiorns, W.D., Methé, E.A., Nierzwickibauer, S.A., and Zehr, J.P. 1997. Bacterial diversity  
in Adirondack mountain lakes as revealed by 16S rRNA gene sequences. Appl.  
576 Environ. Microbiol. **63**: 2957-2960.
- Horn, M., Fritsche, T.R., Linner, T., Gautom, R.K., Harzenetter, M.D., and Wagner, M.  
578 2002. Obligate bacterial endosymbionts of *Acanthamoeba* spp. related to the beta-  
Proteobacteria: proposal of '*Candidatus Procabacter acanthamoebae*' gen. nov., sp. nov.  
580 Int. J. Syst. Evol. Microbiol. **52**: 599-605.
- Jaspers, E., and Overmann, J. 2004. Ecological significance of microdiversity: Identical  
582 16S rRNA gene sequences can be found in bacteria with highly divergent genomes and  
ecophysologies. Appl. Environ. Microbiol. **70**: 4831-4839.
- 584 Kimura, M. 1980. A simple method for estimating evolutionary rates of base substitutions  
through comparative studies of nucleotide-sequences. J. Mol. Evol., **16**: 111-120.
- 586 Kumar S., Tamura, K., Jakobsen, I.B., and Nei, M. 2001. MEGA2: molecular evolutionary  
genetics analysis software. Bioinformatics. **17**: 1244-5.

588 Kurtti, T.J., Munderloh, U.G., Andreadis, T.G., Magnarelli, L.A., and Mather, T.N. 1996.  
Tick cell culture isolation of an intracellular prokaryote from the tick *Ixodes scapularis*.  
590 *J. Invertebr. Pathol.* **67**: 318-321.

Lambert, J.D., and Moran, N.A. 1998. Deleterious mutations destabilize ribosomal RNA  
592 in endosymbiotic bacteria. *Proc. Natl. Acad. Sci. USA* **95**: 4458–4462.

Ludwig, W., Kirchhof, G., Klugbauer, N., Weizenegger, M., Betzl, D., Ehrmann, M.,  
594 Hertel, C., Jilg, S., Tatzel, R., Zitzelsberger, H., Liebl, S., Hochberger, M., Lane, D.,  
Wallnofer, P.R., and Schleifer, K.H. 1992. Complete 23S ribosomal RNA sequences of  
596 gram-positive bacteria with a low DNA G+C content. *Syst. Appl. Microbiol.*, **15**: 487-  
501.

Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, Buchner, A.,  
598 Lai, T., Steppi, S., Jobb, G., Förster, W., Brettske, I., Gerber, S., Ginhart, A. Gross, O.,  
600 Grumann, S., Hermann, S., Jost, R., König, A., Liss, T., Lüßmann, R., May, M.,  
Nonhoff, B., Reichel, B., Strehlow, R., Stamatakis, A., Stuckmann, N., Vilbig, A.,  
602 Lenke, M., Ludwig, T., Bode, A., and Schleifer, K. H. 2004. ARB: a software  
environment for sequence data. *Nucl. Ac. Res.*, **32**: 1363-1371.

604 Moran, N.A. 2002. Microbial minimalism: genome reduction in bacterial pathogens. *Cell*  
**108**:583-586

606 Moran, N.A., Plague, G., Sandström, J. P., and Wilcox, J. A. 2003. A genomic perspective  
on nutrient-provisioning by bacterial symbionts of insects. *Proc. Natl. Acad. Sci. USA*  
608 **100**:1 4543-14548.

Olsen, G.J., Matsuda, H., Hagstrom, R., and Overbeek, R. 1994. FastDNAm1: A tool for  
610 construction of phylogenetic trees of DNA sequences using maximum likelihood.  
*Comput. Appl. Biosci.* **10**: 41-48.

- 612 Pedersen, K., Arlinger, J., Ekendahl, S., and Hallbeck, L. 1996. 16S rRNA gene diversity  
of attached and unattached bacteria in boreholes along the access tunnel to the Aespoe  
614 hard rock laboratory Sweden. *FEMS Microbiol. Ecol.* **19**:249-262.
- Petroni, G., Dini, F., Verni, F., and Rosati, G. 2002. A molecular approach to the tangled  
616 intrageneric relationships underlying phylogeny in *Euplotes* (Ciliophora, Spirotrichea).  
*Mol. Phylogenet. Evol.* **22**: 118–130.
- 618 Petroni, G., Rosati, G., Vannini, C., Modeo, L., Dini, F., and Verni, F. 2003. *In situ*  
identification by fluorescently labeled oligonucleotide probes of morphologically  
620 similar, closely related ciliate species. *Microb. Ecol.* **45**: 156-162.
- Rozas, J., Sanchez-DelBarrio, J.C., Messeguer, X., and Rozas, R. 2003. DnaSP, DNA  
622 polymorphism analyses by coalescent and other methods. *Bioinformatics* **19**: 2496-  
2497.
- 624 Saitou, N., and Nei, M. 1987. The neighbor-joining method: a new method for  
reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**: 406-425.
- 626 Sauer, C., Stackebrandt, E., Gadau, J., Hölldobler, B. and Gross, R. 2000. Systematic  
relationships and cospeciation of bacterial endosymbionts and their carpenter ant host  
628 species: proposal of the new taxon *Candidatus* Blochmannia gen. nov. *Int. J. Syst.*  
*Evol. Microbiol.* **50**: 1877-1886.
- 630 Schauer, M., Kamenik, C., and Hahn, M.W. 2005. Ecological differentiation within a  
cosmopolitan group of planktonic freshwater bacteria (SOL cluster, Saprospiraceae,  
632 Bacteroidetes). *Appl. Environ. Microbiol.* **71**: 5900-5907.
- Semenova, E.A., and Kuznedelov, K.D. 1998. A study of the biodiversity of Baikal  
634 picoplankton by comparative analysis of 16S rRNA gene 5'-terminal regions. *Mol.*  
*Biol.* **32**: 754-760.
- 636 Shigenobu, S., Watanabe, H., Hattori, M., Sakaki, Y. and Ishikawa, H. 2000. Genome  
sequence of the endocellular bacterial symbiont of aphids *Buchnera* sp. *APS. Nature*  
638 **407**: 81-86.

- Šimek, K., Pernthaler, J., Weinbauer, M.G., Horňák, K., Dolan, J.R., Nedoma J., Masin,  
640 M., and Amann, R. 2001. Changes in bacterial community composition and dynamics  
and viral mortality rates associated with enhanced flagellate grazing in a mesoeutrophic  
642 reservoir. *Appl. Environ. Microbiol.* **67**: 2723-33.
- Springer, N., Amann, R., Ludwig, W., Schleifer, K.H., and Schmidt, H. 1996.  
644 *Polynucleobacter necessarius*, an obligate bacterial endosymbiont of the hypotrichous  
ciliate *Euplotes aediculatus*, is a member of the beta-subclass of *Proteobacteria*. *FEMS*  
646 *Microbiol. Lett.* **135**: 333-336.
- Springer, N., Ludwig, W., Amann, R., Schmidt, H.J., Görtz, H.D., and Schleifer, K.H.  
648 1993. Occurrence of fragmented 16S rRNA in an obligate bacterial endosymbiont of  
*Paramecium caudatum*. *Proc. Natl. Acad. Sci. U.S.A.* **90**: 9892-9895.
- 650 Strous, M., Pelletier, E., Mangenot, S., Rattei, T., Lehner, A., Taylor, M. W., Horn, M.,  
Daims, H., Bartol-Mavel, D., Winker, P., Barbe, V., Fonknechten, N., Vallenet, D.,  
652 Segurens, B., Schenowitz-Truong, C., Médigue, C., Collingro, A., Snel, B., Dutilh,  
B.E., Op den Camp, H.J.M., van der Drift, C., Cirpus, I., van de Pas-Schoonen, K.T.,  
654 Harhangi, H.R., van Niftrik, L., Schmid, M., Keltjens, J., van de Vossenberg, J., Kartal,  
B., Meier, H., Frishman, D., Huynen, M.A., Mewes, H.W., Weissenbach, J., Jetten,  
656 M.S.M., Wagner, M. and Le Paslier, D. 2006. Deciphering the evolution and  
metabolism of an anammox bacterium from a community genome. *Nature* **440**: 790-  
658 794.
- Tamas, I., Klasson, L., Canback, B., Naslund, A.K., Eriksson, A.S., Wernegreen, J.J.,  
660 Sandstrom, J.P., Moran, N.A., and Andersson, S.G. 2002. 50 million years of genomic  
stasis in endosymbiotic bacteria. *Science* **296**: 2376-2379.
- 662 Thao, M.L., Gullan, P.J., and Baumann, P. 2002. Secondary (*gamma-Proteobacteria*)  
endosymbionts infect the primary (*beta-Proteobacteria*) endosymbionts of mealybugs  
664 multiple times and coevolve with their hosts. *Appl. Environ. Microbiol.* **68**: 3190-3197.

666 Unterman, B.M., Baumann, P., and McLean, D.L. 1989. Pea aphid symbiont relationships  
established by analysis of 16S rRNAs. *J. Bacteriol.* **171**: 2970-2974.

668 Van Oevelen, S., De Wachter, R., Vandamme, P., Robbrecht, E. and Prinsen, E. 2004.  
'*Candidatus Burkholderia calva*' and '*Candidatus Burkholderia nigropunctata*' as leaf  
gall endosymbionts of African *Psychotria*. *Int. J. Syst. Evol. Microbiol.* **54**: 2237-2239.

670 Vannini, C., Petroni, G., Verni, F., and Rosati, G. 2005 *Polynucleobacter* bacteria in the  
brackish-water species *Euplotes harpa* (Ciliata, *Hypotrichia*). *J. Eukariot. Microbiol.*  
672 **52**: 116-22.

Vannini, C., Rosati, G., Verni, F., and Petroni, G.. 2004. Identification of the bacterial  
674 endosymbionts of the marine ciliate *Euplotes magnicirratu*s (Ciliophora, Hypotrichia)  
and proposal of '*Candidatus Devosia euplotis*'. *Int. J. Syst. Evol. Microbiol.* **54**: 1151-  
676 1156.

von Dohlen, C.D., Kohler, S., Alsop, S.T., and McManus, W.R. 2001. Mealybug beta-  
678 proteobacterial endosymbionts contain gamma-proteobacterial symbionts. *Nature.* **412**:  
433-436.

680 Wu, Q.L., and Hahn, M.W. (2006a) Differences in structure and dynamics of  
*Polynucleobacter* communities in a temperate and a subtropical lake, revealed at three  
682 phylogenetic levels. *FEMS Microb. Ecol.* **57**: 67-79.

Wu, Q.L., and Hahn, M.W. (2006b) High predictability of the seasonal dynamics of a  
684 species-like *Polynucleobacter* population in a freshwater lake. *Environ. Microbiol.* (in  
press). doi: 10.1111/j.1462-2920.2006.01049.x

686 Zientz, E., Dandekar, T., and Gross, R. (2004) Metabolic interdependence of obligate  
intracellular bacteria and their insect hosts. *Microbiol. Mol. Biol. Rev.* **68**: 745-770.

688 Zwart, G., Crump, B. C., Kamst-van Agterveld, M.P., Hagen, F., and Han, S. K. 2002.  
Typical freshwater bacteria: an analysis of available 16S rRNA gene sequences from  
690 plankton of lakes and rivers. *Aquat. Microb. Ecol.* **28**: 141-155.

This work was performed under the auspices of the US Department of Energy's Office of Science, Biological and Environmental Research Program, and by the University of California, Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48, Lawrence Berkeley National Laboratory under contract No. DE-AC02-05CH11231 and Los Alamos National Laboratory under contract No. DE-AC52-06NA25396

**Table 1.** Investigated endosymbionts and their hosts.

| Endosymbiont                        | Host                        | Host strain             | Origin         |                   | Deposition        | Reference               |
|-------------------------------------|-----------------------------|-------------------------|----------------|-------------------|-------------------|-------------------------|
|                                     |                             |                         | Habitat        | Site              |                   |                         |
| <i>Polynucleobacter</i> sp.         | <i>Euplotes harpa</i>       | FSP1.4                  | Brackish water | Pisa, Italy       | -                 | Vannini et al., 2005    |
| <i>Polynucleobacter</i> sp.         | <i>Euplotes harpa</i>       | BOD2                    | Brackish water | Boderne, Denmark  | -                 | Petroni et al., 2002    |
| <i>Polynucleobacter</i> sp.         | <i>Euplotes harpa</i>       | FC1                     | Brackish water | Taranto, Italy    | -                 | Vannini et al., 2005    |
| <i>Polynucleobacter necessarius</i> | <i>Euplotes aediculatus</i> | STIR1                   | Freshwater     | Parma, Italy      | -                 | Petroni et al., 2002    |
| <i>Polynucleobacter necessarius</i> | <i>Euplotes aediculatus</i> | ATCC 30859 <sup>a</sup> | Freshwater     | Marseille, France | ATCC <sup>b</sup> | Heckmann & Schmidt, 198 |

<sup>a</sup> ATCC 30859 is identical with clone 15 of *E. aediculatus* (Heckmann & Schmidt, 1987)

<sup>b</sup> ATCC: American Type Culture Collection

**Table 2.** 16S rRNA similarities between obligate endosymbionts and the known closest related free-living species or strains. Closest relatives of the endosymbionts were found by BLAST. Sequence similarities were determined after alignment of downloaded sequences by using the software program ARB.

| Obligate Endosymbiont                                       | Accession Number | Host  | Closest free-living species/strain        | Phylogenetic distance (16S rRNA similarity) | Reference (for endosymbiont) |
|---|------------------|---|---|---|------------------------------|
| Endosymbiont of <i>Ixodes scapularis</i>                    | AB001518         | Tick ( <i>Ixodes scapularis</i> )             | <i>Flexibacter tractuosus</i>             | 83.1 %                                      | Kurtti et al., 1996          |
| Endosymbiont of <i>Acanthamoeba</i> sp. UWC8                | AF069963         | Amoebae ( <i>Acanthamoeba</i> sp. UWC8)       | <i>Ehrlichia ruminantium</i> <sup>b</sup> | 84.3 %                                      | Fritsche et al., 1999        |
| <i>Candidatus Tremblaya princeps</i>                        | AF476095         | Mealybug ( <i>Pseudococcus viburni</i> )      | <i>Burkholderia cepacia</i>               | 85.6 %                                      | Thao et al., 2002            |
| <i>Caedibacter caryophyllus</i>                             | X71837           | Ciliate ( <i>Paramecium caudatum</i> )        | <i>Azorhizobium caulinodans</i>           | 85.7 %                                      | Springer et al., 1993        |
| <i>Wigglesworthia glossinidia</i>                           | L37342           | Tsetse flies ( <i>Glossina tachinoides</i> )  | <i>Pectobacterium carotovorum</i>         | 88.0 %                                      | Aksoy, 1995                  |
| <i>Candidatus Blochmannia tulipes</i>                       | X92552           | Ant ( <i>Camponotus rufipes</i> )             | <i>Serratia marcescens</i>                | 88.5 %                                      | Sauer et al., 2000           |
| <i>Candidatus Procabacter acanthamoebae</i>                 | AF177427         | Amoebae ( <i>Acanthamoeba</i> sp. UWC12)      | <i>Azovibrio restrictus</i>               | 88.7 %                                      | Horn et al., 2002            |
| <i>Buchnera aphidicola</i> str. APS                         | NC_002528        | Aphid ( <i>Acyrtosiphon pisum</i> )           | <i>Shigella flexneri</i>                  | 89.3 %                                      | Shigenobu et al., 2000       |
| <i>Parachlamydia acanthamoebae</i>                          | Y07556           | Amoebae ( <i>Acanthamoeba castellanii</i> )   | <i>Waddlia chondrophila</i> <sup>b</sup>  | 89.4 %                                      | Everett et al., 1999         |
| <i>Buchnera aphidicola</i> str. Sg                          | NC_004061        | Aphid ( <i>Schizaphis graminum</i> )          | <i>Escherichia coli</i>                   | 89.4 %                                      | Tamas et al., 2002           |
| Trophosome symbiont of <i>Riftia pachyptila</i>             | M99451           | Tube worm ( <i>Riftia pachyptila</i> )        | <i>Thiothrix nivea</i>                    | 90.0 %                                      | Distel et al., 1988          |
| Mussel methanotrophic gill symbiont MAR2                    | U29164           | Hydrothermal vent mussel                      | <i>Methylobacter luteus</i>               | 92.8 %                                      | Distel et al. 1995           |
| Endosymbiont of another bacterial endosymbiont              | AF322016         | Betaproteobacterial endosymbiont <sup>a</sup> | <i>Brenneria quercina</i>                 | 93.6 %                                      | von Dohlen et al., 2001      |
| <i>Candidatus Devosia euplatis</i>                          | AJ548825         | Ciliate ( <i>Euplotes magnicirratu</i> s)     | <i>Devosia riboflavina</i>                | 95.9 %                                      | Vannini et al., 2004         |
| Secondary endosymbiont                                      | M27040           | Aphid ( <i>Acyrtosiphon pisum</i> )           | <i>Serratia plymuthica</i>                | 97.8 %                                      | Unterman et al., 1989        |
| <i>Candidatus Burkholderia calva</i>                        | AY277697         | <i>Psychotria calva</i> (angiosperm plant)    | <i>Burkholderia</i> sp. NF100             | 97.9 %                                      | Van Oevelen et al., 2004     |
| <i>Polynucleobacter</i> sp. Bod2                            | XXX              | Ciliate ( <i>Euplotes harpa</i> )             | <i>Polynucleobacter</i> sp. MWH-Creno-4B4 | 99.5 %                                      | this study                   |
| <i>Polynucleobacter necessarius</i> ATCC 30859 <sup>T</sup> | XXX              | Ciliate ( <i>Euplotes aediculatus</i> )       | <i>Polynucleobacter</i> sp. MWH-Gad-W7    | 99.5 %                                      | this study                   |

<sup>a</sup> of mealybug (*Planococcus citri*)

<sup>b</sup> pathogenic bacterium



**Table 3.** Detection of free-living and endosymbiotic PnecC strains in the pelagic of freshwater lakes. Probe Omi10-mod is specific for the endosymbionts of *Euplotes aediculatus*. Probe PnecC-16S-445 is specific for the entire subcluster C (PnecC) of the *Polynucleobacter* cluster. This subcluster includes endosymbionts of at least two *Euplotes* species, as well as many environmental sequences, and sequences from freshwater isolates. All bacteria that hybridized with probe PnecC-16S-445 were detected outside of protist cells.

| Habitat                        | Cell numbers [ $10^4 \text{ ml}^{-1}$ ] |                   |
|--------------------------------|---|-------------------|
|                                | PnecC-16S-445                           | Omi10-mod         |
| Lake Mondsee                   | 0.4                                     | n.d. <sup>1</sup> |
| Lake Wolfgangsee <sup>2</sup>  | 0.6                                     | n.d.              |
| Lake Irrsee                    | 1.9                                     | n.d.              |
| Lake Krottensee <sup>2</sup>   | 6.6                                     | n.d.              |
| Lake Egelsee (M2) <sup>2</sup> | 8.4                                     | n.d.              |
| Pond Kleine Lacke              | 67.9                                    | n.d.              |

<sup>1</sup> n.d. - no detection

<sup>2</sup> samples from the benthic zone of these habitats were used for attempts to isolate *Euplotes* spp.

**Table 4.** Methods and conditions used in the eight cultivation experiments performed with endosymbionts from two ciliate species.

| <b>Cultivation method</b>    | <b>Initial Medium</b>           | <b>Acclimatization Medium<sup>a</sup></b> | <b>Endosymbionts of</b>                               | <b>Dilutions X Number of inoculated wells</b> |
|------------------------------|---------------------------------|---|---|---|
| Filtration, acclimatization  | IBM medium <sup>b</sup>         | NSY medium                                | <i>E. aediculatus</i> STIR1<br><i>E. harpa</i> FSP1.4 | 3 X 24<br>3 X 24                              |
| Dilution, acclimatization    | IBM medium <sup>b</sup>         | NSY medium                                | <i>E. aediculatus</i> STIR1<br><i>E. harpa</i> FSP1.4 | 3 X 24<br>3 X 24                              |
| Dilution, acclimatization    | ciliate homogenate              | NSY medium                                | <i>E. aediculatus</i> STIR1                           | 3 X 8   |
| Dilution, no acclimatization | NSY medium <sup>c</sup>         | none                                      | <i>E. aediculatus</i> STIR1<br><i>E. harpa</i> FSP1.4 | 3 X 24<br>3 X 24                              |
| Dilution, no acclimatization | ciliate homogenate <sup>c</sup> | none                                      | <i>E. aediculatus</i> STIR1                           | 6 X 4   |

<sup>a</sup> stepwise increase of concentration

<sup>b</sup> enriched with a low concentration of NSY medium

<sup>c</sup> no change of medium in the course of the experiment

## Figure Legends

**Fig. 1.** Geographic location of the sites from which the five endosymbiont bearing *Euplotes* spp. strains (*E. aediculatus*, squares; *E. harpa*, triangles), and the 40 free-living *Polynucleobacter* sp. (circles) strains affiliated with subcluster PnecC were isolated. If more than one isolate was obtained from the same region, the number of habitats (first number) and the number of strains (second number) obtained from the respective region are indicated. The five *Euplotes* spp. cultures harboring the endosymbiotic strains were each obtained from habitats located in different regions. The European area (framed in (A)) from which the majority of investigated strains were obtained is shown in (B) with a higher magnification.

**Fig. 2.** Neighbor-joining tree based on almost complete 16S rRNA sequences (*E. coli* position 39 to 1542) of endosymbiotic (highlighted in bold) and free-living strains affiliated with subcluster C (PnecC) of the *Polynucleobacter* cluster. The tree was calculated with a set of sequences including the shown 45 sequences affiliated with subcluster PnecC, as well as sequences of twelve PnecD strains, one PnecB strain and one *Ralstonia solanacearum* strain (outgroup). The geographic origin of the isolates is indicated in brackets (E, Europe; As, Asia; O, Oceania; SA, South America). Bootstrap (1,000 iterations) values of >60% are shown. The scale bar indicates 1% estimated sequence divergence.

**Fig. 3.** Neighbor-joining tree based on complete 16S-23S internal transcribed spacer (ITS) sequences (approx. 500 bp) of endosymbiotic (highlighted in bold) and free-living strains affiliated with subcluster PnecC of the *Polynucleobacter* cluster. The tree was calculated with 59 sequences of the same organisms as used for the calculation of the 16S rRNA tree (Fig. 2). All other details as described for Fig. 2.

**Fig. 4.** Base pair mismatches in helices of the 16S rRNA as suggested by a secondary structure model provided by the software package ARB (Ludwig et al. 2004). The sequences of the three *E. harpa* endosymbionts differ significantly ( $P < 0.05$ , one way ANOVA) in the number of base pair mismatches from all other groups (asterisk). Only sequences obtained by direct sequencing of the 16S rRNA gene were considered in this analysis. In total, 62 sequences were analyzed. Error bars depict standard deviation. PnecC - subcluster C, PnecD - subcluster D of the *Polynucleobacter* cluster (Hahn 2003), *E. aed.* – *E. aediculatus*, *E. harp.* – *E. harpa*.

**Fig. 5.** (A) Genome size of *Polynucleobacter* strains as determined by pulsed field gel electrophoresis. Data for endosymbiotic strains (grey bars), free-living PnecC strains (black bars), and free-living PnecD strains (hatched bars) are presented. Strains QLW-P1DMWA-1, QLW-P1FAT50D-2, and QLW-P1DNSYA-2 were isolated from the same habitat, share identical 16S rRNA and ITS sequences, as well as RAPD fingerprints. These three strains may represent the same clone. (B) Comparison of the genome size of the investigated *Polynucleobacter* strains and 22 other genome sequenced strains affiliated to the family *Burkholderiaceae*. These strains include members of the genera *Burkholderia*, *Ralstonia* and *Cupriavidus*. Average genome sizes and standard deviations are shown for the two groups. Genome size data of strains affiliated to the other five genera of the family are currently not available.

**Fig. 6.** G+C content of the entire genomes (A) and of the 16S-23S ITS (B) of endosymbiotic and free-living *Polynucleobacter* strains, as well as of other members of the family *Burkholderiaceae*. The genomic G+C content of the *P. necessarius* type strain (endosymbiont of *E. aediculatus*) was determined by Heckmann & Schmidt (1987), and the genomic G+C

content of the free-living strain was determined by genome sequencing. The genomic G+C content of 28 other *Burkholderiaceae* strains affiliated to five different genera was taken from the literature, and is presented as average and standard deviation. The analysis of the G+C content of the ITS sequences was restricted to ITS loci including tRNA-Ala and tRNA-Ile genes.

Fig. 1.




Fig. 2




Fig. 3





**Fig. 4**




Fig. 5




Fig. 6

