

Search for the Production of Narrow $t\bar{b}$ Resonances in 1.9 fb^{-1} of $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$

T. Aaltonen,²⁴ J. Adelman,¹⁴ T. Akimoto,⁵⁶ B. Álvarez González^{s,12} S. Amerio^{y,44} D. Amidei,³⁵ A. Anastassov,³⁹ A. Annovi,²⁰ J. Antos,¹⁵ G. Apollinari,¹⁸ A. Apresyan,⁴⁹ T. Arisawa,⁵⁸ A. Artikov,¹⁶ W. Ashmanskas,¹⁸ A. Attal,⁴ A. Aurisano,⁵⁴ F. Azfar,⁴³ W. Badgett,¹⁸ A. Barbaro-Galtieri,²⁹ V.E. Barnes,⁴⁹ B.A. Barnett,²⁶ P. Barria^{aa,47} V. Bartsch,³¹ G. Bauer,³³ P.-H. Beauchemin,³⁴ F. Bedeschi,⁴⁷ D. Beecher,³¹ S. Behari,²⁶ G. Bellettini^{z,47} J. Bellinger,⁶⁰ D. Benjamin,¹⁷ A. Beretvas,¹⁸ J. Beringer,²⁹ A. Bhatti,⁵¹ M. Binkley,¹⁸ D. Bisello^{y,44} I. Bizjak^{ee,31} R.E. Blair,² C. Blocker,⁷ B. Blumenfeld,²⁶ A. Bocci,¹⁷ A. Bodek,⁵⁰ V. Boisvert,⁵⁰ G. Bolla,⁴⁹ D. Bortoletto,⁴⁹ J. Boudreau,⁴⁸ A. Boveia,¹¹ B. Brau^{a,11} A. Bridgeman,²⁵ L. Brigliadori^{x,6} C. Bromberg,³⁶ E. Brubaker,¹⁴ J. Budagov,¹⁶ H.S. Budd,⁵⁰ S. Budd,²⁵ S. Burke,¹⁸ K. Burkett,¹⁸ G. Busetto^{y,44} P. Bussey,²² A. Buzatu,³⁴ K. L. Byrum,² S. Cabrera^{u,17} C. Calancha,³² M. Campanelli,³⁶ M. Campbell,³⁵ F. Canelli^{14,18} A. Canepa,⁴⁶ B. Carls,²⁵ D. Carlsmith,⁶⁰ R. Carosi,⁴⁷ S. Carrillo^{n,19} S. Carron,³⁴ B. Casal,¹² M. Casarsa,¹⁸ A. Castro^{x,6} P. Catastini^{aa,47} D. Cauz^{dd,55} V. Cavaliere^{aa,47} M. Cavalli-Sforza,⁴ A. Cerri,²⁹ L. Cerrito^{o,31} S.H. Chang,²⁸ Y.C. Chen,¹ M. Chertok,⁸ G. Chiarelli,⁴⁷ G. Chlachidze,¹⁸ F. Chlebana,¹⁸ K. Cho,²⁸ D. Chokheli,¹⁶ J.P. Chou,²³ G. Choudalakis,³³ S.H. Chuang,⁵³ K. Chung,¹³ W.H. Chung,⁶⁰ Y.S. Chung,⁵⁰ T. Chwalek,²⁷ C.I. Ciobanu,⁴⁵ M.A. Ciocci^{aa,47} A. Clark,²¹ D. Clark,⁷ G. Compostella,⁴⁴ M.E. Convery,¹⁸ J. Conway,⁸ M. Cordelli,²⁰ G. Cortiana^{y,44} C.A. Cox,⁸ D.J. Cox,⁸ F. Crescioli^{z,47} C. Cuenca Almenar^{u,8} J. Cuevas^{s,12} R. Culbertson,¹⁸ J.C. Cully,³⁵ D. Dagenhart,¹⁸ M. Datta,¹⁸ T. Davies,²² P. de Barbaro,⁵⁰ S. De Cecco,⁵² A. Deisher,²⁹ G. De Lorenzo,⁴ M. Dell'Orso^{z,47} C. Deluca,⁴ L. Demortier,⁵¹ J. Deng,¹⁷ M. Deninno,⁶ P.F. Derwent,¹⁸ P. Di Canto^{z,47} G.P. di Giovanni,⁴⁵ C. Dionisi^{cc,52} B. Di Ruzza^{dd,55} J.R. Dittmann,⁵ M. D'Onofrio,⁴ S. Donati^{z,47} P. Dong,⁹ J. Donini,⁴⁴ T. Dorigo,⁴⁴ S. Dube,⁵³ J. Efron,⁴⁰ A. Elagin,⁵⁴ R. Erbacher,⁸ D. Errede,²⁵ S. Errede,²⁵ R. Eusebi,¹⁸ H.C. Fang,²⁹ S. Farrington,⁴³ W.T. Fedorko,¹⁴ R.G. Feild,⁶¹ M. Feindt,²⁷ J.P. Fernandez,³² C. Ferrazza^{bb,47} R. Field,¹⁹ G. Flanagan,⁴⁹ R. Forrest,⁸ M.J. Frank,⁵ M. Franklin,²³ J.C. Freeman,¹⁸ I. Furic,¹⁹ M. Gallinaro,⁵² J. Galyardt,¹³ F. Garbersson,¹¹ J.E. Garcia,²¹ A.F. Garfinkel,⁴⁹ P. Garosi^{aa,47} K. Genser,¹⁸ H. Gerberich,²⁵ D. Gerdes,³⁵ A. Gessler,²⁷ S. Giagu^{cc,52} V. Giakoumopoulou,³ P. Giannetti,⁴⁷ K. Gibson,⁴⁸ J.L. Gimmell,⁵⁰ C.M. Ginsburg,¹⁸ N. Giokaris,³ M. Giordani^{dd,55} P. Giromini,²⁰ M. Giunta,⁴⁷ G. Giurgiu,²⁶ V. Glagolev,¹⁶ D. Glenzinski,¹⁸ M. Gold,³⁸ N. Goldschmidt,¹⁹ A. Golossanov,¹⁸ G. Gomez,¹² G. Gomez-Ceballos,³³ M. Goncharov,³³ O. González,³² I. Gorelov,³⁸ A.T. Goshaw,¹⁷ K. Goulianos,⁵¹ A. Gresele^{y,44} S. Grinstein,²³ C. Grosso-Pilcher,¹⁴ R.C. Group,¹⁸ U. Grundler,²⁵ J. Guimaraes da Costa,²³ Z. Gunay-Unalan,³⁶ C. Haber,²⁹ K. Hahn,³³ S.R. Hahn,¹⁸ E. Halkiadakis,⁵³ B.-Y. Han,⁵⁰ J.Y. Han,⁵⁰ F. Happacher,²⁰ K. Hara,⁵⁶ D. Hare,⁵³ M. Hare,⁵⁷ S. Harper,⁴³ R.F. Harr,⁵⁹ R.M. Harris,¹⁸ M. Hartz,⁴⁸ K. Hatakeyama,⁵¹ C. Hays,⁴³ M. Heck,²⁷ A. Heijboer,⁴⁶ J. Heinrich,⁴⁶ C. Henderson,³³ M. Herndon,⁶⁰ J. Heuser,²⁷ S. Hewamanage,⁵ D. Hidas,¹⁷ C.S. Hill^{c,11} D. Hirschbuehl,²⁷ A. Hocker,¹⁸ S. Hou,¹ M. Houlden,³⁰ S.-C. Hsu,²⁹ B.T. Huffman,⁴³ R.E. Hughes,⁴⁰ U. Husemann,⁶¹ M. Hussein,³⁶ J. Huston,³⁶ J. Incandela,¹¹ G. Introzzi,⁴⁷ M. Iori^{cc,52} A. Ivanov,⁸ E. James,¹⁸ D. Jang,¹³ B. Jayatilaka,¹⁷ E.J. Jeon,²⁸ M.K. Jha,⁶ S. Jindariani,¹⁸ W. Johnson,⁸ M. Jones,⁴⁹ K.K. Joo,²⁸ S.Y. Jun,¹³ J.E. Jung,²⁸ T.R. Junk,¹⁸ T. Kamon,⁵⁴ D. Kar,¹⁹ P.E. Karchin,⁵⁹ Y. Kato^{l,42} R. Kephart,¹⁸ J. Keung,⁴⁶ V. Khotilovich,⁵⁴ B. Kilminster,¹⁸ D.H. Kim,²⁸ H.S. Kim,²⁸ H.W. Kim,²⁸ J.E. Kim,²⁸ M.J. Kim,²⁰ S.B. Kim,²⁸ S.H. Kim,⁵⁶ Y.K. Kim,¹⁴ N. Kimura,⁵⁶ L. Kirsch,⁷ S. Klimentov,¹⁹ B. Knuteson,³³ B.R. Ko,¹⁷ K. Kondo,⁵⁸ D.J. Kong,²⁸ J. Konigsberg,¹⁹ A. Korytov,¹⁹ A.V. Kotwal,¹⁷ M. Kreps,²⁷ J. Kroll,⁴⁶ D. Krop,¹⁴ N. Krumnack,⁵ M. Kruse,¹⁷ V. Krutelyov,¹¹ T. Kubo,⁵⁶ T. Kuhr,²⁷ N.P. Kulkarni,⁵⁹ M. Kurata,⁵⁶ S. Kwang,¹⁴ A.T. Laasanen,⁴⁹ S. Lami,⁴⁷ S. Lammel,¹⁸ M. Lancaster,³¹ R.L. Lander,⁸ K. Lannon^{r,40} A. Lath,⁵³ G. Latino^{aa,47} I. Lazzizzera^{y,44} T. LeCompte,² E. Lee,⁵⁴ H.S. Lee,¹⁴ S.W. Lee^{t,54} S. Leone,⁴⁷ J.D. Lewis,¹⁸ C.-S. Lin,²⁹ J. Linacre,⁴³ M. Lindgren,¹⁸ E. Lipeles,⁴⁶ A. Lister,⁸ D.O. Litvintsev,¹⁸ C. Liu,⁴⁸ T. Liu,¹⁸ N.S. Lockyer,⁴⁶ A. Loginov,⁶¹ M. Loretz^{y,44} L. Lovas,¹⁵ D. Lucchesi^{y,44} C. Luci^{cc,52} J. Lueck,²⁷ P. Lujan,²⁹ P. Lukens,¹⁸ G. Lungu,⁵¹ L. Lyons,⁴³ J. Lys,²⁹ R. Lysak,¹⁵ D. MacQueen,³⁴ R. Madrak,¹⁸ K. Maeshima,¹⁸ K. Makhoul,³³ T. Maki,²⁴ P. Maksimovic,²⁶ S. Malde,⁴³ S. Malik,³¹ G. Manca^{e,30} A. Manousakis-Katsikakis,³ F. Margaroli,⁴⁹ C. Marino,²⁷ C.P. Marino,²⁵ A. Martin,⁶¹ V. Martin^{k,22} M. Martínez,⁴ R. Martínez-Ballarín,³² T. Maruyama,⁵⁶ P. Mastrandrea,⁵² T. Masubuchi,⁵⁶ M. Mathis,²⁶ M.E. Mattson,⁵⁹ P. Mazzanti,⁶ K.S. McFarland,⁵⁰ P. McIntyre,⁵⁴ R. McNulty^{j,30} A. Mehta,³⁰ P. Mehtala,²⁴ A. Menzione,⁴⁷ P. Merkel,⁴⁹ C. Mesropian,⁵¹ T. Miao,¹⁸ N. Miladinovic,⁷ R. Miller,³⁶ C. Mills,²³ M. Milnik,²⁷ A. Mitra,¹ G. Mitselmakher,¹⁹ H. Miyake,⁵⁶ N. Moggi,⁶ C.S. Moon,²⁸ R. Moore,¹⁸ M.J. Morello,⁴⁷

J. Morlock,²⁷ P. Movilla Fernandez,¹⁸ J. Mülmenstädt,²⁹ A. Mukherjee,¹⁸ Th. Muller,²⁷ R. Mumford,²⁶ P. Murat,¹⁸ M. Mussini^x,⁶ J. Nachtman,¹⁸ Y. Nagai,⁵⁶ A. Nagano,⁵⁶ J. Naganoma,⁵⁶ K. Nakamura,⁵⁶ I. Nakano,⁴¹ A. Napier,⁵⁷ V. Necula,¹⁷ J. Nett,⁶⁰ C. Neu^v,⁴⁶ M.S. Neubauer,²⁵ S. Neubauer,²⁷ J. Nielsen^g,²⁹ L. Nodulman,² M. Norman,¹⁰ O. Norriella,²⁵ E. Nurse,³¹ L. Oakes,⁴³ S.H. Oh,¹⁷ Y.D. Oh,²⁸ I. Oksuzian,¹⁹ T. Okusawa,⁴² R. Orava,²⁴ K. Osterberg,²⁴ S. Pagan Griso^y,⁴⁴ E. Palencia,¹⁸ V. Papadimitriou,¹⁸ A. Papaikonomou,²⁷ A.A. Paramonov,¹⁴ B. Parks,⁴⁰ S. Pashapour,³⁴ J. Patrick,¹⁸ G. Pauletta^{dd},⁵⁵ M. Paulini,¹³ C. Paus,³³ T. Peiffer,²⁷ D.E. Pellett,⁸ A. Penzo,⁵⁵ T.J. Phillips,¹⁷ G. Piacentino,⁴⁷ E. Pianori,⁴⁶ L. Pinera,¹⁹ K. Pitts,²⁵ C. Plager,⁹ L. Pondrom,⁶⁰ O. Poukhov^{*},¹⁶ N. Pounder,⁴³ F. Prakoshyn,¹⁶ A. Pronko,¹⁸ J. Proudfoot,² F. Ptohosⁱ,¹⁸ E. Pueschel,¹³ G. Punzi^z,⁴⁷ J. Pursley,⁶⁰ J. Rademacker^c,⁴³ A. Rahaman,⁴⁸ V. Ramakrishnan,⁶⁰ N. Ranjan,⁴⁹ I. Redondo,³² P. Renton,⁴³ M. Renz,²⁷ M. Rescigno,⁵² S. Richter,²⁷ F. Rimondi^x,⁶ L. Ristori,⁴⁷ A. Robson,²² T. Rodrigo,¹² T. Rodriguez,⁴⁶ E. Rogers,²⁵ S. Rolli,⁵⁷ R. Roser,¹⁸ M. Rossi,⁵⁵ R. Rossin,¹¹ P. Roy,³⁴ A. Ruiz,¹² J. Russ,¹³ V. Rusu,¹⁸ B. Rutherford,¹⁸ H. Saarikko,²⁴ A. Safonov,⁵⁴ W.K. Sakumoto,⁵⁰ O. Saltó,⁴ L. Santi^{dd},⁵⁵ S. Sarkar^{cc},⁵² L. Sartori,⁴⁷ K. Sato,¹⁸ A. Savoy-Navarro,⁴⁵ P. Schlabach,¹⁸ A. Schmidt,²⁷ E.E. Schmidt,¹⁸ M.A. Schmidt,¹⁴ M.P. Schmidt^{*},⁶¹ M. Schmitt,³⁹ T. Schwarz,⁸ L. Scodellaro,¹² A. Scribano^{aa},⁴⁷ F. Scuri,⁴⁷ A. Sedov,⁴⁹ S. Seidel,³⁸ Y. Seiya,⁴² A. Semenov,¹⁶ L. Sexton-Kennedy,¹⁸ F. Sforza^z,⁴⁷ A. Sfyrta,²⁵ S.Z. Shalhout,⁵⁹ T. Shears,³⁰ P.F. Shepard,⁴⁸ M. Shimojima^q,⁵⁶ S. Shiraishi,¹⁴ M. Shochet,¹⁴ Y. Shon,⁶⁰ I. Shreyber,³⁷ P. Sinervo,³⁴ A. Sisakyan,¹⁶ A.J. Slaughter,¹⁸ J. Slaunwhite,⁴⁰ K. Sliwa,⁵⁷ J.R. Smith,⁸ F.D. Snider,¹⁸ R. Snihur,³⁴ A. Soha,⁸ S. Somalwar,⁵³ V. Sorin,³⁶ J. Spalding,¹⁸ T. Spreitzer,³⁴ P. Squillacioti^{aa},⁴⁷ M. Stanitzki,⁶¹ R. St. Denis,²² B. Stelzer,³⁴ O. Stelzer-Chilton,³⁴ D. Stentz,³⁹ J. Strologas,³⁸ G.L. Strycker,³⁵ D. Stuart,¹¹ J.S. Suh,²⁸ A. Sukhanov,¹⁹ I. Suslov,¹⁶ T. Suzuki,⁵⁶ A. Taffard^f,²⁵ R. Takashima,⁴¹ Y. Takeuchi,⁵⁶ R. Tanaka,⁴¹ M. Tecchio,³⁵ P.K. Teng,¹ K. Terashi,⁵¹ J. Thom^h,¹⁸ A.S. Thompson,²² G.A. Thompson,²⁵ E. Thomson,⁴⁶ P. Tipton,⁶¹ P. Ttito-Guzmán,³² S. Tkaczyk,¹⁸ D. Toback,⁵⁴ S. Tokar,¹⁵ K. Tollefson,³⁶ T. Tomura,⁵⁶ D. Tonelli,¹⁸ S. Torre,²⁰ D. Torretta,¹⁸ P. Totaro^{dd},⁵⁵ S. Tourneur,⁴⁵ M. Trovato^{bb},⁴⁷ S.-Y. Tsai,¹ Y. Tu,⁴⁶ N. Turini^{aa},⁴⁷ F. Ukegawa,⁵⁶ S. Vallecorsa,²¹ N. van Remortel^b,²⁴ A. Varganov,³⁵ E. Vataga^{bb},⁴⁷ F. Vázquezⁿ,¹⁹ G. Velev,¹⁸ C. Vellidis,³ M. Vidal,³² R. Vidal,¹⁸ I. Vila,¹² R. Vilar,¹² T. Vine,³¹ M. Vogel,³⁸ I. Volobouev^t,²⁹ G. Volpi^z,⁴⁷ P. Wagner,⁴⁶ R.G. Wagner,² R.L. Wagner,¹⁸ W. Wagner^w,²⁷ J. Wagner-Kuhr,²⁷ T. Wakisaka,⁴² R. Wallny,⁹ S.M. Wang,¹ A. Warburton,³⁴ D. Waters,³¹ M. Weinberger,⁵⁴ J. Weinelt,²⁷ W.C. Wester III,¹⁸ B. Whitehouse,⁵⁷ D. Whiteson^f,⁴⁶ A.B. Wicklund,² E. Wicklund,¹⁸ S. Wilbur,¹⁴ G. Williams,³⁴ H.H. Williams,⁴⁶ P. Wilson,¹⁸ B.L. Winer,⁴⁰ P. Wittich^h,¹⁸ S. Wolbers,¹⁸ C. Wolfe,¹⁴ T. Wright,³⁵ X. Wu,²¹ F. Würthwein,¹⁰ S. Xie,³³ A. Yagil,¹⁰ K. Yamamoto,⁴² J. Yamaoka,¹⁷ U.K. Yang^p,¹⁴ Y.C. Yang,²⁸ W.M. Yao,²⁹ G.P. Yeh,¹⁸ J. Yoh,¹⁸ K. Yorita,⁵⁸ T. Yoshida^m,⁴² G.B. Yu,⁵⁰ I. Yu,²⁸ S.S. Yu,¹⁸ J.C. Yun,¹⁸ L. Zanello^{cc},⁵² A. Zanetti,⁵⁵ X. Zhang,²⁵ Y. Zheng^d,⁹ and S. Zucchelli^x,⁶
(CDF Collaboration[†])⁶²

¹*Institute of Physics, Academia Sinica, Taipei, Taiwan 11529, Republic of China*

²*Argonne National Laboratory, Argonne, Illinois 60439*

³*University of Athens, 157 71 Athens, Greece*

⁴*Institut de Fisica d'Altes Energies, Universitat Autònoma de Barcelona, E-08193, Bellaterra (Barcelona), Spain*

⁵*Baylor University, Waco, Texas 76798*

⁶*Istituto Nazionale di Fisica Nucleare Bologna, ^xUniversity of Bologna, I-40127 Bologna, Italy*

⁷*Brandeis University, Waltham, Massachusetts 02254*

⁸*University of California, Davis, Davis, California 95616*

⁹*University of California, Los Angeles, Los Angeles, California 90024*

¹⁰*University of California, San Diego, La Jolla, California 92093*

¹¹*University of California, Santa Barbara, Santa Barbara, California 93106*

¹²*Instituto de Fisica de Cantabria, CSIC-University of Cantabria, 39005 Santander, Spain*

¹³*Carnegie Mellon University, Pittsburgh, PA 15213*

¹⁴*Enrico Fermi Institute, University of Chicago, Chicago, Illinois 60637*

¹⁵*Comenius University, 842 48 Bratislava, Slovakia; Institute of Experimental Physics, 040 01 Kosice, Slovakia*

¹⁶*Joint Institute for Nuclear Research, RU-141980 Dubna, Russia*

¹⁷*Duke University, Durham, North Carolina 27708*

¹⁸*Fermi National Accelerator Laboratory, Batavia, Illinois 60510*

¹⁹*University of Florida, Gainesville, Florida 32611*

²⁰*Laboratori Nazionali di Frascati, Istituto Nazionale di Fisica Nucleare, I-00044 Frascati, Italy*

²¹*University of Geneva, CH-1211 Geneva 4, Switzerland*

²²*Glasgow University, Glasgow G12 8QQ, United Kingdom*

²³*Harvard University, Cambridge, Massachusetts 02138*

- ²⁴Division of High Energy Physics, Department of Physics,
University of Helsinki and Helsinki Institute of Physics, FIN-00014, Helsinki, Finland
- ²⁵University of Illinois, Urbana, Illinois 61801
- ²⁶The Johns Hopkins University, Baltimore, Maryland 21218
- ²⁷Institut für Experimentelle Kernphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany
- ²⁸Center for High Energy Physics: Kyungpook National University,
Daegu 702-701, Korea; Seoul National University, Seoul 151-742,
Korea; Sungkyunkwan University, Suwon 440-746,
Korea; Korea Institute of Science and Technology Information, Daejeon,
305-806, Korea; Chonnam National University, Gwangju, 500-757, Korea
- ²⁹Ernest Orlando Lawrence Berkeley National Laboratory, Berkeley, California 94720
- ³⁰University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³¹University College London, London WC1E 6BT, United Kingdom
- ³²Centro de Investigaciones Energeticas Medioambientales y Tecnologicas, E-28040 Madrid, Spain
- ³³Massachusetts Institute of Technology, Cambridge, Massachusetts 02139
- ³⁴Institute of Particle Physics: McGill University, Montréal, Québec,
Canada H3A 2T8; Simon Fraser University, Burnaby, British Columbia,
Canada V5A 1S6; University of Toronto, Toronto, Ontario,
Canada M5S 1A7; and TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
- ³⁵University of Michigan, Ann Arbor, Michigan 48109
- ³⁶Michigan State University, East Lansing, Michigan 48824
- ³⁷Institution for Theoretical and Experimental Physics, ITEP, Moscow 117259, Russia
- ³⁸University of New Mexico, Albuquerque, New Mexico 87131
- ³⁹Northwestern University, Evanston, Illinois 60208
- ⁴⁰The Ohio State University, Columbus, Ohio 43210
- ⁴¹Okayama University, Okayama 700-8530, Japan
- ⁴²Osaka City University, Osaka 588, Japan
- ⁴³University of Oxford, Oxford OX1 3RH, United Kingdom
- ⁴⁴Istituto Nazionale di Fisica Nucleare, Sezione di Padova-Trento, ^yUniversity of Padova, I-35131 Padova, Italy
- ⁴⁵LPNHE, Université Pierre et Marie Curie/IN2P3-CNRS, UMR7585, Paris, F-75252 France
- ⁴⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104
- ⁴⁷Istituto Nazionale di Fisica Nucleare Pisa, ^zUniversity of Pisa,
^{aa}University of Siena and ^{bb}Scuola Normale Superiore, I-56127 Pisa, Italy
- ⁴⁸University of Pittsburgh, Pittsburgh, Pennsylvania 15260
- ⁴⁹Purdue University, West Lafayette, Indiana 47907
- ⁵⁰University of Rochester, Rochester, New York 14627
- ⁵¹The Rockefeller University, New York, New York 10021
- ⁵²Istituto Nazionale di Fisica Nucleare, Sezione di Roma 1,
^{cc}Sapienza Università di Roma, I-00185 Roma, Italy
- ⁵³Rutgers University, Piscataway, New Jersey 08855
- ⁵⁴Texas A&M University, College Station, Texas 77843
- ⁵⁵Istituto Nazionale di Fisica Nucleare Trieste/Udine,
I-34100 Trieste, ^{dd}University of Trieste/Udine, I-33100 Udine, Italy
- ⁵⁶University of Tsukuba, Tsukuba, Ibaraki 305, Japan
- ⁵⁷Tufts University, Medford, Massachusetts 02155
- ⁵⁸Waseda University, Tokyo 169, Japan
- ⁵⁹Wayne State University, Detroit, Michigan 48201
- ⁶⁰University of Wisconsin, Madison, Wisconsin 53706
- ⁶¹Yale University, New Haven, Connecticut 06520
- ⁶²URL: <http://www-cdf.fnal.gov>

We present new limits on resonant $t\bar{b}$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, using 1.9fb^{-1} of data recorded with the CDF II detector at the Fermilab Tevatron. We reconstruct a putative $t\bar{b}$ mass in events with a lepton, neutrino candidate, and two or three jets, and search for anomalous $t\bar{b}$ production as modeled by $W' \rightarrow t\bar{b}$. We set a new limit on a right-handed W' with standard model-like coupling, excluding any mass below $800\text{GeV}/c^2$ at 95% C.L. For any narrow W' -like state with mass above $800\text{GeV}/c^2$, the cross-section is found to be less than 0.28pb at 95% C.L. We also present an exclusion of the W' coupling strength versus W' mass.

PACS numbers: 13.85.Rm; 14.65.Ha; 14.70.Pw; 12.15.Ji; 13.85.Qk

*Deceased

†With visitors from ^aUniversity of Massachusetts Amherst,

Many modifications of the standard model (SM) of particle physics include new, massive, short-lived particles with two-body decays to known fermion pairs. A classic search strategy for these states looks for resonant signals in the spectra of two-body mass distributions. Recent techniques developed to observe electroweak single-top production are well-suited to a search for unexpected $t\bar{b}$ resonances [1]. A $t\bar{b}$ resonance (inclusion of the charge conjugate is implied throughout the text) is predicted by a wide range of models containing a massive charged vector boson, generically referred to as W' . The classic model is a simple extension of the SM to the left-right symmetric group $SU(2)_L \times SU(2)_R \times U(1)$ [2], which adds a right-handed charged boson W_R with universal weak coupling strength and unknown mass. The W' may arise in models with other symmetry extensions: as the excitation of the W -boson in Kaluza-Klein extra dimensions [3], as the techni- ρ of technicolor theories [4], or as a bosonic partner in little Higgs scenarios [5].

The classic limits on W' are derived from searches in the $W' \rightarrow l\nu$ decay channel [6]. For large W' masses, the sensitivity in this channel is diminished by the broad Jacobian lineshapes for the lepton momentum and W' transverse mass. Searches in the $t\bar{b}$ channel [7] avoid this difficulty and also probe models where the couplings are free parameters and the leptonic decay modes may be suppressed. Although we quantify our results using the model of a right-handed W' with SM-like coupling [8], this analysis is sensitive to any narrow state decaying to $t\bar{b}$, including *e.g.* a charged Higgs boson or bound states arising from new dynamics in the third generation. Searches in the $t\bar{b}$ channel complement searches for neutral states coupling to $t\bar{t}$ [9].

In this Letter we present a new search for an s -channel

$W' \rightarrow t\bar{b}$ resonance produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV at the Fermilab Tevatron. The dataset of 1.9 fb^{-1} was recorded with the CDF II detector; a standard coordinate system [10] is used. Our selection is based on the leptonic decay mode $t\bar{b} \rightarrow (\ell\nu b)\bar{b}$, which has been well understood in the search for electroweak single-top production [1]. Events are expected to have a high transverse momentum (p_T) electron or muon candidate, missing transverse energy (\cancel{E}_T) from a neutrino [11], and two or three jets, at least one of which is a b -quark candidate. The dominant background is from W + jet processes and electroweak top-quark production. We reconstruct each event according to our signal hypothesis $W' \rightarrow t\bar{b} \rightarrow (\ell\nu b)\bar{b}$, then search the mass spectrum for a narrow resonance. If no signal is detected, we set limits on $\sigma(p\bar{p} \rightarrow W') \times \text{BR}(W' \rightarrow t\bar{b})$ and on the W' coupling strength $g_{W'}$.

The CDF II detector [12] is a cylindrically-symmetric general-purpose detector. Precision charged-particle tracking is accomplished by layers of silicon microstrip detectors surrounded by a large open-cell drift chamber within a 1.4 T solenoidal magnetic field. Outside the magnet are the electromagnetic and hadronic calorimeters, steel for hadronic shielding, and an exterior layer of muon detectors. The luminosity of the $p\bar{p}$ collisions is measured using gas Cherenkov detectors at small angles.

We select data using online selection criteria which require a high- p_T lepton or large \cancel{E}_T [13]. We identify $t\bar{b} \rightarrow \ell\nu b\bar{b}$ candidates as having an electron or muon with $p_T \geq 20 \text{ GeV}/c$. We also require $\cancel{E}_T \geq 25 \text{ GeV}$ and two or three hadronic jets with $p_T \geq 20 \text{ GeV}/c$ and $|\eta| \leq 2.8$. Jets are clustered in cones of fixed radius $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} \leq 0.4$, and at least one jet is required to be “ b -tagged”, *i.e.* the jet contains a secondary vertex consistent with the decay of a hadron containing a b -quark [14]. Also we exclude events consistent with cosmic rays, Z decay, and QCD multi-jet background [1].

The primary background process is the associated production of a W boson and jets with subsequent leptonic decay of the W boson (W + jets). Approximately 70% of our sample are events where the b -tag singles out W + jets events containing heavy flavor ($W + b\bar{b}$, $W + c\bar{c}$, $W + c$ + jets) or incorrectly b -tagged light flavor (mistags). We establish the normalization of these processes from data, and estimate the fraction of the candidate events with bottom or charm flavor using the ALPGEN Monte Carlo event generator [15]. The mistagging rate for light-flavor jets is estimated from generic jet data [16]. Additional backgrounds including $t\bar{t}$ pair production, s -channel and t -channel single-top production, and diboson processes are modeled using the PYTHIA Monte Carlo event generator [17] are normalized to their next-to-leading-order cross-sections predicted by theory. A small multijet background without leptonic W decay (“non- W ”) arises when a jet is misidentified as a lepton and \cancel{E}_T results from jet energy mismeasurement; this

Amherst, Massachusetts 01003, ^bUniversiteit Antwerpen, B-2610 Antwerp, Belgium, ^cUniversity of Bristol, Bristol BS8 1TL, United Kingdom, ^dChinese Academy of Sciences, Beijing 100864, China, ^eIstituto Nazionale di Fisica Nucleare, Sezione di Cagliari, 09042 Monserrato (Cagliari), Italy, ^fUniversity of California Irvine, Irvine, CA 92697, ^gUniversity of California Santa Cruz, Santa Cruz, CA 95064, ^hCornell University, Ithaca, NY 14853, ⁱUniversity of Cyprus, Nicosia CY-1678, Cyprus, ^jUniversity College Dublin, Dublin 4, Ireland, ^kUniversity of Edinburgh, Edinburgh EH9 3JZ, United Kingdom, ^lUniversity of Fukui, Fukui City, Fukui Prefecture, Japan 910-0017 ^mKinki University, Higashi-Osaka City, Japan 577-8502 ⁿUniversidad Iberoamericana, Mexico D.F., Mexico, ^oQueen Mary, University of London, London, E1 4NS, England, ^pUniversity of Manchester, Manchester M13 9PL, England, ^qNagasaki Institute of Applied Science, Nagasaki, Japan, ^rUniversity of Notre Dame, Notre Dame, IN 46556, ^sUniversity de Oviedo, E-33007 Oviedo, Spain, ^tTexas Tech University, Lubbock, TX 79609, ^uIFIC(CSIC-Universitat de Valencia), 46071 Valencia, Spain, ^vUniversity of Virginia, Charlottesville, VA 22904, ^wBergische Universität Wuppertal, 42097 Wuppertal, Germany, ^{ee}On leave from J. Stefan Institute, Ljubljana, Slovenia,

background is modeled using data. The predicted SM background is 1444 ± 255 events with two jets and 682 ± 83 events with three jets. We observe 1362 events with two jets and 617 events with three jets in data.

According to the proposed W' hypothesis, the W' mass is given by reconstructing $M_{t\bar{b}}$ from the four-momenta of the lepton, neutrino, and two jets. The unmeasured longitudinal neutrino momentum p'_z is quadratically constrained by assigning $M_{l\nu} = M_W = 80.448 \text{ GeV}/c^2$ [18]. We assign p'_z to the smallest real solution or to the real part of complex solutions [19]. The reconstructed W is then combined with the two highest E_T jets, assumed to arise from the b -quarks and corrected to reproduce parton-level energies, to form $M_{t\bar{b}}$.

Our signal model is a W' with purely right-handed decays and SM-like coupling, simulated using PYTHIA. The left-handed case is not considered since the consequent $W - W'$ interference has not been observed in any precision W measurements. Figure 1 shows the $M_{t\bar{b}}$ distribution in data superimposed with the expected signal shape for a $600 \text{ GeV}/c^2$ W' produced with a total cross-section of 9 pb ($\sim 4 \times$ the prediction for a W' with SM-like coupling [8]). The reconstructed width of the signal is dominated by resolution effects, particularly the jet energy resolution [20] and the incorrect assignment of jets from initial or final state radiation. Our test signal is therefore applicable for any W' -like object whose width is small compared to the experimental resolution. The binning is chosen so that background models have a sufficient number of entries in each bin, including the overflow bin for all values above $700 \text{ GeV}/c^2$.

Unlike single-top production, W' production is entirely an s -channel process; contributions from the t and u channels are suppressed by the large W' mass. We simulate a narrow right-handed W' with SM-like coupling and a mass between $300 \text{ GeV}/c^2$ and $950 \text{ GeV}/c^2$ in steps of $100 \text{ GeV}/c^2$ below $600 \text{ GeV}/c^2$ and steps of $50 \text{ GeV}/c^2$ above. This is the mass range to which our analysis is sensitive to changes in the signal distribution. For $M_{W'} = 800 \text{ GeV}/c^2$, our selection efficiency in the $t\bar{b}$ channel is approximately 2.8%: an excess of 10 events would correspond to a cross-section of 0.18 pb .

The branching ratios of a right-handed W' depend on whether decay to ν_R is allowed; we consider both possibilities. If leptonic decay is forbidden, as for a leptophobic W' or when $M_{W'} < M_{\nu_R}$, the $M_{t\bar{b}}$ prediction simply has a slightly larger normalization. For example, if $M_{W'} = 800 \text{ GeV}/c^2$, $\sigma \times \text{BR}(W' \rightarrow t\bar{b})$ is predicted to be 0.337 pb if leptonic decays are forbidden and 0.262 pb if they are allowed.

We set frequentist limits on $W' \rightarrow t\bar{b}$ using the measure CL_s from [22], which is defined as the probability of background plus a specified signal fraction matching the data (P_{S+B}) divided by the probability of a background-only model matching the data (P_B). Sources of uncertainty are treated using a large series of trials ($\sim 50\text{k}$) for




FIG. 1: $M_{t\bar{b}}$ for events with two jets and one b -tag, comparing the shapes between background and signal. Backgrounds are stacked and grouped according to similar shape. A $600 \text{ GeV}/c^2$ W' model is shown with $\sigma \times \text{BR}(W' \rightarrow t\bar{b}) = 9 \text{ pb}$ ($\sim 4 \times$ the prediction for a W' with SM-like coupling).

both cases. Each trial is produced by randomly varying all uncertain parameters in the model prediction within a Gaussian constraint about their nominal values. P_{S+B} is determined from the fraction of the S+B trials with a minimized $\Delta\chi^2 = \chi^2(\text{Data}|S+B) - \chi^2(\text{Data}|B)$ larger than in data; P_B is analogous. The 95% C.L. limit is set by adjusting the signal fraction assumed in the S+B model until $CL_s = 0.05$.

The systematic uncertainties arising from unknown errors in the rates and shapes of our model distributions are estimated by making reasonable variations in the model parameters and resimulating the analysis [1]. Our result is most sensitive to the jet-energy scale and the b -tagging rate at high-energy; all others sources of uncertainty are comparably small. Including all sources of uncertainty in our model increases the expected upper limit on the cross-section by 30-40%.

Applying the full limit procedure, we find no anomalous contribution in the $M_{t\bar{b}}$ spectrum. We set 95% C.L. upper limits on $\sigma \times \text{BR}(W' \rightarrow t\bar{b})$ as listed in Table 1, for a right-handed W' with SM-like coupling. Predicted cross-sections for such a W' [8] are shown in Figure 2: we set new 95% C.L. limits of $M_{W'} > 800 \text{ GeV}/c^2$ including leptonic decays, and $M_{W'} > 825 \text{ GeV}/c^2$ if leptonic decays are forbidden. The best prior result used 0.9 fb^{-1} and found $M_{W'} \geq 768 \text{ GeV}/c^2$ if leptonic decays are forbidden [7].

Although we have quantified our results using the sim-

TABLE I: 95% C.L. limits on $\sigma \times \text{BR}(W' \rightarrow t\bar{b})$ as function of $M_{W'}$ for a right-handed W' with SM-like coupling. The expected limit is quoted with the range of values into which our observation should fall 68% of the time assuming no signal is present. The cross-section for any narrow W' -like signal is less than 0.28 pb for masses above 800 GeV/ c^2 .

$M_{W'}$ (GeV/ c^2)	Expected Limit (pb)	Observed Limit (pb)
300	$1.56^{+0.62}_{-0.45}$	1.59
400	$1.04^{+0.44}_{-0.30}$	1.17
500	$0.74^{+0.35}_{-0.24}$	0.84
600	$0.54^{+0.22}_{-0.17}$	0.44
650	$0.46^{+0.21}_{-0.13}$	0.39
700	$0.40^{+0.17}_{-0.12}$	0.32
750	$0.33^{+0.15}_{-0.09}$	0.28
800	$0.30^{+0.13}_{-0.09}$	0.26
850	$0.28^{+0.13}_{-0.08}$	0.25
900	$0.28^{+0.13}_{-0.08}$	0.26
950	$0.30^{+0.13}_{-0.09}$	0.28




FIG. 2: Expected and observed 95% C.L. limits on $\sigma \times \text{BR}(W' \rightarrow t\bar{b})$ as function of $M_{W'}$ for 1.9 fb^{-1} , along with theoretical predictions. A right-handed W' with SM-like couplings is excluded for W' masses below 800 GeV/ c^2 ,

ple model of a purely right-handed W' with SM-like, this analysis is sensitive to any narrow state decaying to $t\bar{b}$. We find the cross-section for any such narrow state is less than 0.28 pb for masses above 800 GeV/ c^2 .

For a simple s -channel model with effective coupling $g_{W'}$, the cross-section is proportional to $g_{W'}^4$. Relaxing the assumption of the universal weak coupling, our cross-section limits can be rewritten as upper limits on $g_{W'}$ as a function of $M_{W'}$. The excluded region of the $g_{W'} - M_{W'}$ plane is shown in Figure 3, with $g_{W'}$ in units of g_W . At

$M_{W'} = 300 \text{ GeV}/c^2$, we limit (95% C.L.) the effective coupling to be less than 0.40 of the W boson coupling.




FIG. 3: Observed 95% C.L. limits on the coupling strength of a right-handed W' compared to the SM W boson coupling, $g_{W'}/g_W$, as function of $M_{W'}$ for 1.9 fb^{-1} . The shaded region above the dashed lines are excluded.

We thank the Fermilab staff and the technical staffs of the participating institutions for their vital contributions. This work was supported by the U.S. Department of Energy and National Science Foundation; the Italian Istituto Nazionale di Fisica Nucleare; the Ministry of Education, Culture, Sports, Science and Technology of Japan; the Natural Sciences and Engineering Research Council of Canada; the National Science Council of the Republic of China; the Swiss National Science Foundation; the A.P. Sloan Foundation; the Bundesministerium für Bildung und Forschung, Germany; the Korean Science and Engineering Foundation and the Korean Research Foundation; the Science and Technology Facilities Council and the Royal Society, UK; the Institut National de Physique Nucleaire et Physique des Particules/CNRS; the Russian Foundation for Basic Research; the Ministerio de Ciencia e Innovación, and Programa Consolider-Ingenio 2010, Spain; the Slovak R&D Agency; and the Academy of Finland.

-
- [1] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **101**, 252001 (2008).
- [2] J.C. Pati, A. Salam, Phys. Rev. D **10**, 275 (1974) [Erratum-ibid. D **11** 703 (1975)]; R.N. Mohapatra, J.C. Pati, Phys. Rev. D **11**, 566 (1975); G. Senjanovic, R.N. Mohapatra, Phys. Rev. D **12**, 1502 (1975).
- [3] Y. Mimura, S. Nandi, Phys. Lett. B **538**, 406 (2002); G. Burdman, B.A. Dobrescu, E. Ponton, Phys. Rev. D **74**, 075008 (2006).
- [4] E. Malkawi, T. Tait, C.P. Yuan, Phys. Lett. B **385**, 304 (1996); H. Georgi, E.E. Jenkins, E.H. Simmons, Nucl. Phys. B **331**, 541 (1990).
- [5] M. Perelstein, Prog. Part. Nucl. Phys. **58**, 247 (2007).
- [6] V.M. Abazov *et al.* (DØ Collaboration), Phys. Rev. Lett. **100**, 031804 (2008); A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D **75**, 091101 (2007).
- [7] V.M. Abazov *et al.* (DØ Collaboration), Phys. Rev. Lett. **100**, 211803 (2008); D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **90**, 081802 (2003).
- [8] Z. Sullivan, Phys. Rev. D **66** 075011, (2002).
- [9] T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **100**, 231801 (2008); T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. D **77**, 051102 (2008).
- [10] We use coordinates where ϕ is the azimuthal angle, θ is the polar angle with respect to the proton beam axis, transverse energy is $E_T = E \sin(\theta)$, and the pseudorapidity is $\eta = -\ln[\tan(\theta/2)]$.
- [11] Missing transverse energy, \cancel{E}_T , is defined as the magnitude of the vector $-\sum_i E_T^i \vec{n}_i$ where E_T^i are the magnitudes of transverse energy contained in each calorimeter tower i , and \vec{n}_i is the unit vector from the interaction vertex to the tower in the transverse (x, y) plane.
- [12] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 032001 (2005).
- [13] A. Abulencia *et al.* (CDF Collaboration), Phys. Rev. D **74**, 072006 (2006); T. Aaltonen *et al.* (CDF Collaboration), Phys. Rev. Lett. **100**, 211801 (2008).
- [14] We use the SECVTX algorithm described in D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 052003 (2005).
- [15] M.L. Mangano *et al.*, J. High Energy Phys. **0307**, 001 (2003).
- [16] D. Acosta *et al.*, Phys. Rev. D **71**, 052003 (2005).
- [17] T. Sjostrand *et al.*, Comput. Phys. Commun. **135**, 238 (2001).
- [18] D.E. Groom *et al.* (Particle Data Group), Eur. Phys. Jour. C **15** 1 (2000).
- [19] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. D **71**, 012005 (2005).
- [20] A. Bhatti *et al.*, Nucl. Instr. Meth. A **566**, 375 (2006).
- [21] E. Boos, V. Bunichev, L. Dudko, M. Perfilov, Phys. Lett. B **655** 245 (2007).
- [22] T. Junk, Nucl. Instr. Meth. A **434**, 435 (1999); A.L. Read, J. Phys. G **28**, 2693 (2002).