UCRL 134
~~Cy 77/1A~~
c.2

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545

BERKELEY, CALIFORNIA

UCRL - 134
c.2

Cover Sheet
Do not remove

INDEX NO. UCRL-134
This document contains 9 pages
and plates of figures.
This is copy 77 of 93 Series A
Issued to: Int. Div.

RESTRICTED
Classification

Each person who received this document must sign the cover sheet in the space below.

[illegible]

Physics-General

CLASSIFICATION
OF THE DOCUMENT
/ THE DECLASSIFIED

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

ARTIFICIAL COLLATERAL CHAINS TO THE THORIUM
AND ACTINIUM FAMILIES

A. Ghiorso, W. W. Meinke and G. T. Seaborg

Berkeley, California

Special Review of Declassified Reports
Authorized by USDOE JK Bratton
Unclassified TWX P182206Z May 79

REPORT PROPERLY DECLASSIFIED

<i>J. N. Green</i>	<i>8-16-79</i>
Authorized Derivative Classifier	Date
<i>Genia Cohen</i>	<i>8-20-79</i>
By	Date

Physics-General

- 1 -

Standard DistributionCopy Nos.

Argonne National Laboratory	1-8
Armed Forces Special Weapons Project	9
Atomic Energy Commission, Washington	10-11
Battelle Memorial Institute	12
Brookhaven National Laboratories	13-20
Carbide & Carbon Chem. Corp. (K-25 Area)	21-24
Carbide & Carbon Chem. Corp. (Y-12 Area)	25-28
Columbia University (Dunning)	29
General Electric Company	30-33
Hanford Directed Operations	34-38
Iowa State College	39
Los Alamos	40-42
Monsanto Chemical Company, Dayton	43-44
National Bureau of Standards	45-46
Naval Radiological Defense Laboratory	47
NEPA	48
New York Directed Operations	49-50
Oak Ridge National Laboratories	51-58
Patent Advisor, Washington	59
Technical Information Division, ORDO	60-74
UCLA Medical Research Laboratory, (Warren)	75
University of California Radiation Laboratory	76-80
University of Rochester	81-82
Chicago Directed Operations	83
Declassification Procedure	
Declassification Officer	84-87
Publications Officer	88
Patent Dept.	89-90
E. O. Lawrence	91
Area Manager	92
Information Division	93

Total

93

Radiation Laboratory
University of California
Berkeley, California

ARTIFICIAL COLLATERAL CHAINS TO THE THORIUM
AND ACTINIUM FAMILIES

A. Ghiorso, W. W. Meinke and G. T. Seaborg

Department of Chemistry and Radiation Laboratory,
University of California
Berkeley, California

ABSTRACT

We have produced and identified two new series of alpha-particle emitting radioactive elements; one is a "collateral" branch of the actinium ($4n + 3$) radioactive family and the other is collateral to the thorium ($4n$) family. The series are of considerable interest in that they are the first whose early members lie on the neutron deficient side of beta stability. They have been produced in high yield by irradiation of thorium with deuterons of energy about 80 Mev in the Berkeley 184-inch cyclotron. So far as the present observations are concerned both of these series begin with isotopes of protactinium (atomic number 91), although progenitors with higher atomic numbers are to be expected and will possibly be produced and identified. These protactinium isotopes are Pa^{227} and Pa^{228} formed by $d, 7n$ and $d, 6n$ reactions respectively.

Contract No. W-7405-eng-48.

To be published in the Physical Review as a Letter to the Editor.

CLASSIFICATION CANCELLED BY AUTHORITY
OF THE DISTRICT ENGINEER
BY THE DECLASSIFICATION COMMITTEE

UCRL - 134
Page 3

June 18, 1948

ARTIFICIAL COLLATERAL CHAINS TO THE THORIUM
AND ACTINIUM FAMILIES

L. Ghiorso, W. W. Heinke and G. T. Seaborg
Department of Chemistry and Radiation Laboratory,
University of California
Berkeley, California

We have produced and identified two new series of alpha-particle emitting radioactive elements; one is a "collateral" branch of the actinium ($4n + 3$) radioactive family and the other is collateral to the thorium ($4n$) family. The series are of considerable interest in that they are the first whose early members lie on the neutron deficient side of beta stability. They have been produced in high yield by irradiation of thorium with deuterons of energy about 80 Mev in the Berkeley 184-inch cyclotron. So far as the present observations are concerned both of these series begin with isotopes of protactinium (atomic number 91), although progenitors with higher atomic numbers are to be expected and will possibly be produced and identified. These protactinium isotopes are Pa^{227} and Pa^{228} formed by $d, 7n$ and $d, 6n$ reactions respectively.

After bombardments of the order of one-half to four hours duration, the metallic thorium was dissolved and the element protactinium was isolated in essentially weightless fractions.

The decay of the alpha-particles from these was measured both through the use of standard alpha-particle counting devices and also with the help of an alpha-particle pulse analyzer¹ equipped

1. See, e.g., A. Ghiorso, A. H. Jaffey, H. P. Robinson and B. Weissbourd, "An Alpha-Pulse Analyzer Apparatus", Plutonium Project Record 14B, 17.3 (1948) (to be issued).

with a fast sample-changing mechanism. Through the use of the latter, a number of alpha-particle groups were observed and their energies determined.

Prominent soon after bombardment are a number of alpha-particle groups, which decay with the 38-minute half-life of the protactinium parent. These are due to the following collateral branch of the $4n + 3$ radioactive family:

The branch which arises from orbital electron capture by Pa^{227} is not shown. The mass type was identified by observation of the characteristic energy and half-life of the $\text{Bi}^{211}(\text{AcC})$ alpha-particles, the half-life of the beta-particle-emitting $\text{Tl}^{207}(\text{AcC}')$ and the growth of 18.9-day $\text{Th}^{227}(\text{RdAc})$ as an orbital-electron-capture branching decay product of the Pa^{227} (ratio $K/\alpha \approx 0.2$). The energy obtained for these At^{215} alpha-particles

is several hundred kilovolts less than that reported² for At²¹⁵

2. B. Karlik and T. Bernert, Naturwissenschaften 32, 44 (1944).

as formed by the beta-particle branching decay of Po²¹⁵ (AcA).

Identification of members of the series was aided by a simple method of recoil collection. Recoil atoms were collected from a plate which contained the entire series in equilibrium and measurements using the plate upon which these were collected established the half-life and the energy of the Ac²²³ alpha-particles. In a "second order" recoil experiment recoils were collected from the plate containing Ac²²³ (and daughters) in order to check the half-life of the alpha-particles attributed to Bi²¹¹ (AcC) and similarly a third order recoil experiment was effected in order to isolate the beta-particle emitter and prove that it decayed with the known half-life of Tl²⁰⁷ (AcCa). The very short half-lives of the Fr²¹⁹ and At²¹⁵ were estimated and their energies identified through crude coincidence experiments using the pulse analyzer apparatus to operate the driven sweep of a cathode ray oscillograph. The measured half-lives and energies for the members of this series are summarized in Table I below.

After the decay of the above described series, a second group of alpha-particle emitters can be resolved. This second series, which decays with the 22-hour half-life of its prot-actinium parent, is a collateral branch of the 4n radioactive

family as follows:

The branch which arises from orbital electron capture of Pa^{228} and Ac^{224} is not shown. The mass type was identified through observation of the characteristic radioactive properties of the $\text{Bi}^{212}(\text{ThC})$ and its daughters, chemical identification of $\text{Bi}^{212}(\text{ThC})$, the growth of $\text{Th}^{228}(\text{RdTh})$ as an orbital-electron-capture branching decay product of the Pa^{228} (ratio $K/\alpha = \sim 50$) and the growth of $\text{Ra}^{224}(\text{ThX})$ as a similar product of the Ac^{224} (ratio $K/\alpha = \sim 10$). Of interest is the check, within about 0.15 Mev, of the energy of these At^{216} alpha-particles with the energy reported³ for At^{216} as formed by the beta-particle

3. B. Karlik and T. Bernert, *Naturwissenschaften* 31, 492 (1943).

branching decay of $\text{Po}^{216}(\text{ThA})$. The half-life of the Ac^{224} could be measured and the energy of its alpha-particles identified as the result of its collection in recoil experiments. Similarly, the half-life and alpha-particle energy of the Fr^{220} could be determined by second order recoil experiments from plates containing only Ac^{224} (and daughters). The very short half-life of At^{216} was estimated as described above and its energy could be determined by measurements on samples containing its

progenitors. The half-lives and energies are summarized in Table I. The radioactive properties of ThC (and AcC) and daughters are the accepted values taken from the literature⁴.

4. See, e.g., G. T. Seaborg, Rev. Mod. Phys. 16, 1 (1944).

Table I

Isotope	Type of Radiation	Half-life	Energy of Radiation (Mev)
²²⁷ Pa 91	α	38 min.	6.46
²²³ Ac 89	α	~ 2 min.	6.64
²¹⁹ Fr 87	α	$\sim 10^{-4}$ sec.	7.30
²¹⁵ At 85	α	$\sim 10^{-4}$ sec.	8.00
²¹¹ Bi (AcC) 83	α (99.7%)	2.16 min.	6.62
²⁰⁷ Tl (AcC) 81	β^-	4.76 min.	1.47
²⁰⁷ Pb 82	stable		
²²⁸ Pa 91	α	22 hr.	6.09
²²⁴ Ac 89	α	~ 2.5 hr.	6.17
²²⁰ Fr 87	α	~ 30 sec.	6.69
²¹⁶ At 85	α	$\sim 10^{-3}$ sec.	7.79
²¹² Bi (ThC) 83	α (34%)	60.5 min.	6.05
	β^- (66%)		2.20

Table I (cont'd)

Isotope	Type of Radiation	Half-life	Energy of Radiation (Mev)
²⁰⁸ ₈₁ Tl (ThC'')	β^-	3.1 min.	1.82
²¹² ₈₄ Po (ThC')	α	3×10^{-7} sec.	8.78
²⁰⁸ ₈₂ Pb	stable		

These data extend the information on the isotopes of protactinium, actinium, francium and astatine so that more interesting correlations⁵ of mass and atomic numbers, etc., with alpha-

5 See e.g., J. Schintlmeister, Oesterr. Chem. Zt. 41, 315 (1938); A. Berthelot, Jour. de Phys. et Rad. (VIII) 3, 17 (1942).

particle decay energies and half-lives are possible.

The cooperation of Professor R. L. Thornton, Mr. J. T. Vale, and the 184-inch cyclotron group is gratefully acknowledged.

This paper is based on work performed under Contract No. W-7405-eng-48 with ^{the} Atomic Energy Commission in connection with the Radiation Laboratory of University of California, Berkeley, Calif.