

UNIVERSITY OF  
CALIFORNIA

*Radiation  
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-589  
C.2

~~CONFIDENTIAL~~

UCRL-589  
Chemistry-General

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

**DECLASSIFIED**

EXTENSION OF ALPHA- AND BETA-DECAY SYSTEMATICS  
OF PROTACTINIUM ISOTOPES

W. Wayne Meinke and Glenn T. Seaborg

January 30, 1950

CAUTION

This document contains information affecting the National Defense of the United States. Its transmission or the disclosure of its contents in any manner to an unauthorized person is prohibited and may result in severe criminal penalties under applicable Federal laws.

Berkeley, California

DECLASSIFIED

UCRL-589

Chemistry-General

Page 2

Standard Distribution: Series A

Copy Numbers

Argonne National Laboratory	1-8
Armed Forces Special Weapons Project	9
Atomic Energy Commission, Washington	10-11
Battelle Memorial Institute	12
Brookhaven National Laboratory	13-16
Bureau of Ships	17
Carbide & Carbon Chemicals Corporation (K-25)	18-21
Carbide & Carbon Chemicals Corporation (Y-12)	22-25
Columbia University (Failla)	26
Dow Chemical Company	27
General Electric Company, Richland	28-33
Hanford Operations Office	34
Idaho Operations Office	35-36
Iowa State College	37
Knolls Atomic Power Laboratory	38-41
Los Alamos	42-44
Mallinckrodt Chemical Works	45
Massachusetts Institute of Technology (Gaudin)	46
Massachusetts Institute of Technology (Kaufmann)	47
Mound Laboratory	48-50
National Advisory Committee for Aeronautics	51
National Bureau of Standards	52-53
Naval Radiological Defense Laboratory	54
NEPA Project	55
New Brunswick Laboratory	56
New York Operations Office	57-58
North American Aviation, Inc.	59
Oak Ridge National Laboratory	60-71
Patent Branch, Washington	72
Sylvania Electric Products, Inc.	73
Technical Information Branch, ORE	74-88
UCLA Medical Research Laboratory (Warren)	89
University of California Radiation Laboratory	90-93
University of Rochester	94-95
Western Reserve University (Friedell)	96
Westinghouse Electric Corporation	97-98

Total 98

INFORMATION DIVISION  
Radiation Laboratory  
Univ. of California  
Berkeley, California

~~CONFIDENTIAL~~

EXTENSION OF ALPHA- AND BETA-DECAY SYSTEMATICS  
OF PROTACTINIUM ISOTOPES

W. Wayne Meinke and Glenn T. Seaborg  
Radiation Laboratory and Department of Chemistry  
University of California, Berkeley, California

DECLASSIFIED

January 30, 1950

As part of our program for obtaining data to extend the scope of the systematics of alpha-radioactivity<sup>1</sup> and to obtain more data pertaining to the energy surface in the heavy region of elements, we have made some pertinent new measurements on protactinium isotopes.

The heaviest isotope of protactinium hitherto reported is that of mass 234 and hence it would be of interest to know the beta-emission properties of heavier isotopes in order to tie them in with the known radioactive data in this general region. Since low-energy deuteron bombardment of U<sup>238</sup> might be expected to lead to Pa<sup>235</sup> and Pa<sup>236</sup> by (d, $\alpha$ ) and (d, $\alpha$ ) reactions and proton bombardment to Pa<sup>235</sup> by the (p, $\alpha$ ) reaction, these irradiations were made in the 60-inch cyclotron at energies of 19 and 9.5 Mev, respectively.

The protactinium was chemically separated following the bombardment of natural uranium by a procedure which involved a number of manganese dioxide cycles coupled with extractions of protactinium from aqueous into organic solvents. The manganese dioxide cycles consisted of precipitating this compound from the solution of uranium in nitric acid, followed by centrifugation, dissolution of the solid with hydroxylamine solution, dilution, and reprecipitation. The dissolved precipitate from the last cycle was acidified, salted with ammonium nitrate, and the protactinium extracted with diisopropyl ketone, several washings with salted solutions being made to insure good separation from fission products. The protactinium was then washed back into a low acidity aqueous solution and after

acidification was extracted into a benzene solution of thenoyltrifluoroacetone which forms a complex ion with the protactinium. This solution was then evaporated to dryness on a platinum counting plate leaving a weightless deposit of protactinium.

The protactinium from the 19-Mev deuteron bombardment contained  $23.7 \pm 0.5$  minute and 27-day beta-particle emitters. The cross section for the formation of the former is about  $2 \times 10^{-27} \text{ cm}^2$  on the assumption that it is formed from  $\text{U}^{238}$  while that for the latter is about  $4 \times 10^{-27} \text{ cm}^2$  on the assumption that it is due to  $\text{Pa}^{233}$  produced in the reaction  $\text{U}^{235}(\text{d}, \alpha)\text{Pa}^{233}$ . A small amount of two intermediate periods were present but these most probably can be attributed to a small amount of zirconium and niobium fission product contaminants. The 23.7-minute activity was also found in the protactinium fraction from the bombardment of  $\text{U}^{238}$  ( $\text{U}^{238}/\text{U}^{235} = 2300$ ) with 9.5-Mev protons in a yield corresponding to a cross section of about  $3 \times 10^{-29} \text{ cm}^2$ . These observations are consistent with the assignment of the 23.7-minute activity to  $\text{Pa}^{235}$ , produced in the reactions  $\text{U}^{238}(\text{d}, \alpha)\text{Pa}^{235}$  and  $\text{U}^{238}(\text{p}, \alpha)\text{Pa}^{235}$ .

An aluminum absorption curve taken on this 23.7-min. activity shows that the beta-particles have a range of about 600 mg corresponding to an energy of about 1.4 Mev according to the Feather relationship.<sup>2</sup> Gamma-rays are either absent or present in amounts too small to detect by absorption experiments.

If one closes a decay cycle involving  $\text{Pu}^{239}$ ,  $\text{Np}^{239}$ ,  $\text{Pa}^{235}$ , and  $\text{U}^{235}$  using the values 0.7 Mev for the beta-disintegration energy<sup>3</sup> of  $\text{Np}^{239}$ , 5.24 Mev for the alpha-disintegration energy<sup>3</sup> of  $\text{Pu}^{239}$  and 4.6 Mev for the estimated alpha-disintegration energy<sup>1</sup> of  $\text{Np}^{239}$ , a beta-disintegration energy of 1.34 Mev is obtained for  $\text{Pa}^{235}$ . This measurement,

therefore, gives added weight to the somewhat uncertain measured beta- and estimated alpha-disintegration energies of  $\text{Np}^{239}$ , and hence adds to the reliability of data needed for the construction of the energy surface in this region.

We also were interested in determining the partial half-life of  $\text{Pa}^{230}$  for alpha-disintegration. A half-life of some hundreds to thousands of years would be predicted from the alpha-systematics by obtaining the alpha-disintegration energy from mass number versus energy plots (5.5 Mev) and taking into account the prohibition introduced by the presence of two odd nucleons in determining its alpha-disintegration rate.<sup>1</sup> In this experiment the alpha-particles of the  $\text{Pa}^{230}$  were not measured directly but its half-life for alpha-disintegration was inferred by measuring the equilibrium amount of its alpha-disintegration product, the beta-particle emitting 29-hour  $\text{Ac}^{226}$ , on the assumption that the predicted negligible branching decay of  $\text{Ac}^{226}$  by electron capture is correct. (The half-life of  $\text{Ac}^{226}$ , originally reported as 22 hours,<sup>3</sup> has been more recently determined to be 29 hours.<sup>4</sup>)

The actinium was chemically separated from a sample containing a large amount of  $\text{Pa}^{230}$  and the  $\text{Ac}^{226}$  was identified and the amount measured through an alpha pulse analysis in which the alpha-particles of its daughters, 30-minute  $\text{Th}^{226}$  and daughters, were measured. In a bombardment of a thick thorium target with 60-Mev deuterons for 10 microampere-hours in the 184-inch cyclotron a sample of  $\text{Pa}^{230}$  corresponding to  $1.3 \times 10^6$  negative beta disintegrations/minute (as determined by alpha pulse analysis of the daughter  $\text{U}^{230}$  decay chain) was chemically isolated as described above. The actinium was chemically separated from this protactinium fraction by carrying it on lanthanum and cerium fluorides which were

redissolved and reprecipitated several times and further purified after dissolution by removing thorium through repeated zirconium phosphate precipitations. At the end of the procedure, the cerium(III) was oxidized to cerium(IV) with sodium bismuthate, the cerium(IV) removed on a zirconium phosphate precipitate, and the actinium activity precipitated with a small amount of lanthanum fluoride. After dissolution in hydrochloric acid, the soluble chloride of the carrier when evaporated on platinum gave a thin plate suitable for alpha pulse analysis.

The actinium fraction from the protactinium containing  $1.3 \times 10^6$  beta-disintegrations/minute of  $\text{Pa}^{230}$  contained 300 alpha-disintegrations/minute of  $\text{Th}^{226}$  which decayed with the 29-hour half-life of its beta-emitting  $\text{Ac}^{226}$  parent. Correcting for chemical yield (as determined by 10.0-day  $\text{Ac}^{225}$  tracer) and taking into account the fact that  $\text{Pa}^{230}$  has a half-life of 17 days<sup>5</sup> and also decays by electron capture in a ratio of 10 as compared to its decay by negative beta-particle emission,<sup>6</sup> these data lead to a partial half-life for alpha-emission of about 1400 years  $\pm$  20%, which corresponds to a prohibition by a factor of about 100 in the alpha decay, presumably due to the presence in  $\text{Pa}^{230}$  of two odd nucleons.<sup>1</sup>

It is a pleasure to acknowledge the assistance of A. Ghiorso in a number of the measurements. We also wish to express our appreciation to Dr. J. G. Hamilton, B. Rossi, T. M. Putnam, Jr., and the 60-inch cyclotron crew as well as J. Vale and the other members of the 184-inch cyclotron operating group for making the irradiations.

This work was performed under the auspices of the U. S. Atomic Energy Commission.

REFERENCES

- <sup>1</sup>I. Perlman, A. Ghiorso, and G. T. Seaborg, Phys. Rev. 77, 26 (1950).
- <sup>2</sup>N. Feather, Proc. Camb. Phil. Soc. 34, 599 (1938).
- <sup>3</sup>G. T. Seaborg and I. Perlman, Revs. Mod. Phys. 20, 585 (1948).
- <sup>4</sup>K. Street, Jr., private communication.
- <sup>5</sup>M. H. Studier and E. K. Hyde, Phys. Rev. 74, 591 (1948).
- <sup>6</sup>M. H. Studier and R. J. Bruehlman, private communication.