Johnson Creek Artificial Propagation and Enhancement Project Operations and Maintenance Program

Brood Year 2000: Johnson Creek Chinook Salmon Supplementation

Biennial Report 2000 - 2002

May 2003

DOE/BP-00004393-6

This Document should be cited as follows:

Daniel, Mitch, John Gebhards, Robert Hill, "Johnson Creek Artificial Propagation and Enhancement Project Operations and Maintenance Program; Brood Year 2000: Johnson Creek Chinook Salmon Supplementation", 2000-2002 Biennial Report, Project No. 199604300, 34 electronic pages, (BPA Report DOE/BP-00004393-6)

Bonneville Power Administration
P.O. Box 3621
Portland, OR 97208

This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the author's and do not necessarily represent the views of BPA.

BROOD YEAR 2000: JOHNSON CREEK CHINOOK SALMON SUPPLEMENTATION REPORT

JOHNSON CREEK ARTIFICIAL PROPAGATION ENHANCEMENT PROJECT OPERATIONS AND MAINTENANCE PROGRAM

Period Covered: May 2000 through April 2002

Prepared by:

Mitch Daniel,
John S. Gebhards,
and
Robert Hill

Nez Perce Tribe
Department of Fisheries Resources Management
McCall Field Office
P.O. Box 1942
McCall, Idaho 83638

Prepared for:

U.S. Department of Energy
Bonneville Power Administration
Division of Fish and Wildlife
P.O. Box 3621
Portland, Oregon 97208-3621

BPA Project Number: 199604300
BPA Contract Number: 4393

May 2003

TABLE OF CONTENTS

LIST OF TABLES	iii
LIST OF FIGURES	iii
LIST OF APPENDICES	iii
ABSTRACT	1
AWKNOWLEDGEMENTS	1
INTRODUCTION	2
BACKGROUND	2
DESCRIPTION OF PROJECT AREA	3
METHODS AND MATERIALS	5
Adult Take Estimate	5
Trapping Operations	5
Fish Transport and Adult Holding	6
Spawning	6
Nutrient Enhancement	7
Egg Incubation and Rearing	7
Tagging	8
Acclimation and Release	8
RESULTS	10
Adult Take Estimate	10
Adult Salmon Trapping	10
Run Timing, Daily Water Flow and Water Temperature Measurements	11
Tag Detections	12
Length/Age Frequencies	13
Injuries and Pre-spawning Mortality	14

Adult Spawning	14
Fecundity.....	15
Disease Testing	17
Nutrient Enhancement	17
Incubation and Green Egg to Eyed Egg Survival	17
Rearing.....	18
Tagging	18
Smolt Release.....	19
CONCLUSIONS.....	20
LITERATURE CITED	21
APPENDICIES	23

LIST OF TABLES

Table 1: Adult Chinook Trap Numbers by Age*, Sex, and Disposition	10
Table 2: PIT Tag Detections at the Johnson Creek Weir in 2000	12
Table 3: Spawning and Egg Incubation Results	16

LIST OF FIGURES

Figure 1: Map of Johnson Creek Watershed and JCAPE Sites	4
Figure 2: Fish Retained in Proportion to Fish Trapped	11
Figure 3: Run Timing in Relation to Daily Average Water Flow and Temperature	12
Figure 4: Fork Length Frequency of Adult Chinook Trapped in Johnson Creek.....	13

LIST OF APPENDICES

Appendix A: Adult Trapping Summary	24
Appendix B: Hatchery Production Summary For Johnson Creek Brood year 2000 (January, 2001 – March, 2002).....	29

ABSTRACT

The Nez Perce Tribe, through funding provided by the Bonneville Power Administration, has implemented a small scale chinook salmon supplementation program on Johnson Creek, a tributary in the South Fork of the Salmon River, Idaho. The Johnson Creek Artificial Propagation Enhancement project was established to enhance the number of threatened Snake River summer chinook salmon (*Oncorhynchus tshawytscha*) returning to Johnson Creek through artificial propagation.

Adult chinook salmon trapping, broodstock selection, and spawning was first implemented in 1998, did not occur in 1999, and was resumed in 2000. A total of 152 salmon were trapped in Johnson Creek in 2000, of which 73 (25 males, 16 females, and 32 jacks) fish were transported to Idaho Fish and Game's South Fork Salmon River adult holding and spawning facility for artificial propagation purposes. The remaining 79 (29 males, 16 females, and 24 jacks) fish were released above the weir to spawn naturally. A total of 65,060 green eggs were taken from 16 female salmon and transported to the McCall Fish Hatchery for incubation and rearing. Egg counts indicated an average eye-up rate of 86.0% for 55,971 eyed eggs. Average fecundity for Johnson Creek females was 4,066 eggs per female. Juvenile fish were reared indoors at the McCall Fish Hatchery through November 2001. These fish were transferred to outdoor rearing facilities in December 2001 where they remained until release in March 2002. All of these fish were marked with Coded Wire Tags and Visual Implant Elastomer tags. In addition 9,987 were also PIT tagged. Hand counts provided by marking crews were used to amend the number of juvenile salmon released from the original egg count. A total of 57,392 smolts were released into a temporary acclimation channel in Johnson Creek on March 18, 19, 20, 2002. These fish were held in this facility until a fish screen was removed on March 22, 2002 and the fish were allowed to emigrate.

AWKNOWLEDGEMENTS

The Johnson Creek Artificial Propagation Enhancement Project would like to thank all of the people and Agencies that contributed to the success of this project. Jaime Pinkham, Ed Larson, Dave Johnson, and Becky Ashe; Doug Nelson, Carl East, Sarah Aavedal, Ryan Jain, JR Inglis, and Jason Vogel from the Nez Perce Tribe; Ken Kirkman from the Bonneville Power Administration, which provided the funding for this project; the Northwest Power Planning Council; Chris Beasley from Columbia River Inter-tribal Fisheries Commission, who assisted in permit writing; Dan Herrig and his staff at the USFWS, Lower Snake Compensation Office; Gene McPherson and his staff at the McCall Fish Hatchery, who did most of the rearing of the JCAPE fish; Rodney Duke and his crews for CWT and VIE tagging; Herb Pollard from the National Marine Fisheries Service; Doug Munson, Doug Burton, and Tom Rogers from Idaho Fish and Game; and the many other individuals whom we cannot remember at this time that assisted on this project either directly or indirectly.

INTRODUCTION

This brood year report details the methods and results from supplementation activities associated with the Johnson Creek Artificial Propagation Enhancement (JCAPE) project for the 2000 brood year. Methods and results are divided into several categories. These include adult trapping, holding and spawning, incubation, rearing, tagging, and release of the brood year 2000 juveniles as smolts into Johnson Creek. Monitoring and Evaluation (M&E) activities associated with the JCAPE project for the 2000 juvenile and adult brood year are detailed in separate reports.

BACKGROUND

The Johnson Creek summer chinook salmon (*Oncorhynchus tshawytscha*) population has experienced significant decline in population numbers over the past five decades. Escapement levels in Johnson Creek have declined from a recorded high of 486 redds in 1960 to a low of five redds observed in 1995. Since Snake River spring/summer chinook salmon were listed as a threatened species under the Endangered Species Act (ESA) we are in an emergency situation. Unprecedented efforts were needed to prevent species extinction and preserve fish for the future. Artificial propagation programs are one measure to attempt to enhance populations and increase natural production in Snake River tributaries. The National Marine Fisheries Service (NMFS) draft recovery plan (NMFS 1995) states that "supplementation programs should be initiated and/or continued for populations identified as being at imminent risk of extinction, facing severe inbreeding depression, or facing demographic risks".

Due to critically low abundance of summer chinook salmon in Johnson Creek, the Nez Perce Tribe (NPT), through funding provided by the Bonneville Power Administration (BPA), initiated the development of an artificial propagation enhancement project for Johnson Creek in 1996. This decision resulted from a number of factors including: increased emphasis on wild/natural production and stock recovery; consultation and requirements resulting from listing of Snake River chinook populations as threatened under the ESA; and preferred strategies for use of artificial propagation identified in *Wy-Kan-Ush-Mi Wa-Kush-Wit, Spirit of the Salmon* (CRITFC 1996).

The Nez Perce Tribe, as the lead fisheries management agency for the JCAPE project, submitted a Section 10 Permit Application (Lothrop 2000) under the ESA to the NMFS for the purpose of supplementing the Johnson Creek chinook salmon population in 2000. NMFS gave approval for the JCAPE project to trap returning adult salmon in Johnson Creek for the purpose of broodstock collection and subsequent rearing and release of their progeny back into Johnson Creek. The JCAPE project was allowed to rear up to 100,000 smolts per year through this approval.

The United States Fish and Wildlife Service, Lower Snake River Compensation Plan Office (LSRCP) and the Idaho Department of Fish and Game (IDFG), McCall Fish Hatchery were approached about utilizing existing rearing facilities at the McCall Fish Hatchery for the JCAPE project for the 2000 brood year. The LSRCP (owner) and IDFG (operator) of the McCall Fish Hatchery agreed to allow the JCAPE project to incubate and rear up to 100,000 brood year 2000 Johnson Creek fish at the existing McCall Fish Hatchery. In addition, the JCAPE project was allowed to hold and spawn Johnson Creek adult broodstock at the South Fork Salmon River (SFSR) adult trapping and spawning facility that is associated with the McCall Fish Hatchery. The approval by NMFS and the ability to utilize existing facilities at the McCall Fish Hatchery allowed the JCAPE project to collect broodstock in 2000.

DESCRIPTION OF PROJECT AREA

Johnson Creek begins near Deadwood Summit at an elevation of about 7,200 feet and flows northward approximately 35 miles to its confluence with the East Fork South Fork Salmon River near the community of Yellowpine at an elevation of 4,500 feet (Figure 1). Elevations range from 4,500 to 8,500 feet. The sub-basin area is 153,800 acres, primarily National Forest land with some private inholdings. Over half of the sub-basin is inventoried roadless area. The predominant vegetation is mixed conifer forest with interspersed grass and sedge meadows. The upper one-third of the Johnson Creek sub-basin has extensive meadow areas, the middle one-third is characterized by steep cascading sections, and the lower one-third of the sub-basin is lower gradient and located in a wide-bottomed valley. Most of the historic and current chinook salmon spawning and rearing has been observed in this lower section (NOAA 2004 draft). Runoff is characterized by high flows in the spring followed by a decreasing hydrograph with the lowest flows typically in early fall through late winter. Mean annual precipitation varies from between 15 and 58 inches. Most of this precipitation falls in the form of snow from October through April. Summer temperatures can reach near 100 at the lowest elevations, while winter lows are often well below zero (USFS 1995, USFS 1990).

Figure 1: Map of Johnson Creek Watershed and JCAPE Sites

METHODS AND MATERIALS

Adult Take Estimate

The JCAPE project was given approval to rear up to 100,000 Johnson Creek smolts from the 2000 brood year. This level of production would require eggs from up to 32 females. Broodstock for the JCAPE program is collected over the spectrum of the run and across age classes. In order to prevent from biasing one age class or portion of the run, it is necessary to set an adult take level that will meet broodstock and natural spawning needs. This is best done using an adult escapement estimate to set the percentage of fish to be retained as broodstock. In 2000, the adult escapement estimate was based on the number smolts produced from redds in Johnson Creek in 1996 and applying a range of smolt to adult survival rates to the number of smolts produced.

Trapping Operations

Adult salmon trapping activities occur in Johnson Creek approximately five miles upstream of the confluence of the EFSFSR (see Figure 1) at an elevation of 4,889 ft. The weir is located on privately owned property of the Bryant Ranch, and is used by agreement with the owners of the property. The weir is installed as soon water flows are low enough (approximately 700 cfs) to allow for safe installation and operated through mid September until no fish have been trapped for ten (10) consecutive days.

A picket style weir is installed, which consists of a guide fence and an in-stream trap box. The guide fence blocks upstream passage of adult chinook salmon and funnels them into the trap box. The trap box consists of an aluminum framework filled with the same type of pickets as used in the guide fence. The dimensions of the box are approximately 8' x 8' x 8'. The entrance into the trap box was redesigned following the 1998 trapping season during which fish were observed escaping down river out of the trap box through the opening. The entrance was extended upstream with a smaller opening. The trap box is covered with plywood for shade cover and to prevent the possibility of fish jumping out of the trap box. Camouflage burlap is added to the trap box entrance for additional shade cover.

Every day the trap is checked for the presence of adult salmon. Individual fish are netted out of the trap and placed in an anesthetizing tub. Each fish is anesthetized with Tricane Methane Sulfonate (MS-222) at a concentration of 90 mg/l in 30 gallons of water. Once anesthetized, fish are visually and electronically scanned for tags and marks, fork length and mid eye to hyperle length (MEH) are recorded, a sequentially numbered opercle tag is applied, and a paper punch sample of tissue is taken from the caudal fin for genetic analysis.

Adult chinook salmon are retained for broodstock based on age and sex. Age class determination is based on length frequency data from SFSR. Adult salmon that are to be retained as broodstock are given an additional opercle punch and a numbered floy tag corresponding to the numbered opercle tag. Floy tags are used as a back up in case the opercle tag is lost. Each adult retained for broodstock, excluding jacks, is inoculated for bacterial kidney disease with 0.3cc erythromycin.

Fish released above the weir to spawn naturally are held in fresh water upstream of the weir until equilibrium is regained. They are then released and observed until swimming out of sight. Fish released upstream are not injected with erythromycin.

Fish Transport and Adult Holding

Adult chinook retained as broodstock from Johnson Creek are held and spawned at the SFSR adult holding facility (see Figure 1). This satellite facility of the McCall Fish Hatchery is located on the west side of the SFSR, approximately three miles west of Warm Lake, Idaho at an elevation of 5,069 ft. The JCAPE project is allowed to use this facility through an agreement with USFWS, and the IDFG - McCall Fish Hatchery staff.

Fish are transported in an aluminum transport tank mounted on a one-ton flatbed truck. The tank has a capacity of approximately 400 gallons and is equipped with an aerator and a supplemental oxygen supply system. The tank is filled with river water prior to fish work up, and supplied with oxygen at approximately 8 liters per minute. Fish are carried gently in a net or by hand approximately 30 yards to the transport vehicle and placed in the tank. Once the trap is emptied, fish are transported about 35 miles (1.25 hours) to the SFSR facility. They are then netted out of the tank and placed in the raceways with the SFSR fish. The SFSR facility has two raceways, each 90 feet long, 10 feet wide, and 3.5 feet deep, with approximately 3,150 cubic feet of water. Males and females are held separately.

Johnson Creek males and jacks are held in the SFSR male pond until the first spawning sort in early August. At that time they are moved to a temporary 16 foot diameter, 5 foot deep, circular tank. An electric pump supplies the temporary tank with water from the SFSR fish ladder. High and low water alarms are installed in the tank. The tank is covered to provide shade and keep the fish from jumping out. The covers are made of four PVC framed panels, each four feet wide and constructed to approximately match the curvature of the tank. The PVC frames are covered with welded wire fence and then wrapped with plastic tarps. The fence and tarps are attached to the PVC with wire ties and the panels were tied to the tank. Stairs and a work area are positioned next to the tank to provide safe and easy access to the tank. Johnson Creek females remain in the SFSR female pond throughout the spawning season.

Spawning

Adult salmon held at the SFSR facility are typically spawned two (2) days a week from mid August through the first week of September. Prior to spawning each day, a crowd rack is placed in the upper end of the female pond with no fish above it. All SFSR and Johnson Creek females are then crowded from the bottom end of the pond up to the first crowd rack. Individual fish are caught by hand and checked for ripeness, green (un-ripe) females are placed downstream of the second crowd rack and ripe SFSR females are placed upstream of the first crowd rack. Ripe Johnson Creek females are counted and placed in the spawning shed female live-well.

Once the female salmon have been sorted for ripeness, more than twice as many Johnson Creek males than the number of ripe Johnson Creek females are carried gently approximately 20 yards from the circular holding tank to the spawning shed male live-well. Spawning equipment, such as buckets and strainers, are prepared. The females are double checked for ripeness, euthanized with a blow to the head, placed horizontally on a table and rinsed off. A spawning knife is inserted in the females vent prior to picking her up to prevent eggs from discharging. The female

is then picked up and held vertically over a colander, the abdomen is cut to release the eggs. Ovarian fluid is collected at this time for disease testing and the carcass is set aside for further disease sampling. The eggs are then divided equally into the number of colanders matching the number of males to be used. Sperm is then applied, one male per colander. The individual colanders of eggs are placed in separate buckets of fresh water to activate the sperm. After one minute, the eggs of an individual female are combined into one colander and placed back into fresh water for another few minutes to insure fertilization. The eggs are then drained and poured gently into a bucket of water and Argentine at 200 ppm and allowed to water harden for one hour. Ice is added to the water hardening buckets to maintain temperature. After one hour of water hardening, the eggs are rinsed, poured into a numbered egg tube and then stacked in a cooler for the transport. Eggs are driven approximately 55 miles (one hour) to the McCall Fish Hatchery.

Nutrient Enhancement

Carcasses from adult fish spawned artificially are disposed of back into Johnson Creek for decomposition. Natural spawning areas are chosen as disposition points to stimulate natural nutrient enhancement to the stream. Scales and fin ray samples are taken from carcasses to provide specific age information on the Johnson Creek stock. Tails are removed to indicate fish used for artificial production rather than natural spawners. Any carcasses due to mortality other than spawning activities are disposed of in the same manner.

Egg Incubation and Rearing

Eggs collected from Johnson Creek adults are transported to the McCall Fish Hatchery for incubation and subsequent rearing. The McCall Fish Hatchery is located on the north side of the North Fork of the Payette River, in the town of McCall, Idaho at an elevation of 4,971 ft.

Eggs from each individual Johnson Creek female are placed in a single tray of an 8-tray vertical stack incubator and that position is recorded for future reference. Eggs are treated daily with formalin at 1,667 parts per million (ppm) to control fungus. Treatments start two days after the eggs go into the incubators and are continued until the eggs hatch.

Once eggs have absorbed 550 daily temperature units (DTU) (Piper et al, 1982), they are examined to verify they are at a strong eyed stage. If a strong eye development is present, that lot of eggs is shocked. Egg shocking is a process in which an undeveloped or dead egg's fragile inner membrane is broken causing the egg to turn opaque. A live or healthy egg is quite sturdy once it is eyed and will not be hurt from the shocking process. This process consists of pouring a tray of eggs into a watered up trough, siphoning the eggs through a length of garden hose into a bucket, and returning the eggs to the tray and incubator until the next day when they will be picked.

McCall Hatchery personnel separate dead eggs from live eggs and enumerate them with an automatic egg sorter and counter. The egg counter picks eggs up into a wheel that passes them through a light beam. If the light beam does not pass through the dead or opaque egg, a burst of air rejects that egg. The remaining or live eggs pass to the other side of the wheel where a constant air jet ejects the live eggs. As the eggs are ejected, they pass through a counting eye that enumerates both the live and dead eggs. The live eggs are then returned to the incubators.

The eggs remain in the incubators through the sac fry life stage. Chinook salmon eggs generally require 1,000 DTU to hatch. Sac fry generally absorb their yolk sacs at approximately 1,750 DTU. McCall Fish Hatchery protocol calls for fish to be put into early rearing vats once yolk sacs are absorbed and started on feed a few days later.

The early rearing vats, or raceways, are 40' x 4' x 2' (water depth) with 320 cubic feet of rearing space. The vats are set up at half length, or 20' x 4' x 2', with a water flow of 80 gallons per minute (gpm), when the fry are first set out into the vats. Once the fry start feeding aggressively, cleaning baffles are installed. The baffles are reinforced aluminum plates installed into the vats every four feet. They fit into the vat at a slight upstream angle and are two inches off the bottom of the vat. This forces all the water to flow through the bottom two inches, greatly increasing the flow along the bottom and flushing fecal material and uneaten food down through the raceway. The vats are extended to full length and flow is increased to 130 gpm as fish densities increase. The fry remain in the vats until large enough to mark with visual implant elastomer (VIE) tags before being transferred outside to the collection basin. Fish are tagged and transferred to the collection basin when they reach at least 35.0 fish per pound (fpp).

The collection basin is 101' x 15' x 4' (water depth) and straddles the tail end of McCall Fish Hatchery two outdoor rearing ponds. The water from both outdoor rearing ponds flows through the collection basin at a flow of 16 cubic feet per second (cfs). The fish remain in the collection basin until outplanted as smolts in mid March.

Tagging

Johnson Creek fish are marked 100% with coded wire tags (CWT) and VIE tags. CWT tags are injected into the snout and VIE tags are inserted in the adipose tissue behind the eye following standard tagging protocols. CWT and VIE tagging is performed by the IDFG marking crew on two separate occasions. Crews mark all fish and are therefore able to enumerate all the fish at that time. These hand counts are used to update fish production numbers initially provided by the automatic egg counter.

A portion of these fish are also PIT tagged prior to release. PIT tagging is conducted by NPT personnel. Fish are PIT tagged using the protocol described in Nelson and Vogel (1999). During PIT tagging, NPT personnel also conduct a tag retention study of CWT and VIE tags. The retention study involves an individual looking only for CWT tags and another looking only for VIE tags. Tag retention study protocols are described in Nelson and Vogel (1999).

Acclimation and Release

Johnson Creek smolts are transported back to Johnson Creek for release at the end of March or the beginning of April depending on weather and road conditions. Fish are netted out of the collection basin and placed in transport tanks mounted on one-ton flatbed trucks. The tanks have a capacity of approximately 400 gallons and are equipped with an aerator and a supplemental oxygen supply system. Fish are loaded at 300 pounds per tank.

The 1998 brood year was directly released into Johnson Creek. These fish emigrated from Johnson Creek within three days. It was felt that future smolt releases needed to acclimate on Johnson Creek water for a longer period of time. The 2000 brood year fish will be released into a temporary acclimation channel adjacent to Johnson Creek. The acclimation site is a natural

side channel of Johnson Creek and is located just north of the Cox Ranch, seven (7) river miles above the confluence of Johnson Creek and the EFSFSR. The downstream end of the side channel is screened off using metal fence posts and ¼ inch plastic hardware cloth to retard downstream migration of the smolts. Fish are free to move out of the upstream end of the channel as they please. The L-shaped side channel is approximately 120 feet long on the river side and 240 feet on the side closest to the road. The inlet and outlet of the side channel is approximately 50 feet and 75 feet wide, respectively, with varying widths in between. Water depth is approximately six (6) inches at the inlet and four (4) inches at the outlet. The side channel also has several pool areas that allow for holding and resting areas for the smolts. The hardware cloth fence is removed after the fish have been in the pond for 2 to 3 days, allowing the fish to emigrate out of Johnson Creek.

RESULTS

Adult Take Estimate

In 2000, the JCAPE project needed up to 32 adult females to produce the 100,000 Johnson Creek smolts from the 2000 brood year. In order to establish a percentage of the adult run to retain for broodstock, an estimate of adult escapement into Johnson Creek was needed. Adult escapement to Johnson Creek for 2000 was estimated to be between 75 and 150 fish. Assuming an equal sex ratio for the number of returning adults, it was determined that every other adult, by age and sex, would be retained and transported to SFSR adult holding facility for artificial propagation purposes.

Adult Salmon Trapping

Trapping of adult salmon in Johnson Creek occurred for only the second season in 2000. The first trapping season occurred in 1998 when a total of 114 adult summer chinook salmon were trapped. Trapping activities were not conducted in 1999 because a Section 10 permit was not issued by NMFS authorizing direct take/collection of an ESA listed fish, extremely low adult returns, and high water flows would have caused late weir installation, possibly biasing brood stock collection towards the later portion of the run.

The trap was installed in Johnson Creek on June 26, 2000 and removed on September 13, 2000. The first fish was trapped June 29, 2000 and the last fish was trapped September 6, 2000. A total of 152 adult summer chinook were trapped (Appendix A). Of these, 79 fish were released for natural spawning and 73 fish were retained for broodstock (Table 1). Figure 2 depicts the proportion of fish retained for broodstock versus the proportion of fish trapped at the weir over the course of the run. This illustrates that a good cross section of the run was retained for broodstock and was not biased towards any one portion of the adult return. Overall, 48% of the Johnson Creek fish trapped were retained for brood stock.

Mountain Whitefish (*Prosopium williamsoni*), Cutthroat trout (*oncorhynchus clarki*), and Bull trout (*salvelinus confluentus*) were the only other species trapped in the adult trap. The 18 Bull trout trapped ranged in fork length from 39 to 51 cm. Lengths were not taken on the Mountain Whitefish and the Cutthroat were not enumerated.

Table 1: Adult Chinook Trap Numbers by Age*, Sex, and Disposition

Disposition	Age 3	Age 4		Age 5		Total
	Jacks	Males	Females	Males	Females	
Retained	32	22	14	3	2	73
Released	34	24	15 ¹	5	1	79
Total Trapped	66	46	29	8	3	152

* Age 3 < 67 cm, Age 4 = 67cm to 82cm, Age 5 > 82cm.

1 - female trap mortality added to four year old females released.

Figure 2: Fish Retained in Proportion to Fish Trapped

At the time of trapping, caudal fin punches were collected for genetic sampling from adult salmon. A total of 151 fish were sampled at the weir. These samples were turned over to the JCAPE Monitoring and Evaluation program. The samples have been processed; however, the results have not been reviewed and made available at this time.

Run Timing, Daily Water Flow and Water Temperature Measurements

The Johnson Creek adult Chinook salmon return peaked on August 22, 2000. The peak coincided with a drop in water temperature and low water flows, which is similar to the peak in that occurred in 1998 (Daniel and Gebhards 2003). Of 152 fish trapped, 100 (66%) were trapped after August 1. This peak is only a measure of when the fish entered the trap, however, and does not reflect escapement prior to trap installation or fish that spawned below the weir. One mile of Johnson Creek above the weir was snorkeled the day after the trap was installed and two adult chinook were observed. JCAPE Monitoring and Evaluation personnel found 15 unmarked adult chinook carcasses upstream of the Johnson Creek weir in 2000. From this, we know that there were fish upriver of the trap prior to installation. We suspect that these fish passed prior to weir installation.

There appears to be a relationship of water flow, water temperature and when fish moved into the trap. As water flow decreased the number of fish entering the trap increased (Figure 3). Water temperature also seemed to influence adult escapement. There was a sharp increase in the number of fish entering the trap coinciding with a noticeable drop in mean daily water temperature (Figure 3). As in 1998, the largest number of fish trapped in a single day in 2000, coincided with the coldest mean daily water temperature in August. This information will be used to monitor changes in run timing and its relationship to river flows and water temperature as the JCAPE project is implemented.

Figure 3: Run Timing in Relation to Daily Average Water Flow and Temperature

Tag Detections

Each adult chinook trapped is thoroughly checked for tags and marks. The only tags detected in 2000 were PIT tags (Table 2). No CWT, VIE, radio tags, or jaw tags were detected. All of the PIT tags detected in jacks.

Table 2: PIT Tag Detections at the Johnson Creek Weir in 2000

Tag Code	Detection Date	Tag Agency	Tag Date	Tagging Location
52186C2D11	7/11/2000	NPT	9/23/1998	Johnson Cr. Screw Trap
5223004156	8/17/2000	NPT	3/17/1999	Johnson Cr. Screw Trap
53001D4F1D	8/22/2000	NMFS	5/4/1999	Lower Granite Dam

The only marks detected at the trap were adipose fin clips. Several of these fish were from an adult outplant into the Upper EFSFSR. On 7/20/00, IDFG outplanted 100 male and 100 female excess ad-clipped SFSR stock adult chinook into the EFSFSR. Three sites were chosen near the Stibnite mine and Meadow Creek, approximately 12 miles above the confluence with Johnson Creek. Five (5) of these fish, all males, migrated down the EFSFSR and then up Johnson Creek and into the trap. Two were trapped on August 22, 2000, one each on August 25, and 30, 2000, and September 1, 2000. All five were released above the weir with one dying immediately upstream and the other four were not recovered.

Length/Age Frequencies

Of the 152 fish trapped in 2000, 66, (44%) were Age 3 (one ocean) or jacks, 75 (49%) were Age 4 (2 ocean), and only 11 (7%) were Age 5 (3 ocean) (Figure 4). These age class determinations are based solely on length measurements of each fish and may not truly represent the ages of the fish sampled. Age distribution used for the Johnson Creek was based on historic information from IDFG on SFSR adult chinook returns; Age 3 < 67cm, Age 4 = 67cm to 82cm, Age 5 > 82cm. The JCAPE M&E program will use this and future age class composition information to compare and reevaluate age class determinations.

Figure 4: Fork Length Frequency of Adult Chinook Trapped in Johnson Creek

Injuries and Pre-spawning Mortality

Injury to adult chinook salmon sustained during the trapping and holding was minimal, but included jaw and snout damage incurred while being held in the trap box. No direct mortalities occurred during holding in the trap box or during transport. There was one age 4 female that gilled herself between two pickets of the weir. JCAPE personnel unsuccessfully attempted to revive her. There were no other mortalities associated with the weir in 2000.

Females that were retained for but did not contribute to broodstock were considered pre-spawn mortalities. There were no female pre-spawn mortalities in 2000. Adult males and jacks that did not contribute to broodstock and died within two weeks after the first spawn are considered pre-spawn mortalities. Of 25 adult males and 32 jacks retained as broodstock, one (1) adult male and two (2) jacks were considered pre-spawn mortalities for a combined pre-spawn mortality rate of 5.3%.

It was felt that excessive handling stress occurred during the 1998 spawning season when Johnson Creek and SFSR males were held together. Most of the SFSR males had to be handled each time Johnson Creek males were needed for broodstock. By holding Johnson Creek males separately, it was hoped that Johnson Creek broodstock would be readily available and handling stress would be reduced. The temporary circular holding tank at the SFSR facility was an improvement over holding Johnson Creek males with SFSR males during spawning. Problems were observed with the cover placed over the tank to provide shade, security, and serve as a jump screen. Seven (7) mortalities occurred when some of the small fish (jacks) jumped through holes in the tank cover. These holes were repaired and the cover was reinforced to eliminate this problem.

Another problem we experienced with the holding tank was a mechanical malfunction when a downspout fitting in the tank separated from the intake pipe and flooded the top of the covers causing them to collapse. In order to reattach the downspout, the water was turned off to the tank. While the pipe glue was drying several of the adult males and jacks began showing signs of stress due to lack of oxygen. The fish were instantly transferred into the SFSR adult female pond. A total of six males and seven jacks died from this stress. It is believed that the oxygen levels in the tank were much lower than anticipated without the fresh flow of water. All of the adult males that died had been used in spawning the previous day. We believe the combination of stresses from spawning and low oxygen levels were too much for these fish. The surviving adult males and jacks were returned to the tank the following spawn day.

Adult Spawning

Spawn timing for the fish retained as broodstock coincided with that of natural spawners in Johnson Creek, occurring from mid-August to about the first week in September. NPT and IDFG personnel worked cooperatively to spawn both Johnson Creek and SFSR fish. A total of 16 females were spawned in 2000. Fourteen females were spawned with two males each, one female with three males and another female with four males (Table 3). The 16 females were spawned on five different days from August 15 to September 1, 2000. A total of 23 adult males and two (2) jacks were used for spawning. Ten (10) adult males were used more than once. Although 32 jacks were retained for broodstock, the JCAPE spawning protocol (NPT and PRRG 2001) calls for use of about 3% of the jacks ponded. In addition, in an attempt to not overwhelm the remaining natural spawning adults in Johnson Creek, jacks were retained in the same

proportion as the other adults taken for broodstock. The remaining jacks were cryopreserved and later sacrificed. Following fertilization, eggs were transported back to McCall Fish Hatchery at the end of each spawn day.

Multiple males were spawned with each female to ensure fertilization and to increase genetic diversity (Table 3). Only two adult males were not utilized for spawning purposes. One of the males died before being spawned and the other male was trapped after the last female had been spawned. A cryopreservation sample was later collected from this male. Cryopreservation samples were taken from 29 Johnson Creek male salmon (15 adult males and 14 jacks) throughout the course of the spawning season. Samples were collected by personnel from the NPT's Cryopreservation project (BPA Project Number: 199703800).

Fecundity

The 16 Johnson Creek females artificially spawned produced 65,060 green eggs. Eggs from each individual female were incubated in separate incubator trays at McCall Fish Hatchery. Fecundity for the Johnson Creek females ranged from a low of 2,520 to a high of 5,162, with an average fecundity of 4,066. The fecundity of the two five year old females spawned, tag # 0078 and # 0126, was 4,575 and 4,879 respectively (Table 3).

Table 3: Spawning and Egg Incubation Results

Date Spawned	Spawn Lot	Female ID	Males ID	Jack ID	Fecundity	% Egg Eye-Up	# Eyed Eggs
8/15/00	1	0057	0035 0002 0001 0014		3,780	95.0	3,589
8/22/00	3	0064	0099 0097		2,982	89.8	2,677
8/22/00	3	0074	0072 0102		3,782	89.5	3,383
8/22/00	3	0078	0058 0086		4,575	69.3	3,169
8/25/00	4	0011	0035 0066		3,745	79.8	2,988
8/25/00	4	0008	0018 0086		3,060	80.0	2,448
8/25/00	4	0121	0112 0035		5,162	95.3	4,920
8/25/00	4	0087	0120 0079		2,520	72.7	1,831
8/29/00	5	0020	0099	0048	4,234	94.5	4,002
8/29/00	5	0023	0060 0129		3,999	87.2	3,485
8/29/00	5	0051	0014 0001		4,953	91.1	4,511
8/29/00	5	0109	0058	0050	4,489	89.2	4,006
8/29/00	5	0126	0113 0127		4,879	80.0	3,902
8/29/00	5	0131	0102 0120 0112		4,309	97.2	4,190
9/1/00	6	0144	0140 0052		4,318	68.8	2,970
9/1/00	6	0134	0139 0070		4,273	91.3	3,900
Total		16	33	2	65,060		55,971
Average					4,066	86	

Disease Testing

IDFG pathologists collected disease samples from all 16 females, and 27 of the males that were used for artificial spawning for Johnson Creek project. Overall, 43 fish were sampled for BKD, 18 tested negative, 22 tested low, and 3 tested high (greater than 0.25 ELISA OD value). All 16 females tested positive for BKD at low ELISA levels. In spite of these results, no eggs were culled in 2000. Ovarian fluid was also collected from all females, and all tested negative for IHN and IPN. In addition, 51 samples were collected for whirling disease testing. Results from these tests were also negative.

Prior to 1998, no disease history had been established for the Johnson Creek wild/naturally spawning population. Positive BKD samples found in Johnson Creek adult chinook in 1998 and 2000 indicate that BKD was present prior to artificial propagation practices (Daniel and Gebhards 2003).

Nutrient Enhancement

All 73 carcasses of adult and jack salmon used for artificial spawning were returned to Johnson Creek and put in the river to contribute to nutrient enhancement. Natural spawning areas were chosen as disposition points to stimulate natural nutrient enhancement to the stream. Mundy (1998) discusses that smolt productivity may be related to the amount of marine nutrients found in a river system, while Mathisen et al. (1988), the NRC (1996), and Larkin and Slaney (1997) reported the carbon and marine nutrients brought by adult chinook salmon back to their spawning streams and lakes are important for productivity. Schmidt et al. (1998) found that sockeye salmon *Oncorhynchus nerka* carcasses account for an estimated 90% of the phosphorus above the baseline loading. Increased adult abundance and subsequent carcass decomposition will benefit juvenile fish rearing in Johnson Creek and the surrounding ecosystem.

Incubation and Green Egg to Eyed Egg Survival

Eggs taken from Johnson Creek broodstock were incubated at the McCall Fish Hatchery. Eggs eyed-up at approximately 550 DTU and were then shocked, picked, and enumerated. Hatching began at approximately 925 DTU.

Green egg to eyed egg survival ranged from 69.3% to 97.2% with an overall average of 86% resulting in 55,971 eyed eggs (Table 3). No eggs were culled due to high BKD levels. Low eye-up in brood year 1998 was attributed in part to the fact that many of the females were spawned within a very short time after being transported to the holding and spawning facility. However, there did not appear to be a correlation between when a female was transported and poor survival to eye-up in brood year 2000. An increase in fish numbers is indicated in Appendix B between June and July and is attributed to hand counts taken during CWT tagging.

Rearing

Johnson Creek fish are initially reared indoors at the McCall Fish Hatchery. McCall Fish Hatchery personnel applied rearing practices used for SFSR juveniles to the Johnson Creek fish.

Fry were set out in the rearing vats between 1725 and 1750 DTU. Initial feeding began between 1,750 and 1,775 DTU approximately three days after set out. The vats started at half length and were extended to full length when the density index (DI) reached 0.42 around mid April 2001 (Appendix B).

Beginning growth rates were slow, only 0.003- 0.004-inches per day, due to cold water temperatures of only 37°F to 39°F. The fry were started on BioDiet #2 and #3 feed and remained on #3 until they reached 700 fish per pound. BioDiet feed has been used successfully at McCall Fish Hatchery, using modified feed rates. Two medicated feed treatments of Aquamycin 100 were administered in May 2001 and December of 2001 as a preventative measure for BKD. The overall feed conversion for Johnson Creek fish was 1.42:1 (Appendix B).

Johnson Creek fish were reared inside in the rearing vats in order to keep them separate from SFSR fish and until outdoor rearing space was made available in the collection basin. Density indices for the McCall Fish Hatchery are usually kept below 0.35. Density indices for Johnson Creek fish reached 0.54 in September 2001 due to a lack of rearing space. Johnson Creek fish were allowed more rearing space as soon as it became available. Johnson Creek fish were moved to the collection basin in December 2001. They were reared there until their release in March 2002.

Tagging

Brood year 2000 Johnson Creek fish were marked with coded wire tags (CWT – tag code 108971) in the nose and yellow visual implant elastomer (VIE) tags in the adipose tissue behind the left eye. Coded wire tagging was performed in July 2001 and VIE tagging in November 2001 by the IDFG marking crew. At the time of CWT tagging, a hand count of the fish is conducted. This hand count resulted in an increase in the number of fish being reared from 53,813 to 57,876.

In addition, 9,987 (approximately 17%) of these fish were also PIT tagged with 14 mortalities from handling and marking. PIT tagging was conducted by NPT personnel over a two-day period, on January 28 and 29, 2002.

During PIT tagging, NPT personnel also conducted a tag retention study of CWT and VIE tags. A total of 2,269 fish were interrogated for CWT tags and 2,512 fish were interrogated for VIE tags. The retention rate for CWT tags was 96.52 percent and 98.09 percent for VIE tags. Overall, 99.87 percent of the fish were tagged with either CWT, VIE or both. A more detailed summary of the tag retention study is found in Vogel and Nelson (2003 draft).

Smolt Release

Smolts were released into Johnson Creek on March 18, 19, 20, 2002. A total of 57,392 fish were trucked and released by JCAPE and other NPT personnel. At the time of release Johnson Creek smolts averaged 29.3 fpp with an average fork length of 110.0 mm.

Smolts were released into a temporary acclimation channel in Johnson Creek. An artificial barrier at the lower end of a side channel of Johnson Creek was installed to retarded downstream migration for a period of 5 days in order to keep the fish contained and provide an acclimation period. The upper end was left open so fish could leave volitionally. Fish were monitored over the 5 day period and observed exiting through the riffle at the upper end and through small gaps in the barrier fence at the lower end of the channel. Although ice build-up and predators were a concern at the site, NPT personnel monitoring the site reported no major difficulties. The barrier fence was removed March 22 and fish were allowed to emigrate freely into Johnson Creek.

CONCLUSIONS

The second season of implementation of the JCAPE supplementation program was a success. Improvements in field operations in 2000 were made because of experience gained from 1998. These included:

- Modifications to the weir and trap box made trapping more efficient and fish friendly. The modified trap box entrance prevented fish from swimming in and out of the trap box. Shade and jump barriers were also provided.
- The use of the temporary circular tank for holding Johnson Creek males streamlined the spawning process and reduced stress to both SFSR and Johnson Creek males.
- Eggs from each individual female were incubated separately, providing better control of these eggs.
- The brood year 1998 smolts were directly released into Johnson Creek and immediately emigrated from Johnson Creek. This brought about the development of a temporary acclimation channel. Brood year 2000 smolts placed in this acclimation channel were observed moving out of Johnson Creek immediately, but many of the fish remained in the acclimation area several days after the temporary barrier was removed.

There are many more improvements that can be made to add to the continued success of the JCAPE project. Primarily, additional facilities are needed to conduct field operations properly.

- A permanent trapping facility on Johnson Creek would eliminate many of the problems encountered with the current trapping equipment. Trapping would be more efficient, more fish friendly and easier for NPT personnel to install and operate.
- A permanent holding facility would eliminate the need to transport fish. This would greatly reduce the amount of stress placed on fish retained for broodstock. Permanent holding would also eliminate problems encountered with temporary holding facilities at the SFSR facility. Johnson Creek females would not have to cohabitate with SFSR fish in a foreign water source. Males would not have the added stress of moving from fish truck to raceway to circular holding tank. Many of the dangers of using pumps, pipes, and hoses for providing a temporary water source would be eliminated.
- A permanent acclimation facility would be much more efficient and effective than the temporary facility currently being used.

Support and the approval of the agencies governing the activities of JCAPE will continue to be solicited by JCAPE and NPT-DFRM administration. Adaptive management practices will continue to be used to improve the JCAPE project and increase its effectiveness restoring chinook salmon populations in Johnson Creek.

LITERATURE CITED

- BPA (Bonneville Power Administration). 1998. Environmental Clearance Memorandum: Categorical Exclusion for Adult Collection Activities to be conducted on Johnson Creek. May 8, 1998. U.S. Department of Energy, Bonneville Power Administration, Portland, Oregon.
- CRITFC (Columbia River Intertribal Fish Commission). 1996. *Wy-Kan-Ush-Me Wa-Kush-Wit, Spirit of the Salmon*. The Columbia River Anadromous Fish Restoration Plan of the Nez Perce, Umatilla, Warm Springs, and Yakima Tribes. Two Volumes. Columbia River Intertribal Fish Commission. Portland, OR.
- Daniel, M., and J. S. Gebhards. 2003. Brood Year 1998: Johnson Creek Chinook Salmon Supplementation Report. Period Covered: July 1998 through April 2000. Johnson Creek Artificial Propagation Enhancement Project. Operations and Maintenance Program. Nez Perce Tribe Department of Fisheries Resources Management. Prepared for U.S. Department of Energy, Bonneville Power Administration. BPA Project Number 199604300. BPA Contract Number 4393.
- Larkin, G.A., and P.A. Slaney. 1997. Implications of Trends in Marine-Derived Nutrient Influx to South Coastal British Columbia Salmonid Production. *Fisheries*. 22 (11):16-24.
- Lothrop, R.C. 2000. Johnson Creek Artificial Propagation Enhancement Project Section 10 Permit Application. Renewal of permit 1147. Submitted to NMFS as part of ESA Section 10 consultation. CRITFC. Portland, OR.
- Mathisen, O.A., and five others. 1988. Recycling of marine elements transported into freshwater systems by anadromous salmon. *Internationale Vereinigung Für Theoretische und Angewandte Limnologie Verhandlungen*. 23:2,249-2,258.
- Mundy, P.R. 1998. Imnaha River Spring Chinook Fisheries Management Plan Development. Final Report. Prepared for the Nez Perce Tribe. September, 1998.
- Nelson, D.D., and J.L. Vogel. 2003. Draft. Johnson Creek Artificial Propagation Enhancement Monitoring and Evaluation Project. Annual Progress Report. Period Covered: January 1, 1998 through December 31, 1998. Nez Perce Tribe Department of Fisheries Resources Management. Prepared for U.S. Department of Energy, Bonneville Power Administration.
- NMFS (National Marine Fisheries Service). 1995. Summary: Proposed recovery plan for Snake River salmon. U. S. Department of Commerce, National Oceanic and Atmospheric Administration, Portland, Oregon.
- NOAA. 2004 draft. Environmental Assessment of a National Marine Fisheries Service (NMFS) Action to Issue Research/Enhancement Permit 1250 to the Columbia River Inter-tribal Fish Commission (CRITFC) under Section 10(a) (A) of the Endangered Species Act.

- NPT (Nez Perce Tribe) and PRRG (Production and Restoration Research Group). 2001. Johnson Creek Artificial Propagation Enhancement Hatchery and Genetic Management Plan. Nez Perce Tribe Department of Fisheries Resources Management.
- NRC (National Resource Council). 1996. Upstream: Salmon in Society in the Pacific Northwest. Committee on Protection and Management of Pacific Northwest Anadromous Salmonids, Board on Environmental Studies and Toxicology, Commission on Life Sciences. National Academy of Sciences. 452p.
- Piper, R. G., I.B. McElwain, L. E. Orme, J.P. McCraren, L.G. Fowler, J.R. Leonard, 1982. Fish Hatchery Management. Department of the Interior, U.S. Fish and Wildlife Service.
- USFS (United States Forest Service). 1990. Boise National Forest Land and Resource Management Plan (Forest Plan). Boise, Idaho.
- USFS (United States Forest Service). 1995. Upper South Fork Salmon River and Johnson Creek Watershed Analysis. Cascade Ranger District, Boise National Forest and Krassel Ranger District, Payette National Forest.
- Vogel, J.L. and D.D. Nelson. 2003. Draft. Brood Year 2000 Juvenile and Brood Year 2001 Adult Chinook Salmon Report. Johnson Creek Artificial Propagation Enhancement Monitoring and Evaluation Project. Nez Perce Tribe Department of Fisheries Resources Management. Prepared for U.S. Department of Energy, Bonneville Power Administration.

APPENDICES

Appendix A: Adult Trapping Summary

OPERCLE TAG #	DATE TRAPPED	SEX	AGE	FKL (cm)	MEH (mm)	MARKS	DNA SAMPLE TAKEN	ERYTHRO. INJECTION	OPERCLE TAGGED	FLOY TAGGED	DISPOSITION (SFSR=RETAINED FOR BROODSTOCK)
0001	6/29/00	M	4	72	59	NONE	YES	YES	YES	YES	SFSR
0002	6/29/00	M	4	78	63	NONE	YES	YES	YES	YES	SFSR
0003	6/29/00	F	4	74	61	NO	YES	NO	YES	YES	RELEASED
0004	6/29/00	J	3	45	38	NONE	YES	NO	YES	YES	RELEASED
0005	6/30/00	J	3	63	52	NONE	YES	NO	YES	YES	RELEASED
0006	7/1/00	F	4	73	61	AD CLIP	YES	NO	YES	YES	RELEASED
0007	7/1/00	J	3	43	33	NONE	YES	NO	YES	YES	RELEASED
0008	7/2/00	F	4	80	69	NONE	YES	YES	YES	YES	SFSR
0009	7/5/00	J	3	48	39	NONE	YES	NO	YES	YES	SFSR
0010	7/6/00	M	4	79	64	NONE	YES	NO	YES	NO	RELEASED
0011	7/6/00	F	4	75	64	NONE	YES	YES	YES	YES	SFSR
0012	7/6/00	J	3	56	46	AD CLIP	YES	NO	YES	NO	RELEASED
0013	7/6/00	J	3	47	38	NONE	YES	NO	YES	YES	SFSR
0014	7/9/00	M	4	78	64	NONE	YES	YES	YES	YES	SFSR
0015	7/9/00	J	3	52	43	NONE	YES	NO	YES	YES	SFSR
0016	7/9/00	J	3	45	37	NONE	YES	NO	YES	NO	RELEASED
0017	7/9/00	J	3	47	39	AD CLIP	YES	NO	YES	NO	RELEASED
0018	7/10/00	M	4	74	61	NONE	YES	YES	YES	YES	SFSR
0019	7/10/00	J	3	58	49	NONE	YES	NO	YES	YES	SFSR
0020	7/10/00	F	4	80	66	NONE	YES	YES	YES	YES	SFSR
0021	7/11/00	F	4	76	63	NONE	YES	NO	YES	YES	RELEASED
0022	7/11/00	M	4	76	62	NONE	YES	NO	YES	NO	RELEASED
0023	7/11/00	F	4	80	65	NONE	YES	YES	YES	YES	SFSR
0024	7/11/00	J	3	55	44	PIT	YES	NO	YES	YES	SFSR
0025	7/11/00	J	3	45	37	NONE	YES	NO	YES	NO	RELEASED
0026	7/11/00	M	4	79	62	NONE	YES	NO	YES	NO	RELEASED
0027	7/11/00	J	3	53	44	NONE	YES	NO	YES	YES	SFSR
0028	7/12/00	J	3	57	46	NONE	YES	NO	YES	NO	RELEASED
0029	7/12/00	F	4	80	68	NONE	YES	NO	YES	NO	RELEASED
0030	7/12/00	J	3	65	52	NONE	YES	NO	YES	YES	SFSR
0031	7/12/00	J	3	53	45	NONE	YES	NO	YES	YES	SFSR
0032	7/13/00	J	3	58	49	NONE	YES	NO	YES	YES	RELEASED

OPERCLE TAG #	DATE TRAPPED	SEX	AGE	FKL (cm)	MEH (mm)	MARKS	DNA SAMPLE TAKEN	ERYTHRO. INJECTION	OPERCLE TAGGED	FLOY TAGGED	DISPOSITION (SFSR=RETAINED FOR BROODSTOCK)
0033	7/14/00	J	3	55	45	NONE	YES	NO	YES	NO	RELEASED
0034	7/15/00	M	4	74	62	NONE	YES	NO	YES	NO	RELEASED
0035	7/15/00	M	4	77	62	NONE	YES	YES	YES	YES	SFSR
0036	7/15/00	J	3	50	41	AD CLIP	YES	NO	YES	YES	RELEASED
0037	7/15/00	J	3	55	45	NONE	YES	NO	YES	NO	SFSR
0038	7/15/00	J	3	52	42	NONE	YES	NO	YES	NO	SFSR
0039	7/16/00	J	3	56	48	NONE	YES	NO	YES	NO	RELEASED
0040	7/17/00	J	3	47	39	NONE	YES	NO	YES	NO	RELEASED
0041	7/18/00	J	3	55	46	NONE	YES	NO	YES	NO	RELEASED
0042	7/19/00	M	4	80	62	NONE	YES	NO	YES	NO	RELEASED
0043	7/19/00	J	3	50	41	NONE	YES	NO	YES	NO	SFSR
0044	7/20/00	J	3	53	44	NONE	YES	NO	YES	YES	SFSR
0045	7/21/00	J	3	60	48	NONE	YES	NO	YES	YES	SFSR
0046	7/21/00	J	3	58	46	NONE	YES	NO	YES	YES	SFSR
0047	7/24/00	J	3	45	38	NONE	YES	NO	YES	NO	RELEASED
0048	7/25/00	J	3	56	46	NONE	YES	NO	YES	YES	SFSR
0049	7/26/00	J	3	57	46	NONE	YES	NO	YES	NO	RELEASED
0050	7/27/00	J	3	60	51	NONE	YES	NO	YES	YES	SFSR
0051	8/4/00	F	4	75	62	NONE	YES	YES	YES	YES	SFSR
0052	8/6/00	M	4	75	62	NONE	YES	YES	YES	YES	SFSR
0053	8/9/00	J	3	55	46	NONE	YES	NO	YES	NO	RELEASED
0054	8/11/00	F	4	74	63	NONE	YES	NO	YES	NO	RELEASED
0055	8/13/00	J	3	52	42	NONE	YES	NO	YES	NO	SFSR
0056	8/13/00	M	4	74	58	NONE	YES	NO	YES	NO	RELEASED
0057	8/13/00	F	4	75	64	NONE	YES	YES	YES	YES	SFSR
0058	8/14/00	M	4	78	64	NONE	YES	YES	YES	YES	SFSR
0059	8/14/00	M	4	76	60	NONE	YES	NO	YES	NO	RELEASED
0060	8/15/00	M	4	71	55	NONE	YES	YES	YES	YES	SFSR
0061	8/16/00	F	4	81	66	NONE	YES	NO	YES	NO	RELEASED
0062	8/16/00	M	4	77	62	NONE	YES	NO	YES	NO	RELEASED
0063	8/16/00	J	3	55	42	NONE	YES	NO	YES	NO	RELEASED
0064	8/17/00	F	4	75	61	NONE	YES	YES	YES	YES	SFSR
0065	8/17/00	J	3	53	44	PIT	YES	NO	YES	YES	SFSR

OPERCLE TAG #	DATE TRAPPED	SEX	AGE	FKL (cm)	MEH (mm)	MARKS	DNA SAMPLE TAKEN	ERYTHRO. INJECTION	OPERCLE TAGGED	FLOY TAGGED	DISPOSITION (SFSR=RETAINED FOR BROODSTOCK)
0066	8/17/00	M	4	81	64	NONE	YES	YES	YES	YES	SFSR
0067	8/17/00	M	4	73	57	NONE	YES	NO	YES	NO	RELEASED
0068	8/17/00	F	4	70	59	NONE	YES	NO	YES	NO	RELEASED
0069	8/17/00	J	3	56	44	NONE	YES	NO	YES	YES	SFSR
0070	8/18/00	M	4	74	58	NONE	YES	YES	YES	YES	SFSR
0071	8/19/00	M	4	75	58	NONE	YES	NO	YES	NO	RELEASED
0072	8/20/00	M	5	83	65	NONE	YES	YES	YES	YES	SFSR
0073	8/20/00	J	3	58	46	NONE	YES	NO	YES	NO	RELEASED
0074	8/20/00	F	4	74	60	NONE	YES	YES	YES	YES	SFSR
0075	8/20/00	J	3	56	44	NONE	YES	NO	YES	YES	SFSR
0076	8/20/00	J	3	58	47	NONE	YES	NO	YES	NO	RELEASED
0077	8/20/00	J	3	61	50	NONE	YES	NO	YES	YES	SFSR
0078	8/20/00	F	5	82	66	NONE	YES	YES	YES	YES	SFSR
0079	8/20/00	M	4	78	60	NONE	YES	YES	YES	YES	SFSR
0080	8/20/00	J	3	54	44	NONE	YES	NO	YES	NO	RELEASED
0081	8/20/00	J	3	48	39	AD CLIP	YES	NO	YES	NO	RELEASED
0082	8/20/00	J	3	46	38	NONE	YES	NO	YES	YES	SFSR
0083	8/21/00	F	4	77	63	NONE	YES	NO	YES	NO	RELEASED
0084	8/21/00	M	4	76	60	NONE	YES	NO	YES	NO	RELEASED
0085	8/21/00	J	3	58	47	NONE	YES	NO	YES	YES	SFSR
0086	8/21/00	M	4	76	61	NONE	YES	YES	YES	YES	SFSR
0087	8/21/00	F	4	73	59	NONE	YES	YES	YES	YES	SFSR
0088	8/21/00	J	3	51	41	NONE	YES	NO	YES	NO	RELEASED
0089	8/21/00	J	3	55	44	NONE	YES	NO	YES	YES	SFSR
0090	8/21/00	J	3	51	41	NONE	YES	NO	YES	NO	RELEASED
0091	8/21/00	J	3	46	37	NONE	YES	NO	YES	YES	SFSR
0092	8/21/00	J	3	45	37	AD CLIP	YES	NO	YES	NO	RELEASED
0093	8/22/00	J	3	58	46	NONE	YES	NO	YES	YES	SFSR
0094	8/22/00	M	5	84	65	NONE	YES	NO	YES	NO	RELEASED
0095	8/22/00	F	4	80	65	NONE	YES	NO	YES	NO	RELEASED
0096	8/22/00	M	4	78	61	NONE	YES	NO	YES	NO	RELEASED
0097	8/22/00	M	4	78	61	NONE	YES	YES	YES	YES	SFSR
0098	8/22/00	M	4	73	58	NONE	YES	NO	YES	NO	RELEASED

OPERCLE TAG #	DATE TRAPPED	SEX	AGE	FKL (cm)	MEH (mm)	MARKS	DNA SAMPLE TAKEN	ERYTHRO. INJECTION	OPERCLE TAGGED	FLOY TAGGED	DISPOSITION (SFSR=RETAINED FOR BROODSTOCK)
0099	8/22/00	M	4	76	60	NONE	YES	YES	YES	YES	SFSR
0100	8/22/00	M	4	76	59	NONE	YES	NO	YES	NO	RELEASED
0101	8/22/00	F	4	81	66	NONE	YES	NO	YES	NO	RELEASED
0102	8/22/00	M	4	71	57	NONE	YES	YES	YES	YES	SFSR
0103	8/22/00	J	3	60	48	NONE	YES	NO	YES	NO	RELEASED
0104	8/22/00	J	3	56	45	PIT	YES	NO	YES	YES	SFSR
0105	8/22/00	M	4	81	65	AD CLIP	YES	NO	NO	NO	RELEASED
0106	8/22/00	M	4	79	62	AD CLIP	YES	NO	NO	NO	RELEASED
0107	8/22/00	J	3	53	42	NONE	YES	NO	YES	NO	RELEASED
0108	8/23/00	F	5	82	68	NONE	YES	NO	YES	NO	RELEASED
0109	8/23/00	F	4	75	62	NONE	YES	YES	YES	YES	SFSR
0110	8/23/00	J	3	55	44	NONE	YES	NO	YES	NO	RELEASED
0111	8/23/00	J	3	54	43	NONE	YES	NO	YES	YES	SFSR
0112	8/23/00	M	4	77	63	NONE	YES	YES	YES	YES	SFSR
0113	8/23/00	M	4	81	65	NONE	YES	YES	YES	YES	SFSR
0114	8/23/00	J	3	60	49	NONE	YES	NO	YES	NO	RELEASED
0115	8/23/00	M	4	72	58	NONE	YES	NO	YES	NO	RELEASED
0116	8/24/00	M	4	76	62	NONE	YES	NO	YES	NO	RELEASED
0117	8/24/00	M	5	83	65	NONE	YES	NO	YES	NO	RELEASED
0118	8/24/00	F	4	75	61	NONE	YES	NO	YES	NO	RELEASED
0119	8/24/00	M	4	80	64	AD CLIP	YES	NO	YES	NO	RELEASED
0120	8/24/00	M	4	81	64	NONE	YES	YES	YES	YES	SFSR
0121	8/24/00	F	4	77	62	NONE	YES	YES	YES	YES	SFSR
0122	8/24/00	M	5	84	67	NONE	YES	YES	YES	YES	SFSR
0123	8/25/00	F	4	74	62	NONE	YES	NO	YES	NO	RELEASED
0124	8/25/00	M	5	84	67	NONE	YES	NO	YES	NO	RELEASED
0125	8/25/00	M	5	84	67	AD CLIP	YES	NO	YES	NO	RELEASED
0126	8/26/00	F	5	93	78	NONE	YES	YES	YES	YES	SFSR
0127	8/26/00	M	4	78	61	NONE	YES	YES	YES	YES	SFSR
0128	8/27/00	M	4	73	58	NONE	YES	NO	YES	NO	RELEASED
0129	8/27/00	M	4	80	63	NONE	YES	YES	YES	YES	SFSR
0130	8/27/00	M	4	81	64	NONE	YES	NO	YES	NO	RELEASED
0131	8/27/00	F	4	69	57	NONE	YES	YES	YES	YES	SFSR

OPERCLE TAG #	DATE TRAPPED	SEX	AGE	FKL (cm)	MEH (mm)	MARKS	DNA SAMPLE TAKEN	ERYTHRO. INJECTION	OPERCLE TAGGED	FLOY TAGGED	DISPOSITION (SFSR=RETAINED FOR BROODSTOCK)
0132	8/27/00	J	3	55	43	NONE	YES	YES	YES	YES	SFSR
0133	8/27/00	F	4	81	66	NONE	YES	NO	YES	NO	RELEASED
0134	8/27/00	F	4	80	65	NONE	YES	YES	YES	YES	SFSR
0135	8/27/00	J	3	47	38	NONE	YES	NO	YES	NO	RELEASED
0136	8/27/00	J	3	48	39	NONE	YES	YES	YES	YES	SFSR
0137	8/27/00	J	3	52	41	AD CLIP	YES	NO	YES	NO	RELEASED
0138	8/30/00	M	4	73	59	AD CLIP	NO	NO	YES	NO	RELEASED
*****	8/30/00	F	4	77	64	NONE	YES	NO	NO	NO	WEIR MORT
0139	8/31/00	M	4	81	65	NONE	YES	YES	YES	YES	SFSR
0140	8/31/00	M	4	71	57	NONE	YES	YES	YES	NO	SFSR
0141	8/31/00	J	3	66	54	NONE	YES	NO	YES	NO	RELEASED
0142	9/1/00	F	4	73	63	AD CLIP	YES	NO	YES	NO	RELEASED
0143	9/1/00	M	4	79	63	NONE	YES	NO	YES	NO	RELEASED
0144	9/1/00	F	4	78	65	NONE	YES	YES	YES	YES	SFSR
0145	9/1/00	M	4	80	66	AD CLIP	YES	NO	YES	NO	RELEASED
0146	9/1/00	J	3	57	47	NONE	YES	NO	YES	YES	SFSR
0147	9/1/00	J	3	53	44	NONE	YES	NO	YES	NO	RELEASED
0148	9/2/00	J	3	59	49	NONE	YES	NO	YES	YES	SFSR
0149	9/2/00	J	3	52	42	NONE	YES	NO	YES	NO	RELEASED
0150	9/3/00	M	5	83	67	NONE	YES	YES	YES	YES	SFSR
0151	9/6/00	M	5	84	67	NONE	YES	NO	YES	NO	RELEASED