Document ID #P111671
Report covers work performed under BPA contract #4826
Report was completed under BPA contract #30064

Duck Valley Reservoirs Fish Stocking and O&M

Annual Progress Report
Performance Period: October 1, 2006 – September 30, 2007

Prepared by:

Jake Sellman, Project Manager

AND

Tim Dykstra, Director

Department of Fish, Wildlife and Parks
Shoshone-Paiute Tribes
P.O. Box 219
Owyhee, NV 89832

Funded by:

U.S. Department of Energy
Bonneville Power Administration
Division of Fish and Wildlife
Portland, OR 97208-3621

BPA Project Number: 1995-015-06
Contract Number: 4826

Abstract/Executive Summary

*The Duck Valley Reservoirs Fish Stocking and Operations and Maintenance (DV Fisheries) project is an ongoing resident fish program that serves to partially mitigate the loss of anadromous fish that resulted from downstream construction of the hydropower system. The project's goals are to enhance subsistence fishing and educational opportunities for Tribal members of the Shoshone-Paiute Tribes and provide resident fishing opportunities for non-Tribal members. In addition to stocking rainbow trout (*Oncorhynchus mykiss*) in Mountain View, Lake Billy Shaw, and Sheep Creek Reservoirs, the program is also designed to maintain healthy aquatic conditions for fish growth and survival, to provide superior facilities with wilderness qualities to attract non-Tribal angler use, and to offer clear, consistent communication with the Tribal community about this project as well as outreach and education within the region and the local community.*

Tasks for this performance period are divided into operations and maintenance plus monitoring and evaluation. Operation and maintenance of the three reservoirs include fences, roads, dams and all reservoir structures, feeder canals, water troughs and stock ponds, educational signs, vehicles and equipment, and outhouses. Monitoring and evaluation activities included creel, gillnet, wildlife, and bird surveys, water quality and reservoir structures monitoring, native vegetation planting, photo point documentation, control of encroaching exotic vegetation, and community outreach and education.

The three reservoirs are monitored in terms of water quality and fishery success. Sheep Creek Reservoir was very unproductive this year as a fishery. Fish morphometric and water quality data indicate that the turbidity is severely impacting trout survival. Lake Billy Shaw was very productive as a fishery and received good ratings from anglers. Mountain View was also productive and anglers reported a high number of quality sized fish. Water quality (specifically dissolved oxygen and temperature) is the main limiting factor in our fisheries.

Introduction

The Snake River Basin is estimated to have contained 79% of the stream miles suitable for chinook salmon in the Columbia River Basin (Idaho Department of Fish and Game 1985). Using the stream mile estimate, IDFG estimated that 650,000 - 1,030,000 adult chinook, 117,000 - 229,800 steelhead, and 14,400 - 57,400 sockeye were produced annually in the Snake River and its tributaries above what is now Hells Canyon Dam.

With the completion of the Hells Canyon Complex, anadromous salmonids were extirpated from the upper portion of the Snake River and its tributaries. In the Northwest Power and Conservation Council's (NPCC) 1994 Columbia River Basin Fish and Wildlife Program (Program), the NPCC acknowledged that "Salmon and steelhead probably will never be able to return to some areas of the basin because of blockages by dams. These areas include the areas above Chief Joseph and Grand Coulee Dams, the Hells Canyon Complex and other smaller blocked areas." Subsequently, the NPCC suggested that: 1) mitigation in blocked areas is appropriate where salmon and steelhead were affected by the development and operation of the hydroelectric projects and 2) to treat the Columbia River and its tributaries as a system, resident fish substitutions are reasonable for lost salmon and steelhead in areas where in-kind mitigation

cannot currently occur. This suggestion encompassed the Duck Valley Indian Reservation (DVIR).

The DVIR encompasses approximately 289,820 tribally-owned acres surrounded by Idaho to the north and Nevada to the south. Currently, there are approximately 1,800 enrolled Tribal members. The Reservation is in the Middle Snake Province and is within both the Bruneau and Owyhee subbasins. As indicated in Figure 1, the Reservation is both remote and isolated; the closest town centers are Elko, Nevada and Mountain Home, Idaho, both approximately 100 miles from the Reservation's small town of Owyhee. These centers are the closest places to obtain supplies.

Figure 1: Location of the DVIR in relation to the Pacific Northwest

There are three reservoirs on the DVIR that the project oversees: Sheep Creek, Mountain View, and Lake Billy Shaw (Figure 2). All three water bodies are formed by earthen dams and are dependent on the irrigation system that supports the local agriculture. Vegetation in the drainage basins of all three reservoirs mainly consists of a xeric, shrub-steppe vegetative community with intermittent wetland vegetation. Typical vegetation includes Great Basin shrub cover types including various forbs, sagebrush, rabbitbrush, and grasses such as Idaho fescue. Other herbaceous layers have been eliminated and plant diversity has decreased due to grazing (1997 EA and FONSI). The surrounding habitat supports big game (mainly antelope), carnivores, waterfowl, migratory bird species, raptors, reptiles, and amphibians. The three reservoirs are important elements to the DVIR in terms of providing and protecting large tracts of habitat and providing culturally-important fish resources to the local Tribal members.

Sheep Creek Reservoir (SCR) was built in the mid-50's and is the most southern of the three. It is the largest and has a surface area of approximately 855 acres and a maximum depth of 23 feet. This reservoir is fed from the same canal as Lake Billy Shaw and is well known throughout the western states for its highly productive trout fishery. Fly anglers commonly use a fly pattern known as the "Sheep Creek Special," dubbed for its high success at the DVIR reservoirs. Mountain View Reservoir (MVR) was constructed in 1969 and is the only reservoir located on the northern half of the Reservation, which is surrounded by Idaho. It is approximately 640 acres and has a maximum depth of 24 feet. This reservoir is well known for its crystal clear water and reliable fishing experience. Both reservoirs were historically stocked through USFWS funding (Burge and Miller, 1990).

Figure 2: Locations of the three reservoirs on the DVIR

The final and most recently built (1998) reservoir is Lake Billy Shaw (LBS) and was built within the Billy Shaw Slough in the south-central portion of the Duck Valley Indian Reservation, 11 km/7 mi west of the town of Owyhee. Construction was completed in 1998 and the first full fishing season took place in 2003. This reservoir is located slightly south of the Reservation's centerline and is north of SCR. The slough was an intermittent stream which flowed northward in a wide, gently sloped drainage channel in moderately dissected alluvial terrain that had not been developed for agriculture or settlement (1997 EA and FONSI).

Developed and managed for recreational and Tribal subsistence fishing, Lake Billy Shaw is a reservoir with 17.5 miles of shoreline and has a maximum pool elevation of 5,351 feet. The reservoir has a 3,300 acre foot volume with 430 acres of surface area, a maximum depth of 36 feet, and a mean depth of 23 feet; 76% of the reservoir is at 15 feet or greater depth. Inflows to the reservoir include water diverted from the East Fork Owyhee River into the Highline canal, which is then diverted into either Sheep Creek Reservoir or Lake Billy Shaw feed canal (beginning in R 51 E, T 46 N, Section 2). Recorded annual fluctuations can be highly variable (Figure 6, page 14).

The feed canal can carry flows up to 30 cubic feet per second (cfs) without causing erosion damage. Inflows to the reservoir fluctuate drastically according to: time of year, releases from the upstream Wildhorse Reservoir, and upstream irrigation needs. Outflows generally include only surface spillage from the reservoir, although summer evaporation and bank seepage often exceed inflows. Escapement of fish from the reservoir is restricted by a gated outlet and drum screens with paddle wheels at the inlet.

Lake Billy Shaw Reservoir was completed in 1998 to address measures 10.8C.4 and 10.8C.7 of the NPCC's 1994 Program and is presently operated per the "Substitution for Anadromous Fish Losses" and "Resident Fish Losses" provisions of the NPCC's 2000 Program. The NPCC's June 5, 1998, letter to Chairman Paiva and Mr. Robert Lohn states that the goal of Project 1995-015-06 is "to develop, on the Duck Valley Reservation, an enclosed reservoir rainbow trout 'put and take' fishery at Lake Billy Shaw."

During the Three-Step Review for Lake Billy Shaw and initial deliberations with the NPCC and BPA, however, artificial production of native redband trout and associated stocking costs for Lake Billy Shaw were included in Project 1995-006-00 titled "*Shoshone-Paiute /Shoshone-Bannock Joint Culture Facility*" as indicated in the NPCC's May 12, 1998, Decision Memorandum. Nevertheless, the Independent Scientific Review Panel (ISRP) (ISRP 2001-3) recommended to the NPCC that Project 1995-006-00 should not move beyond Step 3 and into construction and suggested that the project sponsors did not adequately address the conditions provided in the NPCC's Step 2 decision. Despite the ODFW, Montana Department of Fish, Wildlife, and Parks, and IDFG's operation of artificial production programs for redband trout (Klamath Hatchery) and Yellowstone cutthroat trout *O. clarki lewisi* (Bluewater Springs, Yellowstone River, and Big Springs trout hatcheries), respectively, the ISRP (ISRP 2001-3) suggested, without references, that hatchery rearing of redband trout and Yellowstone cutthroat trout would be difficult at best, since a high proportion of these projects have not succeeded. Subsequently, the ISRP (ISRP 2001-3) suggested that non-native rainbow trout should be used to meet the production needs of the Duck Valley Indian Reservation and that those needs could be addressed by contracting with private or state aquaculture facilities in southern Idaho.

In recommending funding for Lake Billy Shaw, the NPCC included a stipulation that the Shoshone-Paiute Tribes, BPA, IDFG, Nevada Department of Wildlife, and the ODFW identify an approach that would alleviate problems (e.g., introgression with native redband trout *O. m. gairdneri*) that could arise if hatchery-reared coastal rainbow trout escaped from Lake Billy Shaw. Subsequently, negotiations among the management entities resulted in a decision to stock sterile, hatchery-reared coastal rainbow trout; however, in light of the success that the ODFW has experienced in culturing redband trout, the Shoshone-Paiute Tribes, NPCC, and BPA initiated Project 2000-079-00 titled "*Assess Resident Fish Stocks of the Owyhee/Bruneau Basin, D.V.I.R.*" to identify the genetic structure and purity of redband trout populations throughout the East Fork Owyhee River drainage. Results from this study have enabled the Shoshone-Paiute Tribes to identify populations that could serve as broodstock sources for the development of a redband trout hatchery program in Lake Billy Shaw reservoir, as well as prioritize habitat for protection and enhancement.

Until then, Lake Billy Shaw will be stocked with sterile rainbow trout using funds from this contract with BPA. Guidelines for stocking rates and size of stocked fish are dependent upon reservoir management and vary among waterbodies; however, objectives of most reservoir management programs, regardless of the fish species and location, are to provide a total catch biomass that exceeds the stocked biomass. In April 2002, the Shoshone-Paiute Tribes initiated the stocking of Lake Billy Shaw with BPA funds from the Resident Fish Stocking project (1988-156-00) with the release of 8,000 lbs. of sterile catchable sized (10-14 in.) hatchery-reared rainbow trout. Because Lake Billy Shaw is a new reservoir, it is essential in this project to continue with a long-term study of stocking density, growth, survivorship, and yield to develop guidelines that are not approximations.

The Lake Billy Shaw Project began in 1988 with a feasibility study (Kleinfelder 1988); an Environmental Assessment was completed with a FONSI and the project proceeded to the design phase, finished in 1997 by CH2MHILL. The dam, reservoir, and associated structures were constructed and completed in 1998 by Western Construction; the project was completed ahead of schedule and with no additional costs for completion. The reservoir was initially filled in the spring of 1999, and the borrow areas were reseeded in the fall of that same year. Water quality parameters have been monitored since 1999, and activities to enhance the shorelines have been conducted since 2000 (including native plant transplanting, exotic plant removal, invertebrate sampling and transplanting, exclosure fencing, etc.)

The Lake Billy Shaw Operations and Maintenance (O&M) project went through the peer review with CBFWA (1998) and was recommended for funding in 1999. The project again went through a peer review by CBFWA and the ISRP in 1999 and 2002 and was recommended for funding for the 2000-2002 and 2003-2005 funding cycles.

Lake Billy Shaw reservoir opened October 1, 2002 as a trophy fishery. Previously, funds from BPA project 1988-156-01, Resident Fish Stocking, furnished sterile rainbow trout for Lake Billy Shaw as well as viable rainbow trout for Mountain View and Sheep Creek Reservoirs; however, in August 2003, the Resident Fish Stocking and Lake Billy Shaw O&M (BPA project 1995-015-00) projects were combined. Mid-performance period, the O&M contract title changed to "Duck Valley Reservoirs Fish Stocking and O&M" (still BPA Project 1995-015-00); the contract and performance language were changed to include O&M and fish stocking for all three reservoirs, and a line item to purchase fish for all three reservoirs was added to the budget. This report reflects the above line item, title, and task changes (the Resident Fish Stocking contract was terminated at the time of the merge). The Duck Valley Reservoirs Fish Stocking and O&M project is referred to as "DV Fisheries."

Summary of Work by Work Element

Manage and Administer Project:

Metric and location data entered into Pisces for FY07:

Metric and location data were entered into Pisces for applicable Work Elements in the FY07 Statement of Work. All status reports were submitted and approved. Missing or omitted metric and location data were updated in the final status report (September).

General administrative work performed:

The director, project manager, and secretary submitted purchase orders, payment requests, and invoices. They also addressed employment issues and coordinated with the various Tribal administrative departments.

Keep track of budget:

Detailed account reports were sent by the Finance Department monthly. These reports included expenditures, transfers, and other changes. Microsoft Excel was used to compare and contrast these changes to the approved budget. This spreadsheet was used track money expenditures and monitor remaining money.

Manage sub-contracts with fish suppliers:

The project manager and secretary maintained communication with the two fish suppliers periodically throughout the year. Contracts were revised to match the current fiscal year, reviewed by the Tribal Chairman, and sent to the suppliers. All contracts were approved and accepted by all parties.

Timely and complete submission of invoices:

All invoices were submitted to the Finance Department as soon as they were received. The secretary also retained copies of those invoices. The project manager also created monthly expenditure reports and submitted those to be sent in conjunction with the Tribal invoice to BPA.

Lease, maintain, and repair equipment:

A back-hoe and a dump truck were leased for four months. These and other equipment were maintained and repaired as needed.

Accrual estimate (year-end close):

The project manager developed an accrual estimate and forwarded to the Finance Department for review and submission to BPA.

Accrual estimate (simulated year-end close):

The project manager and a finance department representative developed an accrual estimate and submitted it to BPA.

FY08 SOW/budget package submitted to BPA:

The project manager submitted a draft FY08 SOW and budget pack in July.

Produce Environmental Compliance Documentation:

Ensure that all environmental compliance requirements have been fulfilled:

All environmental compliance requirements were met. This included NEPA and ESA compliance, cultural resource compliance, and obtaining necessary permits to legally perform work.

Outreach and Education:

Volunteer one day environmental education to students:

The project manager spent one afternoon reading fish and wildlife stories to children at the local combined school. These stories were about various species that may be encountered throughout North America. The project manager also put on a fly fishing seminar for children and for adults. In addition to the fishing clinic, he also presented information on the importance of catch-and-release values.

Volunteer one day with Boy Scout Troop activities:

The local Boy Scout Troop was inactive this year due to the Troop leader moving from the DVIR.

Work with local media:

The project manager and director talked with reporters from the Idaho Statesman. They also submitted a report to the Sho-Pai News about the current projects at the Fish, Wildlife, and Parks Department.

Provide information to anglers:

A departmental website was updated on the Shoshone-Paiute Tribes homepage (<http://www.shopaitribes.org/entrymenu.htm>). The fishing pamphlets were updated to show the 2007 changes and approximately 1,500 of these pamphlets were printed and distributed to all of the Tribes' vendors. The fishing permits were also changed to incorporate three types of permits into one type to simplify the permit sale process. The Tribes' added two more fishing permit vendors in Boise, Idaho (Cabela's and The River Keeper).

Participate in community gatherings:

The department put on the annual Arbor Day celebration where they raffled prizes, barbequed, stocked fish into the lakes, and planted trees. The department also helped with the 4th of July Pow-Wow and rodeo as well as the Tribal "Sundance" Ceremony. All D.V. Fisheries employees participated in these gatherings.

Purchase advertisement with flyfishing magazine:

The project purchased advertisements with "Northwest Flyfishing" and "Southwest Flyfishing" magazines. Both magazines are bi-monthly.

Host youth workers:

The department hosted four summer youth workers. All of these youths rotated between projects so they could obtain a well-rounded experience from the department.

Plant Vegetation:

Collect willows, bulrushes, and tules:

Willows were collected by the fisheries technician and a laborer. They planted these willows in the Lake Billy Shaw inlet stream. They planted approximately 0.10 miles of stream. Plants were not collected for Mountain View and Sheep Creek Reservoirs due to funding decisions.

Secure plants using water jet-stinger:

The two employees secured the willows by pushing the willows into the soft mud rather than using the water jet-stinger. They secured the willows in this fashion because of high water flows at time of planting. Later observations showed at least 90% success.

Maintain Vegetation:

Maintain generator at pump house for watering:

The laborers replaced the pressure tank at the Lake Billy Shaw pump house. They also filled the propane tank and ran the pump to keep it in operating condition. Cummins International sent a representative down from Boise to service the pump and generator units.

Water plants as needed:

One laborer would water trees at Lake Billy Shaw at least once per week, particularly during the hot season. Pre-existing plants and trees were watered and maintained at Mountain View and Sheep Creek Reservoir.

Trim plants as needed:

The fish technician and project manager trimmed trees at Lake Billy Shaw during the winter to promote better growth. The laborers trimmed trees of dead branches during the summer months.

Remove Vegetation:

Survey Lake Billy Shaw and associated streams for exotic species:

The fish technician and several laborers spent a couple of days looking at areas around LBS shoreline for Tamarisk and other exotics. They concentrated on areas that had Tamarisk plants present in 2006 surveys (Figure 3). No exotics were identified during these surveys. No surveys were conducted on Mountain View and Sheep Creek due to funding decisions.

Treat Tamarisk and other exotics with herbicide:

No Tamarisk plants were found during surveys.

Remove woody vegetation from Lake Billy Shaw dam:

The fish technician and the laborers spent several weeks chopping and burning brush of the Lake Billy Shaw dam. They also tried to clear vegetation around the toe-drains. This amount of work covered about half of the dam face. Rabbitbrush proved to be a persistent plant and new plants would begin to reestablish the area several weeks after clearing.

Figure 3: Tamarisk plant locations at Lake Billy Shaw in 2006

Operate and Maintain Habitat/Passage:

Maintain alternative water developments:

The lead laborer cleaned out two stock ponds southwest and two northwest of Lake Billy Shaw making them deeper to hold more water. He also placed small to medium sized rip rap at Coyote Hole to protect the impoundment from erosion and stock utilization.

Monitor and remove stock as needed:

The laborers occasionally spotted stock inside the perimeter fencing due to gates being left open. They removed the stock immediately or as soon as they could by ATV, vehicle, or by foot.

Maintain perimeter and inlet fencing at Lake Billy Shaw:

The perimeter fencing was surveyed several times throughout the year. Several stretches of fence were repaired at Coyote Hole by several laborers to prevent stock from entering the reservoir enclosure. In addition, several stretches of damaged fence were repaired at Sheep Creek Reservoir.

Put and Take Fisheries (Lake Billy Shaw):

Inspect dam and its associated structures weekly:

Lake Billy Shaw dam was inspected every two weeks. This inspection involved checking water levels and water clarity in the piezometer wells and checking conditions of the eight toe-drains.

Remove debris from fish screen structures:

The work crews checked and cleaned the fish screens and paddle-wheel structures as needed. This sometimes involved multiple inspections per day during high flow seasons.

Repair or replace paddle wheels at Boyle Creek:

The trash rack installed at Highline Canal was bent by ice and debris but is still in functioning condition. A second paddle-wheel was repaired with support arms between the paddles. This was installed in the early spring. The metal of the paddle wheel arms is soft and the supporting arms helped prevent the paddles from warping and breaking off.

Stock hatchery-reared Rainbow trout:

The fish suppliers delivered and stocked all loads of fish in the spring and fall months (Table 1).

Install 4 portable restrooms:

The crew installed 4 portable toilets at LBS in the first week of April.

Maintain 4 portable restrooms:

The crew periodically cleaned the restrooms and replaced toilet paper supplies. They also facilitated septic pumping through the Tribal water and sanitation department.

Remove 5 portable restrooms:

The 4 restrooms were pumped, removed, and stored for the winter.

Check boat ramps and maintain as needed:

The crew would inspect the boat ramps during water quality sampling as they utilized the ramps at these times. Neither repairs nor maintenance were needed.

Maintain roads around Lake Billy Shaw:

The lead laborer dug drainage ponds alongside the perimeter road at sections consistently flooded by pools of water. He also added gravel to a road section in the southwest portion. The local BIA road crew graded the roads in the spring. They also created widened road shoulders to protect the areas from wildfire due to the high number of damaging fires adjacent to the Reservation.

Maintain water pump at Lake Billy Shaw:

The crew started and used the water pump and generator periodically. The propane tank was filled and a service repair person came and serviced the Cummins/Onan pump and generator.

Remove aquatic vegetation:

The crew did not remove any aquatic vegetation at MVR due to time constraints and infeasible methods of removal. Their efforts most likely would not impact the reservoir significantly. This is a complex issue that has been looked at by other agencies for over 30 years.

Put and Take Fisheries (Sheep Creek Reservoir):

Stock SCR with hatchery-reared Rainbow trout:

The fish suppliers delivered and stocked all loads of fish in the spring and fall months (Table 1).

Removal of aquatic vegetation:

The crew did not remove any aquatic vegetation at MVR due to time constraints and infeasible methods of removal. Their efforts most likely would not impact the reservoir significantly. This is a complex issue that has been looked at by other agencies for over 30 years.

Maintain Campgrounds:

No maintenance on the campgrounds was performed due to FY07 funding decisions.

Table 1: Breakdown of Rainbow trout stocked at the three reservoirs

DATE	FISH FARM	RESERVOIR	WATER TEMP. (DEG)	AIR TEMP.	% OF HATCH?	OFF FEED FOR 48 HRS?	% LUNCH?	TOT. WT.	STERILE OR FERTILE?	CATCHABLE OR FINGERLINGS?	TOTAL COST
April 17, 2007	BCTF	SCR	10	15	Y	Y	N	8,000	F	Catchable	\$14,400
April 26, 2007	BCTF	LBS	11	12	Y	Y	N	8,000	S	Fingerlings	\$22,000
May 1, 2007	BCTF	LBS	12	14	N	Y	N	8,000	S	Catchable	\$16,000
May 4, 2007	BCTF	MVR	10	13	Y	Y	N	8,000	F	Catchable	\$14,400
May 10, 2007	BCTF	MVR	13	16	N	Y	N	8,000	F	Fingerlings	\$22,000
September 19, 2007	BCTF	LBS	52	55	2	Y	N	4,000	S	Catchable	\$8,000
September 20, 2007	CSI	MVR	59	62	Y	Y	N	3,300	F	Catchable	\$5,775
September 21, 2007	CSI	SCR	61	64	Y	Y	N	3,300	F	Catchable	\$5,775
September 26, 2007	BCTF	MVR	54	54	Y	Y	N	8,000	F	Catchable	\$14,400
September 27, 2007	CSI	SCR	59	64	Y	N	N	3,300	F	Catchable	\$5,775
September 28, 2007	CSI	SCR	61	57	Y	Y	N	3,300	F	Catchable	\$5,775
October 4, 2007	CSI	MVR	55	55	N	Y	N	3,300	F	Catchable	\$5,775
October 5, 2007	CSI	MVR	52	50	Y	Y	N	3,300	F	Catchable	\$5,775
October 8, 2007	CSI	MVR	57	55	Y	N	N	3,300	F	Catchable	\$5,775

Put and Take Fisheries (Mountain View Reservoir):

Stock MVR with hatchery-reared Rainbow trout:

The fish suppliers delivered and stocked all loads of fish in the spring and fall months (Table 1).

Removal of aquatic vegetation:

The crew did not remove any aquatic vegetation at MVR due to time constraints and infeasible methods of removal. Their efforts most likely would not impact the reservoir significantly. This is a complex issue that has been looked at by other agencies for over 30 years.

In regards to aquatic vegetation removal at all three reservoirs, Burge and Miller (1993) suggested several methods to manage the aquatic macrophytes in the DVIR reservoirs. These methods were mechanically removing plants using an aquatic weed harvester, using biological control measures such as grass carp (*Ctenopharyngodon idella*), or drawing down water to desiccate and kill the plants. Mechanical removal may prove to be costly in terms of labor and equipment costs. Grass carp may prove to be a poor choice as colder water temperatures decrease ingestion rates, these fish do not prefer milfoil and will forage on native species of

plants first, and risk of exotic establishment to other ecosystems runs high. Finally, water draw downs may decrease the reservoirs' ability to sustain trout fisheries. A final option may be the use of aquacides. However since fish from all three reservoirs are consumed by humans and other wildlife species and have a high affinity to absorb toxins within their flesh, this option is one that has many unknown risks and therefore is not being pursued at this time.

Maintain Campgrounds:

No maintenance on the campgrounds was performed due to FY07 funding decisions, except turning water and electricity on and off.

Collect/Generate/Validate Field and Lab Data:

Take photo-points of LBS shoreline and canal to monitor annual changes:

The fish technician and crew updated the photopoints in June. They are in digital format on the project manager's computer.

Take GPS locations of plants:

No GPS map was created this year because of no significant changes in plant density.

Record piezometer, toe-drain, and water levels at LBS:

The fish technician and the crew recorded piezometer, toe-drain, and water levels every two-weeks. Figure 6 shows changes in water levels at all three reservoirs. Typical spring water refills each reservoir to just over full pool status. Over the next few months the water slowly seeps or evaporates, reducing the overall amount of water in each reservoir. During the fall months, the irrigation district may put additional but not always sufficient water into the reservoirs before winter freeze-over.

Figure 6: Changes in water levels over time.

Monitor water quality at reservoirs:

The fish technician and laborers sampled water quality every two weeks using an instrument built by YSI. The primary characters sampled were dissolved oxygen, temperature, pH, turbidity, and chlorophyll-a. Dissolved oxygen and temperatures are the two main limiting factors of Rainbow trout in the three reservoirs. They are shown in the following six charts.

The 2007 spring and summer were quite warm and had very little precipitation and cloud cover. These dry, hot summers make it very hard on the fisheries because of high water temperatures and increased biological oxygen demand. Furthermore, hot dry weather causes excessively high demand for water by ranchers and other irrigators. This high demand makes it difficult to obtain water for the reservoirs. Another problem associated with long periods of hot weather and sunshine is the potential for algae blooms to occur. These algae blooms can prove to be very detrimental to lentic fisheries. An example of this occurred in a large water impoundment adjacent to the Duck Valley Indian Reservation. Wildhorse Reservoir, managed by the Nevada Division of Wildlife, experienced an algae bloom towards the end of August. This algae bloom resulted in very low dissolved oxygen levels throughout the water column and a subsequent large fish kill (http://www.ndow.org/about/news/pr/082907_Wildhorse.shtml). See attached article in appendix.

Figure 7 (a-f): Temperatures and dissolved oxygen for the three reservoirs.

For such a hot, dry summer the temperatures at LBS surprisingly never exceeded 25 Celsius. This may be attributed water being delivered to the reservoir by the Tribal irrigation department during the hottest portions of the summer. Dissolved oxygen levels became critical only in the deepest depths of water from the end of July to the end of August. DO levels in the first three meters of water stayed at adequate levels throughout the entire year. Rainbow trout were excluded from depths 4 meters and greater from the surface from the end of July towards the end of August.

Mountain View Reservoir exhibits similar seasonal changes at LBS. Water temperatures did not exhibit stratification as much as the other two reservoirs. The water in the hypolimnion (deeper than 3 meters) became somewhat anoxic during mid-July through early August. Water shallower than 3 meters stayed well above critical dissolved oxygen levels. This indicates that the late summer months were not as limiting to trout as seen in previous years and trout were only excluded from the hypolimnion for approximately a month.

Sheep Creek Reservoir exhibited similar stratification and seasonal shifts as the other two reservoirs. However, the surface water temperatures were highly variable and may be attributed to the highly turbid characteristics of the water (see figure 8). Because the water is turbid, sunlight is mostly absorbed at the surface and does not penetrate into deeper depths easily. Therefore, heat energy associated with sunlight affects the first meter of water more drastically than deeper depths. Dissolved oxygen seasonal changes were similar to the other reservoirs. Hypolimnetic water became anoxic from mid-July through the end of August, while more shallow depths maintained levels suitable for trout survival.

Figure 8: Turbidity measurements of the three reservoirs in 2007.

Turbidity measurements (Figure 8) at the three reservoirs show that Mountain View and Lake Billy Shaw had much clearer water than Sheep Creek. Initial field surveys, conducted by Brown and Caldwell (Boise Office), and soil/water analysis, conducted by the Environmental Science Corporation in Juliet, Tennessee, provided some insights to the turbidity problem. Soil and water samples showed that the particles contributing to turbidity are not algal but geological in nature. The particles largely consist of extremely small clay smectite minerals which may be negatively charged in nature and surrounded by clouds of positively charged ions. These clouds of positive ions magnetically repel other similar particles and preclude aggregation. The smectite clay minerals may settle out of the water column over several years but the shallowness of Sheep Creek, the repelling charged ions around the particles, and the various disturbing factors of water runoff, wind action, and boat propellers may negate the ability of these particles to settle out. Brown and Caldwell found that the clay minerals may be added by all shoreline areas but a sample collected from the western bluffs adjacent to the reservoirs closely resembled the particle composition in the water indicating that surface runoff may be the major contributor to the turbidity. Initial mitigation recommendations include increasing the vegetative buffer areas along the shoreline, further study to understand the usefulness of adding flocculants, and implementing utilization of flocculants depending on the study results (Kerner and Gagnon 2007).

Lake Billy Shaw experienced a spike in turbidity in August which may be attributed to free floating algae growth. Mountain View also experienced elevated turbidity in September and again may be attributed to algae growth and proliferation. In both reservoirs, the increase in turbidity was concurrent with an increase in chlorophyll-a levels.

Record high detail water quality:

The fisheries technician and the crew assisted the local TEPP department with high detail water quality sampling. The TEPP released the water quality details to the project (Table 2).

Table 2: High detailed water quality sampling performed by TEPP in 2007

Sample ID	Latitude	Longitude	Date Sampled	NO3/NO2 mg/L	NO3 mg/L	NO2 mg/L	Ortho-P mg/L	T-Phos mg/L	NH3 mg/L	TKN mg/L	CO3 mg/L	HCO3 mg/L	SO4 mg/L	Cl mg/L
MVR N Inlet	42.02.30	116.09.12	7/30/2007	< 0.01	< 0.01	< 0.01	< 0.003	0.11	0.03	1.56	197	67	5.8	6
MVR @ Middle	42.02.63	116.09.51	7/30/2007	< 0.01	< 0.01	< 0.01	< 0.003	0.055	0.04	1.21	22	56	5.9	5.9
MVR Near Dam	42.03.23	116.10.23	7/30/2007	< 0.01	< 0.01	< 0.01	< 0.003	0.025	0.02	1	10.1	90	6.2	6
LBS Near Dam	41.58.36	116.14.34	7/30/2007	< 0.01	< 0.01	< 0.01	< 0.003	0.05	0.02	1.83	188	88	7.7	4.9
LBS Near Middle	41.57.84	116.14.52	7/30/2007	< 0.01	< 0.01	< 0.01	< 0.003	0.055	0.02	1.82	1	88	7.8	4.9
LBS Near Inlet	41.58.40	116.14.53	7/30/2007	< 0.01	< 0.01	< 0.01	0.003	0.055	0.02	1.83	0	88	8.0	4.1
LBS Near Inlet	41.58.40	116.14.53	7/30/2007	< 0.01	< 0.01	< 0.01	0.003	0.055	0.02	1.83	0	88	8.0	4.1
SCR Near Inlet	41.52.83	116.15.52	7/30/2007	0.02	0.01	0.01	0.024	0.11	0.04	0.99	0.56	99	10.3	4.2
SCR N Middle	41.51.88	116.15.87	7/30/2007	0.04	0.02	0.02	0.029	0.1	0.08	0.83	0	101	11.2	4.2
SCR Near Dam	41.51.50	116.16.14	7/30/2007	0.04	0.02	0.02	0.032	0.12	0.09	0.86	0	101	11.2	4.2
Sample ID	Ca mg/L	Mg mg/L	Na mg/L	K mg/L	Hardness mg/L	TDS mg/L	SiO2 mg/L	Alkalinity mg/L	Fecal col/100mL	E. coli col/100mL	TSS mg/L	Lab pH SU	Lab EC uS/cm	Turbidity NTU
MVR N Inlet	13.8	7.9	15.1	4	67	114	7.2	87.8	12	< 2	20	9.4	198	14
MVR @ Middle	12.4	7.6	15	3.5	62	115	12.5	84.3	2	< 2	87	9.5	193	26
MVR Near Dam	15	7.8	14.5	3.9	70	120	9.4	90.7	< 2	< 2	4	9.1	206	6
LBS Near Dam	17.3	4.8	8.1	2.8	84	111	8.2	75.4	< 2	< 2	8	8.1	178	22
LBS Near Middle	18	4.8	8.1	2.8	88	112	8.8	76.1	< 2	< 2	8	8.1	179	8
LBS Near Inlet	17.2	6.1	11.3	3.1	84	112	8.3	76.4	< 2	< 2	2	8.2	180	4
LBS Near Inlet	17.4	6.1	11.3	3.1	84	112	8.4	76.4	< 2	< 2	2	8.1	181	4
SCR Near Inlet	20	4.8	10.3	4.1	70	128	22.1	82.8	< 2	< 2	8	8.4	199	32
SCR N Middle	20.6	5	10.3	4.3	72	129	22.4	82.8	< 2	< 2	3	8	198	25
SCR Near Dam	20.4	5	10.2	4.3	72	129	22.4	82.8	4	< 2	4	8	199	25
Sample ID	Hg ug/L	Sam. Depth feet	Field Temp C	DO mg/L	Field pH SU	Field EC uS/cm	ORP mv	BP mm Hg	Field NTU					
MVR N Inlet	< 0.2	3.3	21.8	7.1	8.16	200	149	628						
MVR @ Middle	< 0.2	3.3	21.7	6.8	9.42	195	92	628	4					
MVR Near Dam	< 0.2	3.3	22.9	8.6	8.93	207	75	628						
LBS Near Dam	< 0.2	3.3	22.4	8.7	8.23	193	89	628	12					
LBS Near Middle	< 0.2	3.3	22.7	11.3	8.44	194	89	628						
LBS Near Inlet	< 0.2	3.3	22.8	10.4	8.38	198	200	628	12					
LBS Near Inlet	< 0.2	3.3	22.8	10.4	8.38	198	200	628	12					
SCR Near Inlet	< 0.2	3.3	24.1	8.2	7.85	203	186	631	21					
SCR N Middle	< 0.2	3.3	23.9	6.8	7.7	203	1	628						
SCR Near Dam	< 0.2	3.3	22.8	7.6	7.47	203	170	628						

Conduct wildlife surveys: Informal bird surveys were conducted but no new species were identified. However, a hummingbird feed at the project manager's residence showed three hummingbird species (Rufous, Black-chinned, and Broad-tailed) to be present. This location is near Lake Billy Shaw (Table 3). No new wildlife species were identified at LBS (Table 4).

Table 3: Bird species identified at LBS to date.

Common Name	Scientific Name	Common Name	Scientific Name
1 American Avocet	<i>Recurvirostra americana</i>	36 Killdeer	<i>Charadrius vociferans</i>
2 American Coot	<i>Fulica americana</i>	38 Lesser Scaup	<i>Anas platyrhynchos</i>
3 American Hareling	<i>Fulica macrura</i>	37 Merganser	<i>Lophodytes cucullatus</i>
4 American White Pelican	<i>Pelecanus erythrorhynchos</i>	39 Marsh Wren	<i>Chondestes schach</i>
5 American Wren	<i>Troglodytes aedon</i>	38 Northern Harrier	<i>Circus hudsonius</i>
6 Bald Eagle	<i>Haliaeetus leucocephalus</i>	40 Northern Flicker	<i>Colaptes auratus</i>
7 Black-bellied Plover	<i>Pluvialis dominica</i>	41 Northern Shrike	<i>Lanius borealis</i>
8 Black-bellied Plover	<i>Pluvialis dominica</i>	42 Osprey	<i>Pandion haliaetus</i>
9 Black-crowned Night Heron	<i>Nycticorax nycticorax</i>	43 Red-bellied Diver	<i>Colymbus auratus</i>
10 Black-necked Stilt	<i>Himantopus mexicanus</i>	44 Purple Finch	<i>Spizella socialis</i>
11 Bonaparte Gull	<i>Arenaria interpres</i>	45 Common Raven	<i>Corvus corax</i>
12 Bonaparte Gull	<i>Arenaria interpres</i>	46 Redhead	<i>Colinus virginianus</i>
13 Bonaparte Gull	<i>Arenaria interpres</i>	47 Red-tailed Hawk	<i>Bubo virginianus</i>
14 Bonaparte Gull	<i>Arenaria interpres</i>	48 Red-tailed Hawk	<i>Bubo virginianus</i>
15 Bonaparte Gull	<i>Arenaria interpres</i>	49 Red-tailed Hawk	<i>Bubo virginianus</i>
16 Bonaparte Gull	<i>Arenaria interpres</i>	50 Red-tailed Hawk	<i>Bubo virginianus</i>
17 Bonaparte Gull	<i>Arenaria interpres</i>	51 Red-tailed Hawk	<i>Bubo virginianus</i>
18 Bonaparte Gull	<i>Arenaria interpres</i>	52 Red-tailed Hawk	<i>Bubo virginianus</i>
19 Bonaparte Gull	<i>Arenaria interpres</i>	53 Red-tailed Hawk	<i>Bubo virginianus</i>
20 Bonaparte Gull	<i>Arenaria interpres</i>	54 Red-tailed Hawk	<i>Bubo virginianus</i>
21 Bonaparte Gull	<i>Arenaria interpres</i>	55 Red-tailed Hawk	<i>Bubo virginianus</i>
22 Bonaparte Gull	<i>Arenaria interpres</i>	56 Red-tailed Hawk	<i>Bubo virginianus</i>
23 Bonaparte Gull	<i>Arenaria interpres</i>	57 Red-tailed Hawk	<i>Bubo virginianus</i>
24 Bonaparte Gull	<i>Arenaria interpres</i>	58 Red-tailed Hawk	<i>Bubo virginianus</i>
25 Bonaparte Gull	<i>Arenaria interpres</i>	59 Red-tailed Hawk	<i>Bubo virginianus</i>
26 Bonaparte Gull	<i>Arenaria interpres</i>	60 Red-tailed Hawk	<i>Bubo virginianus</i>
27 Bonaparte Gull	<i>Arenaria interpres</i>	61 Red-tailed Hawk	<i>Bubo virginianus</i>
28 Bonaparte Gull	<i>Arenaria interpres</i>	62 Red-tailed Hawk	<i>Bubo virginianus</i>
29 Bonaparte Gull	<i>Arenaria interpres</i>	63 Red-tailed Hawk	<i>Bubo virginianus</i>
30 Bonaparte Gull	<i>Arenaria interpres</i>	64 Red-tailed Hawk	<i>Bubo virginianus</i>
31 Bonaparte Gull	<i>Arenaria interpres</i>	65 Red-tailed Hawk	<i>Bubo virginianus</i>
32 Bonaparte Gull	<i>Arenaria interpres</i>	66 Red-tailed Hawk	<i>Bubo virginianus</i>
33 Bonaparte Gull	<i>Arenaria interpres</i>	67 Red-tailed Hawk	<i>Bubo virginianus</i>
34 Bonaparte Gull	<i>Arenaria interpres</i>	68 Red-tailed Hawk	<i>Bubo virginianus</i>

Table 4: Wildlife species identified at LBS to date

Common Name	Scientific Name
1 American Badger	<i>Taxidea taxus</i>
2 American Beaver	<i>Castor canadensis</i>
3 Coyote	<i>Canis latrans</i>
4 Deer mouse	<i>Peromyscus maniculatus</i>
5 Least Chipmunk	<i>Tamias minimus</i>
6 Long-tailed Weasel	<i>Mustela frenata</i>
7 Meadow vole?	<i>Microtus pennsylvanicus</i>
8 Montane vole	<i>Microtus montanus</i>
9 Mountain (Nuttall's) Cottontail	<i>Sylvilagus nuttallii</i>
10 Mule Deer	<i>Odocoileus hemionus</i>
11 Muskrat	<i>Ondatra zibethicus</i>
12 Night Snake	<i>Hypsiglena torquata</i>
13 Northern grasshopper mouse?	<i>Onychomys leucogaster</i>
14 Pronghorn	<i>Antilocapra americana</i>
15 Racer	<i>Coluber constrictor</i>
16 Sagebrush Lizard	<i>Sceloporus graciosus</i>
17 Striped Whipsnake?	<i>Masticophis bilineatus</i>
18 Western Fence Lizard	<i>Sceloporus occidentalis</i>
19 Western Rattlesnake	<i>Crotalis viridis</i>
20 White-tailed Jackrabbit	<i>Lepus townsendii</i>

Monitor fisheries at all reservoirs:

The three reservoir fish populations were monitored through length and weight indices. Length and weight data on Rainbow trout were collected through two means, gill netting and creel surveys.

Figure 9a: Length frequencies by percent of trout at LBS

At Lake Billy Shaw, the length frequency histogram (Figure 9a) shows the reservoir is growing fish into good length ranges for trophy status. The small percentages of the stocked sized trout indicate that recently stocked fish grew into larger size classes quickly. The greatest percent of fish caught were in the 15 to 17.9 inch fish. This is good because as a trophy fishery anglers will catch fish in these size classes about 60 to 65% of the time. Many fish in LBS are also making it into the 19 to 21 inch range. About 15% of the fish caught were 19 to 20.9 inches in length. One concern is the lack of fish greater than 21 inches but this may be attributed to lack of adequate sample size during gill netting. Angler reports also supported this lack of very large trout. However, some anglers reported catching fish greater than 22 inches. During gill net

sampling in the fall fewer numbers of trout were caught than wanted during. A second sample was taken but only resulted in two additional trout.

Figure 9b: Average relative weight (Wr) values by length class at LBS

Relative weight values (Figure 9b) at LBS showed that trout were slightly less rotund than what the standard trout is. The extreme values >100 and <80 may be attributed to small sample size because only one or two individual fish in those length groups were caught. Overall, the relative weight values indicate that the trout in LBS are less rotund than they should be. This is believed to be caused by a slight overstocking of the reservoir. The exponential value shown in Figure 9c (2.3144) also supports the overstocking hypothesis. Values of 3.000 indicate fish grow isometrically. Values less than 3.000 indicate that fish are becoming longer faster than putting on weight.

Figure 9c: Length-weight regression of trout at LBS

Overall fish at LBS are still healthy despite the lower relative weight averages and the low length-weight regression exponent. Anglers reported fish that were strong and hard fighters.

Neither lesions nor signs of poor health were reported by anglers. Also, signs of bad health were not seen in any of the gill net sample fish.

Mountain View length frequency histogram showed that the reservoir produced very well in 2007 (Figure 10a). Many fish were living and growing into larger size classes despite the high harvest rates on the reservoir. Of the harvested fish at MVR, 34 percent were 13 to 13.9 inches, 33 percent were 14 to 15.9 inches, and about 17 percent were 16 to 20.9 inches. The high catch rates for 14 to 20.9 inch fish ensures that anglers visiting the reservoir will have very good chances at catching nice sized fish. One other pleasant aspect shown by the histogram is the 2 percent of fish caught being greater than 24 inches in length.

Figure 10a: Length frequency histogram at MVR

Relative weight values (Figure 10b) for trout in Mountain View show that these trout are in excellent condition up to 55 cm (~22 inches) in length. The average W_r values for fish shorter than 55 cm are greater than 90. Average W_r values for fish shorter than 35 cm (14 inches) are all over 100. These values indicate that the fish are in superb condition and there are no factors limiting their condition. These high values indicate that the reservoir would be able to handle increased fish stocking.

Figure 10b: Average relative weight values by length class at MVR

The length-weight regression of Rainbow trout at MVR also supports the idea that more fish may be stocked into the reservoir. The exponent value of 2.5885 was increased from 2006 (2.4738) but decreased from 2005 (2.6011). The variation in the L-W regression exponent may reflect varying environmental conditions but the fairly high value for the three years indicates that the fish grew somewhat isometrically. The high amount of harvest in MVR reduced the amount of intraspecific competition thus allowing more fish to be stocked.

Figure 10c: Length-Weight regression of trout at MVR

The length frequency histogram for Sheep Creek Reservoir (Figure 11a) provides information that is disconcerting. The frequencies indicate that fish 11 to 13.9 inches are the only size of fish in the reservoir. Very few fish or no fish 14 inches or larger were caught by anglers. In 2005

and in 2006 there were a significant number of fish 15 to 15.9 inches caught. There should be a large percent of fish that carry over into larger length groups in 2007 but there is not. The absence of fish larger than 13.9 indicates that there is a large mortality occurrence in fish soon after stocking. The turbidity problem associated with SCR is believed to be the main limiting factor for trout. The turbidity may affect the trout in several ways. First, the turbidity may reduce the ability of trout to affectively hunt for food as visibility is reduced to several feet. Second, the poor light penetration through the water column reduces the reservoir's ability to grow phytoplankton and subsequent higher food levels such as zooplankton and macroinvertebrates. Finally, the suspended clay minerals may negatively impact fish physiological processes and organs. For example, gills may be constantly irritated by the particles and gas exchange may be compromised.

Figure 11a: Length frequency histogram at SCR

The average relative weight values of trout in SCR (Figure 11b) were somewhat perplexing when first looked in numerical form. Because of the turbidity problems of the water quality, it was assumed that the W_r values would be fairly low. However, when these data were plotted on a regression chart, the reason for the high W_r values presented itself. All of the fish caught were recently stocked fish (11 to 13.9 inches) and the high W_r values may be attributed to the optimal growing conditions associated with raceway rearing.

The length-weight regression (Figure 11c) also reflects the relatively good condition (2.5915) of fish but also reflects that mainly stock sized fish were caught. Both the W_r and L-W regression emphasize the distressing occurrence of fish mortality.

Figure 11b: Average relative weight values by length class at SCR

Figure 11c: Length-weight regression of trout at SCR

The creel data collected by the employees provided an overview of the fishing pressure, success, and enjoyment. Table 5 shows estimates of angler-days, effort, harvest, and fishing rating. This table also shows a breakdown where anglers are coming from and how these anglers learned about the fishing at Duck Valley Indian Reservation.

Mountain View Reservoir had the highest number of angler days (2,151) and effort (8,615 hours). The number of anglers counted on the reservoirs is expressed in angler-days because some of the anglers stay for multiple days. MVR also had the highest number of fish harvested (3,285 fish) and a rating of 3.09. The fishing rating was based on a five number scale (1 is very bad, 2 is poor, 3 is fair, 4 is good, and 5 is excellent) and MVR had an average of 3.09. Lake Billy Shaw had a angler-day count of 1,060 with a total estimated effort of 5,495 hours of effort and 7 fish harvested which reflects the catch-and-release regulations. Fishing experience rating at LBS indicated that anglers enjoyed themselves the most at LBS. Sheep Creek Reservoir received the least amount of fishing pressure and may be attributed to the turbid water and lack of larger fish. This historically well-known fishery only received 447 angler-days and 1,849 hours of effort. Only 490 fish were harvest from the reservoir and anglers rated the fishing between poor and fair. For all three fisheries, most visitors came from Idaho or Nevada. Large numbers came from California and Oregon and lesser numbers came from Utah, Washington, and other states. One foreigner from Romania was surveyed on the reservoirs. Most anglers knew about Duck Valley Indian Reservation just by being here before but many first time visitors learned about the area through friends and family.

Table 5: Creel data summary and averages

Reservoir	Count of Angler Days	Total Estimated Effort	Estimated Harvest	Fishing experience rating
LBS	1060	5495.3	7	3.37
MVR	2151	8615.4	3285	3.09
SCR	447	1849.4	490	2.43
State	Count Of State	Learning Pathway	Number of Anglers	
California	56	Been Here Before	433	
Foreign Country	1	DMR Website	11	
Idaho	740	Family Member	16	
Nevada	148	Friend	53	
		Magazine		
Oregon	41	Advertisement	9	
Other State	16	Newspaper Article	1	
Utah	18	Other	2	
Washington	9	Sporting Good Store	6	

Produce Plan:

Work with regional biologists and CBFWA to update plans:

No Resident Fish Committee project review and update meetings were held nor attended.

Produce Pisces Status Report:

Status reports for the twelve months were completed and submitted to BPA.

Produce Annual Report:

Submit draft of FY07 annual report to BPA for review:

The FY07 annual report rough draft was worked on and completed in October.

Finalize FY07 annual report and post on BPA's website:

The finalized version of the FY07 annual report was submitted and posted on BPA's website.

References:

- Anderson, R.O. and R.M. Neumann. 1996. Length, weight, and associated structural indices. Pages 454-463 in B. R. Murphy and D. W. Willis, editors. Fisheries techniques, 2nd edition. American Fisheries Society, Bethesda, Maryland.
- Kerner, D.A. and M.E. Gagnon. 2007. Sheep Creek Reservoir initial turbidity analysis: Duck Valley Indian Reservation, Owyhee, Nevada. Brown and Caldwell, Boise Office.
- Burge, H.L. and W. H. Miller. 1993. Fishery Management on the Duck Valley Indian Reservation, Shoshone-Paiute Tribe. USFWS 1992 Annual Report. 12 pp.
- Ney, J.J. 1999. Practical Use of Biological Statistics. Pages 171-172 in C.C. Kohler and W.A. Hubert, editors. Inland fisheries management in North America, 2nd edition. American Fisheries Society, Bethesda, Maryland.
- Cooperrider, A.Y., R.J. Boyd, and H.R. Stuart, eds. 1986. Inventory and monitoring of wildlife habitat. U.S. Dept. Inter., Bur. Land Manage. Service Center, Co. xviii, 858pp.
- Bookhout, T.A., Editor. 1996. Research and management techniques for wildlife and habitats. Fifth ed., rev. The Wildlife Society, Bethesda, Md. 740pp.

Appendix:

About NDOW

News & Media

[Press Releases](#) | [On the Wildside of Life - Radio & TV](#)

Date: 8/29/07

Contact: Joe Doucette

Phone: (775) 777-2305

WILDHORSE RESERVOIR FISH KILL

Nevada Department of Wildlife fisheries biologist Chris Drake discovered a major fish die-off when he traveled to Wildhorse Reservoir after receiving a call from a concerned angler regarding dead fish at the popular fishery.

"We estimate that approximately 8,000 to 11,000 trout and 750 to 1,000 perch died over the last couple of days," says Drake. "We believe it is due to the lack of dissolved oxygen within the water."

While there have been occasional smaller fish kills over the last few years during the heat of the summer, Drake says that this die-off is more significant than is normal. The problem appears to be related to several factors.

The combination of unusually hot weather and more than the usual number of sunny days this summer has produced above normal algae growth in the lake. As algae grows, oxygen is released into the water for aquatic organisms to use. However, when the algae dies, as it did just recently, and settles to the bottom and decomposes, it becomes anaerobic and the oxygen is depleted as the dead plant matter breaks down.

"The dissolved oxygen (DO) level was 1.1 to 1.6 parts per million (ppm) on the surface and 0.7 to 1.4 ppm on the bottom when I tested on Monday," said Drake. "Trout require DO levels of 3.0 ppm or higher for normal respiratory function."

Coupled with the low levels of DO are low lake levels and warm water, which doesn't hold as much dissolved oxygen to begin with. The end result was high stress to the trout without suitable habitat and a significant fish kill occurred. Drake estimates that the majority of the fish washing up on shore were trout (approximately 95%), consisting of rainbow, tiger and a few brown trout. The average size of the trout was 16 to 17 inches, with fish up to 22 inches being found dead in deeper water. Some yellow perch were also found to be stressed and dying, with the average size being around 9 inches.

With the large number of fish lining the shorelines, anglers and campers can expect conditions to be extremely smelly due to the decomposition of the fish. NDOW will continue to monitor the extent of the fish kill and will perform population surveys later this fall once the weather cools down.

Drake says that fish may continue to die if current weather conditions don't change and that fishing will be poor over the next two to three weeks until temperatures start to cool down as fish are in survival mode and not eating. He also wants to remind anglers that all fishing regulations are still in effect and that restocking of desired game fish will resume once water quality levels are safe and productive and will be determined by current reservoir storage and anticipated runoff in 2008.

Survival of smaller fish in the reservoir is expected to be fair to good as low DO fish kills generally target the larger fish due to their higher metabolic rate and greater oxygen demands. NDOW records indicate that in an average year approximately 100,000 trout are stocked into Wildhorse Reservoir.

The Nevada Department of Wildlife (NDOW) protects, restores and manages fish and wildlife, and promotes fishing, hunting, and boating safety. NDOW's wildlife and habitat conservation efforts are primarily funded by sportsmen's license and conservation fees and a federal surcharge on hunting and fishing gear. Support wildlife and habitat conservation in Nevada by purchasing a hunting, fishing, or combination license. For more information, visit www.ndow.org.