Hood River Steelhead Genetics Study

Relative Reproductive Success of Hatchery and Wild Steelhead in the Hood River

Final Report 2002 - 2003

May 2003

DOE/BP-00009245-1

This Document should be cited as follows:

Blouin, Michael, "Hood River Steelhead Genetics Study; Relative Reproductive Success of Hatchery and Wild Steelhead in the Hood River", 2002-2003 Final Report, Project No. 198805312, 27 electronic pages, (BPA Report DOE/BP-00009245-1)

Bonneville Power Administration
P.O. Box 3621
Portland, OR 97208

This report was funded by the Bonneville Power Administration (BPA), U.S. Department of Energy, as part of BPA's program to protect, mitigate, and enhance fish and wildlife affected by the development and operation of hydroelectric facilities on the Columbia River and its tributaries. The views in this report are the author's and do not necessarily represent the views of BPA.

Hood River Steelhead Genetics Study

Relative Reproductive Success of Hatchery and Wild Steelhead in the Hood River

Final report on work conducted under

**BPA Intergovernmental Contract Number 00009245,
Project Number 1988-053-12**

and

ODFW Interagency Agreement Number 001-2007s

OSU Technical Contact and Principal Investigator

Dr. Michael Blouin, Associate Professor
Dept. Zoology, Oregon State University
Corvallis, OR 97331
Tel: 541-737-2362
Fax: 541-737-0501
blouinm@science.oregonstate.edu
Date: 5/1/03

OVERVIEW

Motivation for the project

There is a considerable interest in using hatcheries to speed the recovery of wild populations. The Bonneville Power Administration (BPA), under the authority of the Northwest Power Planning Act, is currently funding several hatchery programs in the Columbia Basin as off-site mitigation for impacts to salmon and steelhead caused by the Columbia River federal hydropower system. One such project is located on the Hood River, an Oregon tributary of the Columbia. These hatchery programs cost the region millions of dollars. However, whether such programs actually improve the status of wild fish remains untested. The goal of this project was to evaluate the effectiveness of the Hood River hatchery program as required by the Northwest Power Planning Council Fish and Wildlife Program, by the Oregon Plan for Coastal Salmonids, by NMFS ESA Section 4(d) rulings, and by the Oregon Department of Fish and Wildlife (ODFW) Wild Fish Management Policy (OAR 635-07-525 through 529) and the ODFW Hatchery Fish Gene Resource Management Policy (OAR 635-07-540 through 541).

This work was funded by the Bonneville Power Administration through BPA Intergovernmental Contract 9245 (Project # 1988-053-12), and through ODFW Interagency agreement No. 001-2007s.

Brief summary of project and results

The Hood River supports two populations of steelhead, a summer run and a winter run. They spawn only above the Powerdale Dam, which is a complete barrier to all salmonids. Since 1991 every adult passed above the dam has been measured, cataloged and sampled for scales. Therefore, we have a DNA sample from every adult steelhead that went over the dam to potentially spawn in the Hood River from 1991 to the present. Similar numbers of hatchery and wild fish have been passed above the dam during the last decade. During the 1990's "old" domesticated hatchery stocks of each run (multiple generations in the hatchery, out-of-basin origin; hereafter H_{old}) were phased out, and conservation hatchery programs were started for the purpose of supplementing the two wild populations (hereafter "new" hatchery stocks, H_{new}). These samples gave us the

unprecedented ability to estimate, via microsatellite-based pedigree analysis, the relative total reproductive success (adult-to-adult production) of hatchery (H_{old} or H_{new}) and wild (W) fish for two populations, over multiple brood years. Our analyses of samples from fish that bred in the early to mid 1990's show that fish of "old" hatchery stocks have much lower total fitness than wild fish (17% to 54% of wild fitness), but that "new" stocks have fitness that is similar to that of wild fish (ranging from 85% to 108% of wild fitness, depending on parental gender and run year). Therefore, our results show that the decision to phase out the old, out-of-basin stocks and replace them with new, conservation hatchery stocks was well founded. We also conclude that the H_{new} fish are leaving behind substantial numbers of wild-born offspring. The similar fitnesses of H_{new} and W fish suggests that wild-born offspring of H_{new} fish are unlikely to have negative genetic effects on the population when they in turn spawn in the wild. We will test this hypothesis once enough F2 offspring have returned. Another interesting result is that we were unable to match a large fraction of the unclipped, returning fish with parents from their brood year. Furthermore, we were missing more fathers than mothers. Because we sampled almost every possible anadromous parent, these results suggest that non-anadromous trout or precocious parr may be obtaining a substantial number of matings. Substantial reproduction by precocious parr could be one unintended consequence of the hatchery program.

Personnel and project Coordination:

The genetics pedigree work was carried out at Oregon State University by Drs. M. Blouin and W. Ardren (formerly a postdoc in Dr. Blouin's laboratory, now at the USFWS Abernathy Fish Genetics lab). This project was coordinated with the Hood River steelhead hatchery and research project, funded by Bonneville Power Administration and administered and implemented by the Oregon Department of Fish and Wildlife (Rod French and Erik Olsen, supervisor and database manager). The project was also coordinated with Dr. Thomas E. Morse, Bonneville Power Administration (BPA), Fish and Wildlife Division, and with Kathryn Kostow, ODFW.

INTRODUCTION

Using hatcheries to enhance wild populations

Hatchery supplementation is the use of captive propagation to increase the size of an endangered wild population (also known as supportive breeding or as a conservation hatchery program; Ryman et al., 1995). The concept is to take part of a small wild population into captivity, protect their offspring through the high-mortality early life stages, and then allow them to spawn naturally when they return as adults. The return of their offspring should increase the size of the wild population. Whether such supplementation programs actually work (increase the size of wild populations) is not clear (Waples 1999). While we may observe that a wild population's size increases or decreases after hatchery adults were allowed to spawn in the same stream, we have not been able to determine which fish among the spawners actually produced returning adults. It is possible that the hatchery fish are making little contribution to the next generation, or even worse, are dragging down the fitness of the wild population for genetic or ecological reasons (Waples 1991; 1999; Fleming and Petersson, 2001; Lynch and O'Hely, 2001). On the other hand, the hatchery fish may indeed be leaving behind offspring that return to breed successfully and contribute to the next generation. Until now we have had no way of determining which of these scenarios is true.

Background on the Hood River basin and stocks

The Hood River supports wild runs of winter and summer steelhead. Breeding areas for winter and summer fish are segregated, with summer fish breeding in the West Fork of the Hood River and winter fish breeding in the remaining tributaries (Fig. 1). The Powerdale Dam at mile 4.0 on the river is a complete barrier to migrating salmon. Facilities include an adult trap and sorter built by BPA. The trap is used for all broodstock collection, for monitoring hatchery and wild adults, and for controlling entry of hatchery fish into natural production areas (a photo of the dam and of the inside of the fish handling facility can be seen at http://oregonstate.edu/~blouinm/Hood%20RiverProject_files/slide0001.htm). This facility provides the unique opportunity to handle the entire population of returning adults every year. Since 1991 every adult passed over the dam has been catalogued, measured

and sampled for scales. Traps for sampling juveniles have been in place in the main stem and at the outlets of all the main tributaries since the mid 1990s. The dam is scheduled to be removed in 2010, although the actual date has not yet been determined. The dam is being kept in place until that time in order to facilitate several ongoing research projects in the basin, including this one.

Winter run

Winter run hatchery stock 13 (a domesticated, out-of-basin, multi-generation hatchery stock from Big Creek) was previously stocked in the basin but was phased out in 1991. It was replaced by conservation hatchery stock 50, which uses wild Hood River broodstock each generation and was implemented for the purposes of supplementing the wild winter population. The first generation of stock 50 adults began returning in appreciable numbers in 1995 (Fig. 2). Since then the number of H_{new} fish passed above the dam has been limited to no more than the number of wild fish passed (Table 1). This protocol created an ideal opportunity to evaluate the relative reproductive success of each type of fish spawning in the wild.

Summer run

Summer run hatchery stock 24 (a domesticated, out-of-basin, multigeneration hatchery stock from Skamania) was phased out in 1998 and replaced by summer conservation hatchery stock 50 in 1998. The protocols for this summer-run supplementation program are the same as for the winter run program. Skamania stock 24 are still planted below the dam to provide a sports fishery, but none are allowed above the dam.

Here we use the abbreviation H_{old} to refer to “old” hatchery stocks 24 and 13, and H_{new} to refer to the “new” conservation hatchery winter stock 50 and summer stock 50. Run years begin in the fall and breeding occurs in the spring and summer of the following year. Thus, for example, parents of run year 1991-1992 produce offspring whose brood year (date of birth) is 1992. For simplicity’s sake we will, for example, refer to 91 as both the parental run year and offspring birth year.

Figure 1. Study site. Powerdale dam is a complete barrier to salmonids at mile 4.0. Summer steelhead breed in the West Fork, while winter steelhead breed in the Middle Fork, East Fork, and Neal Creek. Juvenile traps are located just above the dam and at the base of each of the main branches within the system.

Table 1. Summary of numbers of wild (unclipped) and hatchery (clipped) fish passed above Powerdale Dam. These counts do not include fish taken for broodstock, which we also genotyped. Run year refers to the fall season (e.g. the 91-92 run year fish began arriving fall of 1991 and continued arriving into 1992). In 1992 the H_{old} stock 13 program was phased out for winter run, and the winter run conservation hatchery program was begun. Those H_{new} winter stock 50 fish began returning to spawn in the wild in 94-95. The old summer stock 24 program was phased out and the new summer run conservation hatchery program was begun in 1998. Those H_{new} summer fish began returning in 01-02. Highlighted cells have been genotyped. The rest will be typed as part of the work proposed here. The key parental years for which we have preliminary comparisons of hatchery and wild parental fitness are highlighted in bold. For example, to compare the fitness of the 212 W and 161 H_{new} winter fish that went upstream in the 95-96 run year, we matched them against all *unclipped* winter fish that returned in 1998 to 2001 and whose scale ages indicated they were born in 1996 (i.e. from the 95-96 run year parents; see also Fig. 2) (We did not genotype the H fish from 98-01 for current analyses because they can't be the offspring of fish that bred in wild).

WINTER RUN

run year	wild fish passed	hatchery fish passed: H _{old} stock 13	hatchery fish passed: H _{new} stock 50
91-92	632	273	
92-	350	5	
93-	304	2	
94-	160	0	6
95-	212	0	161
96-	242	0	249
97-	184	0	162
98-	258	0	186
99-	875	0	222
00-	883	0	657
01-	954	0	682

SUMMER RUN

run year	wild fish passed	hatchery fish passed: H _{old} stock 24	hatchery fish passed: H _{new} stock 50
92-93	489	1722	
93-	243	1105	
94-	218	1635	
95-	132	520	
96-	182	1312	
97-	65	447	
98-	100	4	
99-	148	0	
00-	179	0	
01-	415	0	127
02- (03 to date)	540	0	492

Figure 2. Example of when offspring of each winter run parental breeding year are expected to return. Circles represent run years (e.g. 92-93 means the fish that returned in fall of 1992 through winter of 1993, and spawned in 1993. For simplicity, we will say spawning took place and their offspring were born in 92). Lines and numbers represent the percentage of babies born in a given breeding year that will return in each of the subsequent years. For example, 6% of the offspring born in the wild in 96 are expected to return to Powerdale Dam as unmarked adults in year 98, 61% are expected to return in year 99, and so on (the timing of return of hatchery fish is different from that of wild fish). Solid lines represent hatchery fish, dotted lines represent fish born in the wild. These numbers are based on age distributions of wild adults returning to Powerdale dam. Descendents of the first generation of winter run conservation hatchery stock 50 fish are illustrated as an example. Those hatchery fish spawned in nature mostly in years 95 and 96. Their F1 offspring are almost all returned by 01, and > 90% of F2's born in 99 and 00 are back by 05. The study proposed here involves genotyping samples collected through the 08-09 run year in order to insure a large sample of F2's from several F1 breeding years.

Specific questions asked

Adult offspring returning to the dam in the late 1990's and early 2000's were matched back their parents that were sampled in previous years (Fig. 2, Table 1; dates of birth from scale aging). From these data we asked the following questions.

(1) What is the mean and year-to-year variance in relative reproductive success (adult to adult production) of conservation hatchery-origin (H_{new}) and wild-origin (W) fish that spawned naturally in the Hood River?

The relative rate of adult-to-adult production by H_{new} and W fish is the key unknown parameter needed for predicting the demographic effects of hatchery supplementation on wild populations. Here we measured the parameter ($H_{new}:W$ fitness) for each of three years of winter fish (run years 95, 96, and 97; Table 1).

(2) Are “new” hatchery stocks closer in fitness to wild fish than “old” hatchery stocks?

Theory and substantial circumstantial evidence suggest that “old” hatchery stocks will have substantially lower total fitness than “new” hatchery stocks in the wild (Lynch and O’Hely, 2001; Fleming and Petersson, 2001). However, there has never been a direct test of this hypothesis, nor are there any empirical data on how much better the “new” stocks should perform. Here we compared the $H_{new}:W$ relative fitnesses with that of $H_{old}:W$. For winter run we have one run year of H_{old} vs. W (91), and for summer run we have data on two years (95 and 96) (Table 1).

Methods

(1) Sampling:

All handling of fish, phenotypic data collection, and sampling of scales and fin snips was done at the Powerdale Dam by ODFW staff. All fish approaching the dam were shunted into a trap and lifted into a building built specifically for the purpose of handling these fish. After being measured and sampled, each fish was either recycled downstream (e.g. extra hatchery fish), taken as broodstock or put above the dam to continue on to the spawning grounds. Sampling and database management protocols have been in place since project inception. Thus, we have an extensive database on the size, run timing, age

and freshwater residency (from scales), gender, fin clip and disposition (i.e. taken for broodstock, recycled, etc...) of every fish for which we also have pedigree data. We successfully genotyped between 97% and 99% of the steelhead collected for broodstock or passed over the dam each year. Therefore, we have an almost complete sample of all potential anadromous spawners from each run year.

(2) Molecular Methods:

We used a standard chelex protocol to extract DNA from fin snips or scales. Note that we obtain high quality DNA template from the scale samples, even those from the early 1990's. All extractions are done in 96-well plates.

From an initial set of 27 microsatellite loci that are known to work well in steelhead, we chose a set of eight loci based on the criteria that they amplify well, can be scored unambiguously in two sets of four multiplexed loci, and lack high-frequency null alleles. These eight loci provide a total power to reject a false parent-offspring pair via simple exclusion of > 0.9999 . Furthermore, our ground truthing experiments (see below) demonstrate a very low empirical rate of false parentage inclusion, and high power to exclude all but the true parents.

Template plates are pooled into 384-well plates for PCR, and those are pooled into four-locus, 384-well plates for multiplex scoring on an ABI 3100 16-channel capillary electrophoresis system. We use a Hydra-96 liquid handling robot (Robins Scientific) for all pipetting procedures involving plates (i.e. for all procedures following the initial handling of the scale or fin snip). Adding this device to our lab has cut sample handling errors down to virtually zero.

Table 2. Example of diversity at the eight loci we use (from a sample of 2,487 winter run fish. Summer fish have similar levels of diversity).

Locus	Number of Alleles	Expected Heterozygosity
Omy1001	28	0.91
Omy1011	27	0.90
Omy77	20	0.89
One108	33	0.92
One2	67	0.94
Rt191	35	0.93
Ssa407	30	0.90
Str2	44	0.95

(3) *Data analysis:*

Estimation of individual fitnesses

The eight-locus genotypes are merged with ODFW's database in Microsoft Access. For each returning fish (putative offspring) we search for it's mother and father in the year in which it was born (based on scale aging), plus or minus one year to allow for aging errors. From preliminary matching tests using a wider window we found only a few percent of fish are mis-aged by one year, and none are mis-aged by two years. For parent-offspring matching we use standard likelihood-based parentage analysis with an empirically-determined genotyping error rate (Marshall et al., 1998). Note that the presence of trout or precocious parr in the system is not be a problem, even if they obtain matings with anadromous parents. Our main question is to compare the average fitness of anadromous H and W parents, where fitness is defined as production of returning anadromous adults. Only offspring that assign to parents are relevant to the main question, and we have large sample sizes of those. Note that for run years 95 and 96 we should have approximately 99% and 96% of all offspring returned and genotyped. For 97 we should have approximately 66% in hand and genotyped (Fig. 2). Therefore, keep in mind that for 97 our comparison assumes there is no difference in year of return between offspring born of H_{new} and W parents.

Parameters vs. parameter estimates

For each run year we are interested in making inference about the fitness of the anadromous hatchery and wild fish that went above the dam. Because we sample all the fish at the dam each year, the fitness values we obtain for each type of parent (H or W) in a given run year are the parameters for that year, not estimates of the parameter. In other words, the parents that return each run year *are* the population of inference, not a sample from some larger population of hatchery and wild fish to which we wish to make inference for that year (the fitness of an individual fish may be measured with error owing to mis-assignment of offspring, but that is an issue of precision of measurement). On the other hand, the fitness estimates obtained for a given year can be considered to be a sample from some larger universe of run years. Because the ultimate goal here is to estimate H:W relative fitness for use in modeling conservation hatchery programs in general, the key values of interest are the mean and variance of H:W fitness among run years. So our main focus here is in measuring H:W fitness in as many run years as possible. In this report we provide data on H_{new} vs. W for three run years.

(4) Ground truthing:

All fish taken for broodstock are also genotyped. Therefore, as a form of ground truthing we ran fin-clipped returning adults from four brood years through the parentage analyses. For these analyses the fish taken for broodstock were included in the pool of wild potential parents. Ninety-six percent of the clipped returning fish were unambiguously matched back to a single mother-father pair in their expected brood year, and in every case our hatchery records show that that male-female pair was indeed crossed in the hatchery. The remaining unassigned, clipped offspring mismatch all potential parents at multiple loci and so are probably stray hatchery fish from out of the basin. Clipped and unclipped fish were treated identically during all stages of data collection. Therefore, we should have the same power to find the parents of unclipped returnees if their parents are in the parent pool.

Results

(1). Fraction of putative offspring that were assigned

We were able to unambiguously assign a mother-father pair to fewer than 40% of the putative offspring, a single parent to around 50% of them, and no parents to around 10-20%, depending on run year (Figure 3). From our ground-truthing experiments we know that the unassigned offspring are not all the result of experimental error. Because we sampled essentially all anadromous parents, these results suggest that missing parents may be non-anadromous, resident trout or precocious parr. Offspring for which neither parent could be found might also be wild strays or unclipped hatchery fish. Note, however, that for purposes of comparing the relative reproductive success of anadromous hatchery and wild fish, only those offspring that match to anadromous parents are relevant to this study. Again, the relevant measure of fitness here is number of returning anadromous adults produced per spawner. The sources of missing parents are a separate issue that we will address in the future.

(2). Analyses for individual fish of each sex:

Winter run: H_{old} vs. W and H_{new} vs. W

The 91 year gives a comparison of H_{old} vs. W, and the 95, 96 and 97 years give comparisons of H_{new} vs. W. In 91 the H_{old} had 35% the fitness of wild fish (Fig. 4; Table 3). In 95 the H_{new} had 85% the fitness of wild fish, in 96 they had 85-90%, and in 97 they had 90-108% the fitness of wild fish (Table 3). These results are consistent with the opinion that “new” hatchery stocks perform much better than “old” hatchery stocks. They also show that the relative performance of H_{new} fish might vary substantially from year to year. The caveat for the 97 run year is that although we have plenty of offspring back from that run year, they represent only 66% of those expected to have been produced by the 97 parents fish (see Fig. 2). Therefore, we are assuming that wild-born offspring of H_{new} parents do not return earlier than offspring of wild parents. Else the relative success of the H_{new} parents has been overestimated. We will verify the result when we genotype the 02-03 returnees.

Summer run H_{old} vs. W

We matched summer-run offspring back to their putative parents that spawned in the 95 and 96 run years. Both years involve comparisons of H_{old} vs. W. The relative fitness of H_{old} vs. W was 45-54% in 95 and only 17-30% in 96 (Table 3). These results again suggest that H_{old} have low fitness, and also show how variable the relative fitness of hatchery fish may be from year to year. In this case it is interesting that the relative performance of the H_{old} fish was lowest in the 96 run year when almost twice as many summer fish were on the spawning grounds. It may be that the relative success of hatchery fish is lower under more stressful conditions.

Figure 3. Fraction of offspring that were assigned to one parent, both parents or neither parent, by parental run year. For example, 33% of offspring born in 1992 (1991 run year) were matched to a single mother-father pair, 41% matched to a mother only, 11% to a father only, and 14% to neither parent. The offspring that match no parents could be unclipped hatchery fish or out-of-basin strays. The fact that some fish are missing single parents suggests that resident fish of some sort are breeding with anadromous fish.

Parentage Assignment Rate

Figure 4. Example of distributions of number of offspring per male or female breeder (1991 run year). Distributions for other years have similar shapes, although the right skew and variance increases with the mean (Table 3). Notice that most adults produced zero returning offspring. See Table 3 for summary data by parental gender and year.

Table 3. Analysis of average number of offspring produced per potential spawner of each type in each run year. n = number of adults collected at the dam (potential spawners) and genotyped. "Avg. # offspring per adult" = average fitness (number of offspring matched back) per potential spawner. "H/W relative fitness" is the average # of offspring per hatchery adult divided by the avg. # per wild adult. Discrepancies between the total number of fish in this table and in Table 1 in some years result because (1) for this analysis we excluded hatchery fish that had been recycled once or twice before being put above the dam (because that may have affected their fitness), and (2) we excluded a few parents for whom we did not have complete genotypes.

WINTER RUN

Run Year	H _{old} (stock 13)			Wild			
1991-92	n	Avg. # offspring per adult (variance)		n	Avg. # offspring per adult (variance)		H/W relative fitness
males	165	0.24 (0.28)		247	0.68 (1.10)		0.35
females	99	0.23 (0.36)		379	0.68 (1.02)		0.34
Run Year	H _{new} (stock 50)			Wild			
1995-96							
males	90	3.66 (19.6)		78	4.31 (25.05)		0.85
females	65	4.39 (16.4)		132	5.05 (25.23)		0.87
Run Year	H _{new} (stock 50)			Wild			
1996-97							
males	95	2.28 (6.01)		93	2.54 (8.03)		0.90
females	153	2.08 (3.71)		148	2.47 (7.04)		0.85

WINTER RUN continued

1997-98							
males	53	2.08 (3.19)		64	2.30 (5.26)		0.90
females	107	2.71 (6.89)		109	2.50 (6.05)		1.08

SUMMER RUN

Run Year	H _{old} (stock 24)			Wild			
1995-96							
males	211	0.72 (1.07)		44	1.34 (3.44)		0.54
females	297	0.45 (0.67)		86	1.01 (3.75)		0.45
Run Year	H _{old} (stock 24)			Wild			
1996-97							
males	474	0.45 (0.69)		61	1.48 (4.45)		0.30
females	766	0.20 (0.32)		121	1.18 (3.70)		0.17

(3) *Analyses of parental pairs (performance of HxH, HxW and WxW crosses):*

We could estimate the proportions of each type of observed cross expected if H and W fish mate randomly, and then compare those proportions to the observed proportions of parental pairs we actually detected of each type. But because we cannot count pairs that left no surviving offspring, there is no way to disentangle non-random mating from differences in parental fecundity or offspring survival (you would need to observe matings to do that). If we restrict our analysis to pairs that left *at least one surviving offspring*, then we can calculate the relative fitness of each type of cross for that truncated dataset. Any difference here is necessarily owing to offspring survival or parental fecundity because we have restricted the inference to those fish that, by definition, mated. This analysis probably underestimates the fitness differences among the three types of pairs because we have no zero-offspring class. Nevertheless, even for this restricted dataset our results show that $H_{old} \times H_{old}$ crosses always did worse than $H_{old} \times W$, which in turn were worse than $W \times W$ crosses (see Table 4, WINTER RUN 91 and SUMMER RUN 95-96). This result indicates that breeding by H_{old} fish in the wild may indeed be dragging down the fitness of wild fish. In contrast, $H_{new} \times H_{new}$ and $H_{new} \times W$ crosses did quite well, equalling or exceeding $W \times W$ fitness in some years, although the differences are quite variable from year to year (see Table 4, WINTER RUN 95-97). Of more importance, however, these results show that all three types of crosses ($H_{new} \times H_{new}$, $H_{new} \times W$, and $W \times W$) occur in the wild and produce substantial numbers of surviving F1 offspring.

Table 4. Analysis of average number of offspring per type of parental pair (HxH, HxW or WxW) for pairs *that left at least one offspring*. n = number of pairs of that type unambiguously identified as leaving at least one offspring. The two reciprocal H x W crosses were pooled to increase the sample sizes.

WINTER RUN

Type of cross	1991 Avg. # (n)	Fitness of the cross relative to W x W	Type of cross	1995 Avg. # (n)	Fitness of the cross relative to W x W	1996 Avg. # (n)	Fitness of the cross relative to W x W	1997 Avg. # (n)	Fitness of the cross relative to W x W
W x W	1.33 (87)	1.00	W x W	2.53 (72)	1.00	1.54 (57)	1.00	1.24 (34)	1.00
W x H _{old}	1.09 (11)	0.82	W x H _{new}	2.05 (78)	0.81	1.58 (90)	1.03	1.50 (47)	1.21
H _{old} x H _{old}	NA (0)	NA	H _{new} x H _{new}	1.29 (21)	0.51	1.42 (38)	0.92	1.56 (25)	1.26

SUMMER RUN

	Type of cross		1995 Avg. # (n)	Fitness of the cross relative to W x W		1996 Avg. # (n)	Fitness of the cross relative to W x W
	W x W		1.75 (4)	1.00		1.46 (13)	1.00
	W x H _{old}		1.29 (17)	0.74		1.24 (50)	0.85
	H _{old} x H _{old}		1.0 (10)	0.57		1.05 (63)	0.72

CONCLUSIONS

1. Fish from old hatchery stock consistently have very low fitness (usually less than 50% that of wild fish) when breeding in the wild. The fact that $H_{old} \times W$ crosses consistently produce fewer offspring than $W \times W$ crosses (Table 4) suggests that having H_{old} breeders in a system might lower the fitness of the wild population. Whether the surviving wild-born offspring of such crosses re-establish “wild” levels of fitness after one full generation of selection in nature remains to be tested.

2. Fish from new, conservation hatchery stock have fitness that is about equal to that of wild fish (less than wild in two years, greater than in the third year). The same pattern is apparent whether one examines the relative fitness of individual parents or that of pairs that left at least one offspring. The similar fitnesses $H_{new} \times W$ and $W \times W$ pairs, suggests that having H_{new} fish in the system is probably not obviously dragging down the fitness of the wild population for genetic reasons (as might have been expected under some models; e.g. Lynch and O’Hely, 2001). Thus, the conservation hatchery program appears to have added a demographic boost to the population without having obvious negative genetic consequences - at least in regards the effects of domestication selection and mutation accumulation that should occur in the hatchery. We have not yet conducted a formal analysis of the effect of the hatchery program on the effective size of the wild population (e.g. Ryman et al., 1995), but the high levels of microsatellite diversity we still observe in both runs suggest that reduced effective size is not a problem.

3. The surprisingly large number of missing parents, and the fact that most missing parents are fathers (Fig. 3), suggests that precocious parr or resident trout are obtaining matings that produce anadromous offspring. Alternate explanations for offspring that lack both parents include a large number of unclipped hatchery fish or wild strays entering the system.

Future Work

We hope to continue genotyping fish through the rest of this decade. Additional questions we plan to address include:

(1) Do F1 progeny (born in the wild) of $H_{new} \times W$, $H_{new} \times H_{new}$ and $W \times W$ winter run parents differ in their production of F2 progeny?

We know from our current analyses that all three types of matings occur on the spawning ground, and that all three types of mating produce offspring that return to spawn as adults. F2 offspring of those winter F1s that spawned in the late 1990s are now returning (see Fig. 2). If we continue sampling through the end of the decade we will have a large number of returned F2s from multiple brood years with which to test the relative fitness of different types of F1s (Fig. 2). Given the apparently high fitness of H_{new} hatchery fish, our expectation is that the three types of wild-born F2's will have similar fitnesses.

(2) Selection to maintain the difference between summer and winter runs:

What is the rate of hybridization between the runs? What are the phenotypes (run time, size, freshwater residency) and actual fitnesses of any hybrids?

(3) Selection on measurable phenotypic traits:

We can use standard selection gradient analysis (Lande and Arnold, 1983) to analyze fitness as a function of body size, run time, age and freshwater residency (known from scales), after controlling for hatchery/wild genetic background.

(4) Quantitative genetic parameter estimation:

From our pedigrees we can estimate the heritabilities of, and genetic correlations among any measurable phenotypic traits. We can also estimate the average breeding value for each trait in individuals of $H \times H$ and $W \times W$ genetic background, in order to test whether genetic changes in the hatchery, and subsequent mating with wild fish, could be changing phenotypic distributions in the wild population (Ford, 2001).

(5) Parental contributions of resident, non-anadromous fish

We sample all potential breeding adults passed over the dam, and we know from our ground truthing experiments the expected rate of mismatching owing to experimental error. Therefore, unassigned offspring are either wild strays from out of the basin, or were parented by resident fish (non-anadromous *O. mykiss*, or precocious parr). We will use likelihood methods (Rannala and Mountain, 1997) to attempt to determine the most likely source of missing parents (of offspring that only match to a single known parent), and whether fish lacking both parents are most likely to be Hood River wild, Hood River hatchery (unclipped) or immigrants from adjacent steelhead populations. Because we sample all anadromous parents, the Hood River is an ideal system in which to ask questions about the rate of parentage from resident fish and about the sources of those fish.

(6) Effective size estimation

From the pedigrees we can obtain direct estimates of the effective size (N_e) of each population over time. These data will be used to estimate the impact of hatchery programs on the effective size of the wild population and to provide basic parameter estimates such as the variance in family sizes (number of returning adults) for hatchery broodstock, for H fish in the wild, and for W fish in the wild. These are important parameters that are unknown for most populations and can be very useful for estimating N_e and the effects of supplementation in other steelhead populations (e.g. *sensu* Ryman et al., 1995). We can also use our system to evaluate the accuracy of indirect methods for estimating effective size (e.g. Waples, 2002; Anderson et al., 2000). If the indirect methods give very different values from the pedigree-based estimates, then we can ask what assumptions of the indirect methods cause the difference. Note that because of our ability to sample all potential anadromous parents, we can take into account the contributions of non-anadromous, resident fish in our calculations.

References Cited

- Anderson, E.C et al. 2000. Monte Carlo evaluation of the likelihood for N_e from temporally-spaced samples. *Genetics* 156: 2109-2118.
- Fleming, I.A. and E. Petersson. 2001. The ability of released hatchery salmonids to breed and contribute to the natural productivity of wild populations. *Nordic Journal of Freshwater Research* 75:71-98.
- Ford, M.J. 2001. Selection in captivity during supportive breeding may reduce fitness in the wild. *Conservation Biology* 16:815-825.
- Lande, R. and S.J. Arnold. 1983. The measurement of selection on correlated characters. *Evolution* 36:1210-1226.
- Lynch, M and M. O'Hely. 2001. Supplementation and the genetic fitness of natural populations. *Conservation Genetics* 2:363-378
- Marshall et al. 1998. Statistical confidence for likelihood-based paternity inference in natural populations. *Molecular Ecology* 7:639-655.
- Rannala, B. and J. L. Mountain. 1997. Detecting immigration by using multilocus genotypes. *Proc. Natl. Acad. Sci. USA* 94: 9197-9221
- Ryman, N. et al. 1995. Supportive breeding and variance effective population size. *Conservation Biology*. 9: 1619-1628.
- Waples, R. 2002. Effective size of fluctuating salmon populations. *Genetics* 161:783-791.