**SNAKE RIVER SOCKEYE SALMON
CAPTIVE BROODSTOCK PROGRAM
RESEARCH ELEMENT**

**ANNUAL PROGRESS REPORT
January 1, 2007—December 31, 2007**

Photo by Jeff Heindel

Prepared by:

**Mike Peterson, Fisheries Research Biologist
Kurtis Plaster, Senior Fisheries Technician
Laura Redfield, Fisheries Technician
Jeff Heindel, Conservation Hatchery Supervisor
and
Paul Kline, Principal Fisheries Research Biologist**

**IDFG Report Number 08-10
June 2008**

Snake River Sockeye Salmon Captive Broodstock Program Research Element

2007 Annual Project Progress Report

Part 1—Project Overview

**Part 2—*Oncorhynchus nerka* Population Monitoring and Redfish
Lake Sport Fishery Investigations**

**Part 3—Sockeye Salmon Smolt Monitoring and Evaluation And
Sockeye Salmon Spawning Investigations and Unmarked
Juvenile Out-migrant Monitoring**

Part 4—Predator Surveys

By

**Mike Peterson
Kurtis Plaster
Laura Redfield
Jeff Heindel
Paul Kline**

**Idaho Department of Fish and Game
600 South Walnut Street
P.O. Box 25
Boise, Idaho 83707**

To:

**U.S. Department of Energy
Bonneville Power Administration
Division of Fish and Wildlife
P.O. Box 3621
Portland, OR 97283-3621**

**Project Number 1991-07-200
Contract Number 37981**

**IDFG Report Number 08-10
June 2008**

TABLE OF CONTENTS

	<u>Page</u>
EXECUTIVE SUMMARY	1
PART 1—PROJECT OVERVIEW.....	3
BACKGROUND	3
SNAKE RIVER SOCKEYE SALMON CAPTIVE BROODSTOCK PROGRAM.....	4
PROJECT GOALS.....	5
PROJECT OBJECTIVES.....	5
STUDY AREA	6
2006 and 2007 Captive Broodstock Program Egg and Juvenile Supplementation.....	7
LITERATURE CITED	12
PART 2—ONCORHYNCHUS NERKA POPULATION MONITORING AND REDFISH LAKE SPORT FISHERY INVESTIGATIONS.....	17
INTRODUCTION	17
METHODS	17
<i>Oncorhynchus Nerka</i> Population Monitoring.....	17
Redfish Lake Sport Fishery Investigations.....	18
RESULTS AND DISCUSSION	18
<i>Oncorhynchus Nerka</i> Population Monitoring.....	18
Redfish Lake	18
Alturas Lake	19
Pettit Lake	19
Redfish Lake Sport Fishery Investigations.....	20
LITERATURE CITED	26
PART 3—SOCKEYE SALMON JUVENILE OUT-MIGRANT AND ADULT SPAWNING MONITORING AND EVALUATION	27
INTRODUCTION	27
SOCKEYE SALMON JUVENILE OUT-MIGRANT MONITORING AND EVALUATION	27
ADULT SPAWNING MONITORING AND EVALUATION.....	27
METHODS	28
SOCKEYE SALMON JUVENILE OUT-MIGRANT MONITORING AND EVALUATION	28
Redfish Lake Creek Trap	28
Alturas Lake Creek Trap	29
Pettit Lake Creek Trap	29
Salmon River and Redfish Lake Creek Smolt Groups	29
Mainstem Snake and Columbia River Dams	29
ADULT SPAWNING MONITORING AND EVALUATION.....	30
Sockeye Salmon Spawning Investigations	30
Unmarked Juvenile Out-migrant Monitoring.....	30
Adult Trapping on Redfish Lake Creek	30
RESULTS AND DISCUSSION	31
SOCKEYE SALMON JUVENILE OUT-MIGRANT MONITORING AND EVALUATION	31
Redfish Lake Creek Trap	31

Table of Contents, continued.

	<u>Page</u>
Alturas Lake Creek Trap	32
Pettit Lake Creek Trap	32
Salmon River and Redfish Lake Creek Smolt Groups	32
Mainstem Snake and Columbia River Dams	33
SOCKEYE SALMON SPAWNING INVESTIGATIONS AND UNMARKED JUVENILE	
OUT-MIGRANT MONITORING	33
Sockeye Salmon Spawning Investigations	33
Unmarked Juvenile Out-migrant Monitoring.....	34
LITERATURE CITED	43
PART 4—PREDATOR SURVEYS.....	45
INTRODUCTION	45
METHODS	45
Adult Bull Trout Trapping Redfish Lake Creek.....	45
RESULTS AND DISCUSSION	46
Fishhook Creek.....	46
Alpine Creek	46
Bull Trout Capture, Mark, and Haul Operations Redfish Lake Creek	46
LITERATURE CITED.....	51
ACKNOWLEDGEMENTS	52
APPENDICES.....	53

LIST OF TABLES

	<u>Page</u>
Table 1. Eyed-egg production by IDFG and NOAA facilities for the Snake River sockeye salmon captive broodstock program 1991—2007.....	8
Table 2. Snake River sockeye salmon captive broodstock program egg and fish reintroduction history.....	8
Table 3. Physical and morphometric characteristics of three study lakes located in the Sawtooth Valley, Idaho.....	9
Table 4. Sockeye salmon releases to Sawtooth Valley waters in 2006.	9
Table 5. Sockeye salmon releases to Sawtooth Valley waters in 2007.	10
Table 6. Estimated <i>O. nerka</i> population, density, and biomass for Redfish, Alturas, and Pettit lakes, 1990 to 2007.....	22
Table 7. Estimated 2007 <i>O. nerka</i> abundance, density (fish/ha), and biomass (kg/ha) by age class in Redfish, Alturas, and Pettit lakes.....	23
Table 8. Estimated kokanee escapement to Fishhook Creek 1991 to 2007 and Alturas Lake Creek 1992 to 2007. Data obtained from the Shoshone-Bannock Tribes.	23
Table 9. Estimated angler effort on Redfish Lake for the 2007 fishing season.	24
Table 10. Historical kokanee catch rates, kokanee harvest estimates, bull trout catch rates, and angler effort for the Redfish Lake fishery.	24
Table 11. Catch rates (fish/hour) for summer 2007 on Redfish Lake categorized by day type and species.	24
Table 12. Estimated number of fish harvested and released on Redfish Lake during the summer 2007.	24
Table 13. Out-migration estimate for natural origin sockeye salmon smolts captured at the Redfish Lake Creek trap from April 14 to June 13, 2007.	35
Table 14. Summary of 2007 sockeye salmon smolt out-migration information (by release strategy) at trap locations and at Lower Granite Dam (LGR). Sawtooth Fish Hatchery (SFH) was the rearing location for the fall-direct released (FDR) presmolts and Salmon River smolt release group. Oxbow Fish Hatchery (OFH) was the rearing location for the Redfish Lake Creek smolt release.....	36
Table 15. Estimated overwinter survival until out-migration for Sawtooth Fish Hatchery-reared presmolts released in the fall to Redfish, Alturas, and Pettit lakes.	37
Table 16. Redfish Lake Sockeye Salmon Captive Broodstock Program prespawn adult release history.....	37
Table 17. Summary of sockeye salmon redd observations in Redfish Lake in 2007.	38
Table 18. Bull trout adult fish counts and redd counts in index sections of Fishhook Creek from 1998 to 2007.....	48
Table 19. Bull trout adult fish counts and redd counts in index sections of Alpine Creek from 1998 to 2007.	48

LIST OF FIGURES

	<u>Page</u>
Figure 1. Map of the upper Salmon River watershed located in the Sawtooth Valley, Idaho.	11
Figure 2. Estimated bull trout catch rates from 1996-2007 within Redfish Lake. The trend line is used to show the population increase within the lake. The equation for the trend line is $y = 0.0117x + 0.103$ with an R^2 of 0.17.	25
Figure 3. Daily capture of natural origin and hatchery origin sockeye salmon smolts (unexpanded) at the Redfish Lake Creek trap during the 2007 out-migration.	39
Figure 4. Length frequency of natural origin sockeye salmon smolts ($n = 491$) collected at Redfish Lake Creek trap in 2007.	40
Figure 5. Length frequency of fall-direct released sockeye salmon smolts ($n = 609$) collected at Redfish Lake Creek trap in 2007.	40
Figure 6. Spawning locations for sockeye salmon in Redfish Lake: 1) Sockeye beach, 2) the south beach near the slide, 3) the area near the U.S. Forest Service transfer camp dock, and 4) West side spawning area used in 2007.	41
Figure 7. Natural origin sockeye salmon smolt out-migration estimated at Redfish Lake Creek, Alturas Lake Creek, and Pettit Lake Creek traps from 1991 to 2007 (juvenile out-migrant traps on Pettit Lake Creek were not operated every year).	42
Figure 8. Location bull trout redd index sections in Fishhook Creek in 2007.	49
Figure 9. Location of bull trout redd index sections in Alpine Creek in 2007.	50

LIST OF APPENDICES

Appendix A. Fork length, weight (g) and age of <i>O. nerka</i> captured during midwater trawls conducted during September 2007 on Redfish, Pettit, and Alturas lakes.	54
Appendix B. Arrival dates at Lower Granite Dam for PIT-tagged sockeye salmon smolts during the 2007 migration year.	59

EXECUTIVE SUMMARY

On November 20, 1991, the National Oceanic Atmospheric Administration listed Snake River sockeye salmon *Oncorhynchus nerka* as endangered under the Endangered Species Act of 1973. In 1991, the Shoshone-Bannock Tribes (SBT) and Idaho Department of Fish and Game (IDFG) initiated the Snake River Sockeye Salmon Captive Broodstock Program to conserve and rebuild populations in Idaho. Restoration efforts are focused on Redfish, Pettit, and Alturas lakes within the Sawtooth Valley. The first release of hatchery-produced adults occurred in 1993. The first release of juvenile sockeye salmon from the captive broodstock program occurred in 1994. In 1999, the first anadromous adult returns from the captive broodstock program were recorded when six jacks and one jill were captured at the IDFG Sawtooth Fish Hatchery.

In 2007, progeny from the captive broodstock program were released using four strategies: 1) eyed-eggs were planted in Pettit Lake in November; 2) age-0 presmolts were released to Alturas, Pettit, and Redfish lakes in October; 3) age-1 smolts were released into Redfish Lake Creek and the upper Salmon River in May; and 4) hatchery-produced adult sockeye salmon were released to Redfish Lake for volitional spawning in September.

Oncorhynchus nerka population monitoring was conducted on Redfish, Alturas, and Pettit lakes using a midwater trawl in September 2007. Population abundances were estimated at 73,702 fish for Redfish Lake, 124,073 fish for Alturas Lake, and 14,746 fish for Pettit Lake.

Angler surveys were conducted from May 26 through August 7, 2007 on Redfish Lake to estimate kokanee harvest. On Redfish Lake, we interviewed 102 anglers and estimated that 56 kokanee were harvested. The calculated kokanee catch rate was 0.03 fish/hour for each kokanee kept.

The juvenile out-migrant trap on Redfish Lake Creek was operated from April 14 to June 13, 2007. We estimated that 5,280 natural origin and 14,256 hatchery origin sockeye salmon smolts out-migrated from Redfish Lake in 2007. The hatchery origin component originated from a 2006 fall presmolt direct-release. The juvenile out-migrant traps on Alturas Lake Creek and Pettit Lake Creek were operated by the SBT from April 19 to May 23, 2007 and April 18 to May 29, 2007, respectively. The SBT estimated 1,749 natural origin and 4,695 hatchery origin sockeye salmon smolts out-migrated from Pettit Lake and estimated 8,994 natural origin and 6,897 hatchery origin sockeye salmon smolts out-migrated from Alturas Lake in 2007. The hatchery origin component of sockeye salmon out-migrants originated from fall presmolt direct-releases made to Pettit and Alturas lakes in 2006.

In 2007, the Stanley Basin Sockeye Technical Oversight Committee (SBSTOC) chose to have all Snake River sockeye salmon juveniles (tagged and untagged) transported due to potential enhanced survival. Therefore, mainstem survival evaluations were only conducted to Lower Granite Dam. Unique PIT tag interrogations from Sawtooth Valley juvenile out-migrant traps to Lower Granite Dam were utilized to estimate survival rates for out-migrating sockeye salmon smolts. Survival rate comparisons were made between smolts originating from Redfish, Alturas, and Pettit lakes and the various release strategies. Alturas Lake hatchery origin smolts tagged at the out-migrant trap recorded the highest survival rate of 78.0%.

In 2007, 494 hatchery origin adult sockeye salmon were released to Redfish Lake for natural spawning. We observed 195 areas of excavation in the lake from spawning events. This

was the highest number of redds observed in Redfish Lake since the program was initiated. Suspected redds were approximately 3 m x 3 m in size and were constructed by multiple pairs of adults.

To monitor the predator population found within the lakes, we monitored bull trout spawning in Fishhook Creek, a tributary to Redfish Lake; and in Alpine Creek, a tributary to Alturas Lake. This represented the tenth consecutive year that the index reaches have been surveyed on these two streams. Adult counts (41 adults) and redd counts (22 redds) in Fishhook Creek increased compared to counts conducted since monitoring began in 1998. Beginning in 2007, we also surveyed an additional trend site in Fishhook Creek resulting in observing 43 adult bull trout and 30 additional redds. Bull trout numbers (13 adults) and the number of redds observed (18 redds) have gradually increased in Alpine Creek compared to counts from initial monitoring.

Authors:

Mike Peterson
Fisheries Research Biologist

Kurtis Plaster
Senior Fisheries Technician

Laura Redfield
Fisheries Technician

Jeff Heindel
Conservation Hatchery Supervisor

Paul Kline
Principal Fisheries Research Biologist

PART 1—PROJECT OVERVIEW

BACKGROUND

The Idaho Department of Fish and Game (IDFG) initiated the Snake River sockeye salmon *Oncorhynchus nerka* captive broodstock program in May 1991 in response to the decline of anadromous returns to the Sawtooth Valley in central Idaho. Waples et al. (1991) described Snake River sockeye salmon as a species on the threshold of extinction, and it was listed as endangered under the Endangered Species Act (ESA) on November 20, 1991 (ESA; 16 U.S.C.A. §§1531 to 1544). Snake River sockeye salmon are one of 27 stocks of Pacific salmon and steelhead (*Genus: Oncorhynchus*) in the Columbia River basin currently listed as threatened or endangered under the Endangered Species Act (www.nwr.noaa.gov/1salmon/salmesa/index).

Historically, Redfish, Alturas, Pettit, Stanley, and Yellowbelly lakes supported sockeye salmon in the Sawtooth Valley (Chapman et al. 1990; Evermann 1895; Bjornn et al. 1968) (Figure 1). Historical observations and discussions with local residents by Evermann (1895; 1896) described the Sawtooth Valley lakes as being important spawning and rearing areas for sockeye salmon; actual adult escapement enumeration or estimations were not conducted. Adult sockeye salmon escapement to Redfish Lake was enumerated from 1954 through 1966 by the IDFG, University of Idaho, and the United States Bureau of Commercial Fisheries. During this time, adult escapement ranged from a high of 4,361 in 1955 to a low of 11 in 1961 (Bjornn et al. 1968). Adult escapement enumeration was reinitiated in 1985 by the IDFG. Between 1985 and 1990, 62 adults were estimated to have returned to the Sawtooth Valley. No redds or anadromous adults were identified in Redfish Lake in 1990. Hydropower development, water withdrawal and diversions, water storage, harvest, predation, and inadequate regulatory mechanisms were outlined as factors contributing to the Snake River sockeye salmon's decline (Federal Register 1991).

The National Marine Fisheries Service (NMFS) listed Snake River sockeye salmon as an endangered species under the 1973 Endangered Species Act (as amended in 1978) in November of 1991 (Federal Register 1991). A population is considered a distinct population segment and, hence, a species for purposes of the Endangered Species Act if it represents an evolutionary significant unit of the biological species (Waples 1991). To be considered an evolutionary significant unit, a stock must satisfy two criteria: 1) it must be reproductively isolated from other conspecific population units; and 2) it must represent an important component in the evolutionary legacy of the biological species (Waples 1991). At the time of listing, the Redfish Lake sockeye salmon population was the only remaining sockeye salmon population of the Snake River sockeye salmon stock. Snake River sockeye salmon are one of three remaining stocks of sockeye salmon in the Columbia River system; the other two stocks, Okanogan Lake sockeye salmon and Wenatchee Lake sockeye salmon, are located in tributaries of the upper Columbia River. Approximately 1,127 river kilometers separate Snake River sockeye salmon from the nearest sockeye salmon populations in the upper Columbia River. Additionally, there has been no reported evidence of straying of sockeye salmon from the upper Columbia River into Redfish Lake (Waples et al. 1991; Winans et al. 1996). Mitochondrial DNA analyses completed in 2003 confirmed the genetic isolation of the upper Columbia River stocks from the Snake River sockeye salmon stock (Faler and Powell 2003).

Sockeye salmon returning to Redfish Lake travel a greater distance from the Pacific Ocean (1,448 river kilometers) and to a higher elevation (2,138 meters) than any other sockeye salmon population in the world. Additionally, Redfish Lake supports the species' southernmost

population within its recognized range (Burgner 1991). Together these characteristics presented a strong argument for the ecological uniqueness of the Snake River habitat and for the unique adaptive genetic characteristics of the Snake River sockeye salmon stock (Waples et al. 1991).

Genetic investigations conducted during and after Snake River sockeye salmon were listed further refined genetic relationships between anadromous sockeye salmon, residual sockeye salmon, and resident kokanee present in Redfish Lake. The presence of all three of these life history strategies of *O. nerka* in Redfish Lake complicated the ESA listing. Anadromous *O. nerka* (sockeye salmon) spawn on the shoals of the lake in October and November. Juveniles out-migrate during the spring from their nursery lake at age-1 or age-2 and remain in the ocean for two to three years before returning to their natal area to spawn. Residual *O. nerka* (sockeye salmon) spawn with anadromous sockeye salmon on the shoals of the lake in October and November. Residual sockeye salmon spend their entire life in their nursery lake. Variable proportions of anadromous and residual progeny may conform to a residual life history pattern. Resident *O. nerka* (kokanee) also complete their lifecycle in freshwater. They remain in Redfish Lake until maturation and spawn in Fishhook Creek, a tributary creek to Redfish Lake, in August and September. Kokanee are nonindigenous to Redfish Lake and were periodically stocked from a range of hatchery sources beginning in 1930 and continuing through 1972 (Bowler 1990). Redfish Lake anadromous sockeye salmon, residual sockeye salmon, and out-migrants were determined to be genetically similar, whereas kokanee were found to be genetically different (Brannon et al. 1992, 1994; Cummings et al. 1997; Waples et al. 1997). Because of their genetic similarity, residual sockeye salmon were added to the ESU listing in 1992.

SNAKE RIVER SOCKEYE SALMON CAPTIVE BROODSTOCK PROGRAM

The Snake River sockeye salmon captive broodstock program collected fish from the following sources for broodstock and reintroduction purposes: 1) anadromous adult returns that were trapped between 1990 and 1998 and retained for hatchery spawning, 2) residual adults that were trapped between 1992 and 1995 and retained for hatchery spawning, and 3) smolts that were trapped between 1991 and 1993, reared until maturity, and spawned in the hatchery. Fifth generation lineages of the founders are currently in captive broodstock culture. Both IDFG and National Oceanic and Atmospheric Administration (NOAA) Fisheries maintain Snake River sockeye salmon captive broodstocks. Groups of fish are reared at two facilities to avoid the potential catastrophic loss of the unique genetics of the stock. Idaho Department of Fish and Game rears annual captive broodstocks from the egg stage to maturity at Eagle Fish Hatchery in Eagle, Idaho (Johnson 1993; Johnson and Pravecek 1995, 1996; Pravecek and Johnson 1997; Pravecek and Kline 1998; Kline and Heindel 1999; Kline et al. 2003a, 2003b; Kline and Willard 2001; Willard et al. 2003a; Baker et al. 2005a, 2005b). Additionally, NOAA Fisheries rears annual captive broodstocks from the egg stage to maturity at the Manchester Research Station and Burley Creek Hatchery near Seattle, Washington (Flagg 1993; Flagg and McAuley 1994; Flagg et al. 1996, 2001; Frost et al. 2002).

The IDFG and NOAA Fisheries captive broodstock programs have produced approximately 3,846,973 eyed-eggs from 1991 through 2007 (Table 1). Each year approximately 800 eggs are selected to be reared in the hatchery as an annual captive broodstock; the remaining eggs are utilized for reintroduction into the habitat.

The development of captive broodstock program reintroduction plans follows a “spread-the-risk” philosophy incorporating multiple release strategies and multiple lakes (Hebdon et al. 2004). Progeny from the captive broodstock program are reintroduced to Sawtooth Valley waters at different life stages using a variety of release options including: 1) eyed-egg plants to in-lake incubator boxes in November and December, 2) presmolt releases direct to lakes in October, 3) smolt releases to outlet streams in May, and 4) prespawn adult releases (hatchery-reared) direct to lakes in September. All hatchery-reared presmolt and smolt release groups are uniquely marked to identify release strategy and release origin. To date, approximately 2,624,154 sockeye salmon eggs and fish have been reintroduced to Sawtooth Valley waters (Table 2). Overall survival to release (over different strategies) from the eyed-egg stage of development has averaged just under 68% for the program.

Life history traits exhibited by wild sockeye salmon and habitat carrying capacity evaluations conducted by the Shoshone-Bannock Tribes (SBT) (Teuscher and Taki 1995, 1996; Taki and Mikkelsen 1997; Taki et al. 1999; Griswold et al. 2000; Lewis et al. 2000; Kohler et al. 2001, 2002) influence annual release plans. Bjornn et al. (1968) conducted smolt out-migration and adult return monitoring between 1954 and 1966; their work provides a detailed account of sockeye salmon life history in Redfish Lake. Smolt out-migration from Redfish Lake begins in early April, peaks in mid-May, and is complete by mid-June. Smolts out-migrate at either age-1 or age-2; the proportion of age-1 and age-2 smolts varies every year. During their 11-year study, Bjornn et al. (1968) noted a dominance of age-1 smolts for six of the 11 years. Age-2 smolts ranged from 2% to 77% of the total out-migration over the course of the early monitoring effort. Age-2 smolts are common in many other sockeye salmon lakes. The reasons for the additional freshwater residence time are unclear (Burgner 1991). Smolt fork length ranges between 45 mm and 120 mm. Adult sockeye salmon begin arriving at Redfish Lake Creek in mid-July and continue escapement through early September. Sockeye salmon spawn over submerged beach substrate of the lake; spawning peaks in mid-October. Returning adults are primarily two-ocean fish (Bjornn et al. 1968).

Efforts to prevent extinction of the Snake River sockeye salmon are coordinated through the Stanley Basin Sockeye Technical Oversight Committee (SBSTOC), a team of biologists representing IDFG, the SBT, NOAA Fisheries, and the University of Idaho. The Bonneville Power Administration provides coordination for the SBSTOC process.

PROJECT GOALS

The immediate goal of the program is to utilize captive broodstock technology to conserve the population's unique genetics. Long-term goals include increasing the number of individuals in the population to address NOAA's interim abundance guidelines and to provide sport and treaty harvest opportunity. Draft ESA delisting criteria for Snake River sockeye salmon includes the return of 1,000 adults to Redfish Lake, 500 adults to Pettit Lake, and 500 adults to Alturas Lake for two generations (NMFS 2002). Interim abundance targets must be met without relying on hatchery production (e.g., natural origin adults).

PROJECT OBJECTIVES

1. Develop captive broodstocks from Redfish Lake sockeye salmon, culture broodstocks, and produce progeny for reintroduction.

2. Determine the contribution hatchery-produced sockeye salmon make toward avoiding population extinction and increasing population abundance.
3. Describe *O. nerka* population characteristics for Sawtooth Valley lakes in relation to carrying capacity and broodstock program reintroduction efforts.
4. Utilize genetic analysis to discern the origin of wild and broodstock sockeye salmon to provide maximum effectiveness in their utilization within the broodstock program.
5. Transfer technology through participation in the technical oversight committee process, provide written activity reports, and participate in essential program management and planning activities.

Idaho Department of Fish and Game's participation in the Snake River Sockeye Salmon Captive Broodstock Program includes two areas of effort: 1) sockeye salmon captive broodstock culture, and 2) sockeye salmon research and evaluations. Although objectives and tasks from both components overlap and contribute to achieving the same goals, work directly related to sockeye salmon captive broodstock culture appears under a separate cover (Baker *in review*). Research and evaluation activities associated with Snake River sockeye salmon are permitted under the ESA (NOAA) Nos. 1120, 1124, and 1481 (for a review see Kline 1994; Kline and Younk 1995; Kline and Lamansky 1997; Hebdon et al. 2000; Hebdon et al. 2002; Hebdon et al. 2003; Willard et al. 2003b; Willard et al. 2005). This report details fisheries research information collected between January 1 and December 31, 2007. Research information includes *O. nerka* population monitoring in Sawtooth Valley lakes, sport fishery evaluation on Redfish Lake, smolt out-migration monitoring and evaluation at lake outlets, telemetry studies of mature adult sockeye salmon released to Sawtooth Valley lakes for natural spawning, and predator investigations in tributaries to Redfish and Alturas lakes.

STUDY AREA

The program's recovery efforts focus on Redfish, Pettit, and Alturas lakes in the Sawtooth Valley located within the Sawtooth National Recreation Area (Figure 1). These lakes provide critical spawning and rearing habitat under the ESA listing. Lakes in the Sawtooth Valley are glacial-carved and considered oligotrophic. The three lakes range in elevation from 1,996 m (Redfish Lake) to 2,138 m (Alturas Lake) and are located 1,448 km (Redfish Lake) to 1,469 km (Alturas Lake) from the Pacific Ocean. Redfish Lake is the largest of the three lakes (615 ha), Pettit Lake is the smallest (160 ha), and Alturas Lake (338 ha) is intermediate in surface area (Table 3). Reintroduction efforts have been ongoing in Redfish Lake since 1993, Pettit Lake since 1995, and Alturas Lake since 1997.

In addition to *O. nerka*, numerous native and nonnative fish reside in the study lakes and streams within the Sawtooth Valley. Native fish present in Sawtooth Valley waters include: Chinook salmon *O. tshawytscha*, rainbow trout/steelhead *O. mykiss*, westslope cutthroat trout *O. clarkii lewisi*, bull trout *Salvelinus confluentus*, sucker *Catostomus spp.*, northern pikeminnow *Ptychocheilus oregonensis*, mountain whitefish *Prosopium williamsoni*, redbside shiner *Richardsonius balteatus*, dace *Rhinichthys spp.*, and sculpin *Cottus spp.* Nonnative species present in Sawtooth Valley waters include lake trout *S. namaycush* (Stanley Lake only), and brook trout *S. fontinalis*. Rainbow trout are released into Pettit, Alturas, and Stanley lakes in the

summer to increase sportfishing opportunities. Sportfishing on Pettit, Alturas, and Stanley lakes is covered by Idaho's statewide general fishing regulations, which allow harvest of six trout per day (excluding bull trout, which must be released if caught) and 15 kokanee per day with no seasonal closures. Sportfishing regulations on Redfish Lake restrict kokanee fishing/harvest to January 1 through August 7 to protect residual sockeye salmon. No trout have been stocked in Redfish Lake since 1992.

2006 and 2007 Captive Broodstock Program Egg and Juvenile Supplementation

All hatchery origin sockeye salmon released to Sawtooth Valley waters were adipose fin-clipped to distinguish hatchery rearing origin and/or release strategy. A subsample of some of the release groups was PIT tagged prior to release.

In 2006, 193,808 sockeye salmon and 184,596 sockeye salmon eyed-eggs were released into Sawtooth Valley waters from the captive broodstock program (Table 4). All presmolts released in 2006 were brood year 2005 (BY05) age-0 fish reared at IDFG Sawtooth Fish Hatchery (SFH) and were adipose fin clipped (AD). Redfish Lake received 61,804 presmolts (1,021 PIT tagged), Alturas Lake received 26,994 presmolts (1,016 PIT tagged), and Pettit Lake received 18,494 presmolts (1,021 PIT tagged) in October by direct lake releases. Smolts were released to the Salmon River on May 4, 2006 and to Redfish Lake Creek on May 10, 2006. A total of 39,622 BY04 smolts were released above the SFH weir. All fish released were ad-clipped and coded-wire tagged (AD/CWT), and 1,011 were PIT tagged before release. A total of 46,430 Oregon Department of Fish and Wildlife (ODFW) Oxbow Fish Hatchery (OFH) reared smolts were released into Redfish Lake Creek, with adipose and left ventral fin clips (AD/LV/CWT) for rearing site designation, of which 1,011 were PIT tagged before release. Four hundred sixty-four hatchery-produced adult sockeye salmon (all reared at NOAA Burley Creek Hatchery) were released to Redfish Lake for volitional spawning in September. In November and December 2006, 79,908 eyed-eggs (32,069 reared at NOAA Burley Creek Hatchery and 47,839 reared at Eagle Fish Hatchery [EFH]) were planted in Pettit Lake, and 104,688 eyed-eggs (77,482 reared at NOAA Burley Creek Hatchery and 27,206 reared at EFH) were planted in Alturas Lake.

In 2007, 184,275 sockeye salmon and 51,008 sockeye salmon eyed-eggs were released into Sawtooth Valley waters from the captive broodstock program (Table 5). All presmolts released in 2007 were age-0 fish from brood year 2006 (BY06) reared at SFH. Redfish Lake received 62,015 presmolts (988 PIT tagged), Alturas Lake received 9,977 presmolts (1,005 PIT tagged), and Pettit Lake received 10,113 presmolts (993 PIT tagged) in October by direct lake releases. Smolts were released to the Salmon River on May 8, 2007. A total of 101,676 BY05 smolts were released at two separate release sites (47,094 below the river water intake to SFH, and 54,582 into Redfish Lake Creek below the IDFG weir site). Smolts were reared at three locations: SFH, EFH, and OFH. All sockeye smolts released were coded-wire tagged, differentially marked (OFH fish were AD/RV clipped, SFH fish were AD-clipped, EFH fish were AD-clipped), and had representative groups PIT tagged (1,019 OFH, 1,025 SFH, 329 EFH). Four hundred ninety-four hatchery-produced adult sockeye salmon were released to Redfish Lake for volitional spawning in September. In November, 51,008 eyed-eggs reared at NOAA Burley Creek Hatchery were planted in Pettit Lake.

Table 1. Eyed-egg production by IDFG and NOAA facilities for the Snake River sockeye salmon captive broodstock program 1991—2007.

Brood year	Eyed-eggs produced by IDFG	Eyed-eggs produced by NOAA
1991	1,978	0
1992	36	0
1993	13,647	0
1994	259,536	48,000
1995	3,006	0
1996	110,756	381,500
1997	152,760	171,965
1998	15,580	47,533
1999	63,168	65,400
2000	253,047	94,500
2001	121,320	90,859
2002	66,324	60,516
2003	303,983	139,359
2004	140,823	135,699
2005	145,207	143,362
2006	258,342	190,603
2007	175,810	192,354
Total:	2,085,323	1,761,650

Table 2. Snake River sockeye salmon captive broodstock program egg and fish reintroduction history.

Year of reintroduction	Eyed-eggs	Presmolts	Smolts	Hatchery-reared adults	Anadromous Adults
1993	0	0	0	20	0
1994	0	14,119	0	65	0
1995	0	91,572	3,794	0	0
1996	105,000	1,932	11,545	120	0
1997	105,767	255,711	0	120	0
1998	0	141,871	81,615	0	0
1999	20,311	40,271	9,718	21	3
2000	65,200	72,114	148	271	200
2001	0	106,166	13,915	79	14
2002	30,924	140,410	38,672	190	12
2003	199,666	76,788	0	315	0
2004	49,134	130,716	96	241	0
2005	51,239	72,108	78,330	173	0
2006	184,596	107,292	86,052	464	0
2007	51,008	82,105	101,676	494	0
Total	862,845	1,333,175	425,561	2573	229

Table 3. Physical and morphometric characteristics of three study lakes located in the Sawtooth Valley, Idaho.

Surface Area (ha)	Elevation (m)	Volume (m ³ x 10 ⁶)	Mean Depth (m)	Maximum Depth (m)	Drainage Area (km ²)
615	1,996	269.9	<u>Redfish Lake</u> 44	91	108.1
338	2,138	108.2	<u>Alturas Lake</u> 32	53	75.7
160	2,132	45.0	<u>Pettit Lake</u> 28	52	27.4

Table 4. Sockeye salmon releases to Sawtooth Valley waters in 2006.

Release Location	Strategy (Brood Year)	Release Date	Number Released	Marks ^a	Number PIT-tagged	Mean Release Weight (g)	Rearing ^b Location
Salmon River (above SFH weir)	smolt (2004)	05/04/2006	39,622	AD/CWT	1,011	17.5	SFH
Redfish Lake Creek (Below weir)	Smolt (2004)	05/10/2006	46,430	ADRV/CWT	1,011	52.8	OFH
Alturas Lake (direct lake)	presmolt (2005)	10/02/2006	26,994	AD	1,016	6.1	SFH
Pettit Lake (direct lake)	presmolt (2005)	10/02/2006	18,494	AD	1,021	6.2	SFH
Redfish Lake (direct lake)	presmolt (2005)	10/03/2006	61,804	AD	1,021	6.2	SFH
Redfish Lake (direct lake)	adult (2002)	09/08/2006	38	AD	38	3,300.0	NOAA-MML
	(2002)	09/08/2006	11	AD	11	2,300.0	NOAA-BC
	(2003)	09/08/2006	174	AD	174	1,950.0	NOAA-MML
	(2003)	09/08/2006	144	AD	144	1,600.0	NOAA-BC
	(2003)	09/08/2006	73	AD	73	1,600.0	NOAA-BC
	(2004)	09/08/2006	16	AD	16	600.0	NOAA-MML
	(2004)	09/08/2006	8	AD	8	600.0	NOAA-BC
Pettit Lake (direct lake)	eyed-egg (2006)	11/21/2006	47,839	NA	NA	NA	EAG
	(2006)	12/6/2006	32,069	NA	NA	NA	NOAA-BC
Alturas Lake (direct lake)	eyed-egg (2006)	11/28/2006	38,992	NA	NA	NA	NOAA-BC
	(2006)	11/28/2006	17,884	NA	NA	NA	EAG
	(2006)	12/6/2006	38,490	NA	NA	NA	NOAA-BC
	(2006)	12/6/2006	9,322	NA	NA	NA	EAG

^a AD = adipose fin-clip, ADRV/CWT = adipose fin, right ventral fin-clip, coded wire tagged.

^b SFH = Idaho Department of Fish and Game Sawtooth Fish Hatchery; OFH = Oregon Department of Fish and Wildlife Oxbow Fish Hatchery; EAG = Idaho Department of Fish and Game Eagle Fish Hatchery; NOAA-BC = National Oceanic and Atmospheric Administration Burley Creek Hatchery; NOAA-MML = National Oceanic Administration Manchester Marine Lab.

Table 5. Sockeye salmon releases to Sawtooth Valley waters in 2007.

Release Location	Strategy (Brood Year)	Release Date	Number Released	Marks ^a	Number PIT-tagged	Mean Release Weight (g)	Rearing Location
Salmon River (above SFH weir)	Smolt (2005)	05/08/2007	46,765 329	AD/CWT AD	1,025 329	21.7 69.0	SFH EAG
Redfish Lake Creek (Below weir)	Smolt (2005)	05/08/2007	54,582	ADRV/CWT	1,019	51.6	OFH
Alturas Lake (direct lake)	Presmolt (2006)	10/02/2007	9,977	AD	1,005	7.1	SFH
Pettit Lake (direct lake)	Presmolt (2006)	10/03/2007	10,113	AD	993	7.0	SFH
Redfish Lake (direct lake)	Presmolt (2006)	10/03/2007	62,015	AD	988	6.7	SFH
Redfish Lake (direct lake)	Adult (2002)	09/05/2007	2	PIT	2	1,500.0	NOAA-BC
	(2003)	09/05/2007	53	PIT	53	3,200.0	NOAA-BC
	(2004)	09/05/2007	126	PIT	126	1,300.0	NOAA-BC
	(2004)	09/05/2007	182	PIT	182	1,300.0	NOAA-BC
	(2004)	09/06/2007	90	PIT/AD	90	1,126.0	IDFG-EAG
	(2004)	09/10/2007	41	PIT/AD	41	1,087.0	IDFG-EAG
Pettit Lake (direct lake)	Eyed-egg (2007)	11/27/2007	51,008	NA	NA	NA	NOAA-BC

^a AD = adipose fin-clip, ADRV/CWT = adipose fin, right ventral fin-clip, coded-wire tagged.

^b SFH = Idaho Department of Fish and Game Sawtooth Fish Hatchery; OFH = Oregon Department of Fish and Wildlife Oxbow Fish Hatchery; EAG = Idaho Department of Fish and Game Eagle Fish Hatchery; NOAA-BC = National Oceanic and Atmospheric Administration Burley Creek Hatchery; NOAA-MML = National Oceanic Administration Manchester Marine Lab.

Figure 1. Map of the upper Salmon River watershed located in the Sawtooth Valley, Idaho.

LITERATURE CITED

- Baker, D., J. Heindel, J. Redding, and P. A. Kline. In review. Snake River sockeye salmon captive broodstock program, hatchery element, 2006. Project no. 91-72. Portland, Oregon, Bonneville Power Administration, Annual Report.
- Baker, D., J. Heindel, J. Redding, and P. A. Kline. 2005a. Snake River sockeye salmon captive broodstock program, hatchery element, 2003. Project no. 91-72. Portland, Oregon, Bonneville Power Administration, Annual Report.
- Baker, D., J. Heindel, J. Redding, and P. A. Kline. 2005b. Snake River sockeye salmon captive broodstock program, hatchery element, 1999. Project no. 91-72. 2005. Portland, Oregon, Bonneville Power Administration, Annual Report.
- Bjornn, T. C., D. R. Craddock, and D. R. Corley. 1968. Migration and survival of Redfish Lake, Idaho, sockeye salmon, *Oncorhynchus nerka*. Transactions of the American Fisheries Society 97:360-375.
- Bowler, B. 1990. Additional information on the status of Snake River sockeye salmon. Idaho Department of Fish and Game. Boise, Idaho.
- Brannon, E. L., A. L. Setter, T. L. Welsh, S. J. Rocklage, G. H. Thorgaard, and S. A. Cummings. 1992. Genetic analysis of *Oncorhynchus nerka*. Project no. 199009300. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Brannon, E. L., T. Welsh, R. Danner, K. Collins, M. Casten, G. H. Thorgaard, K. Adams, and S. Cummings. 1994. Genetic analysis of *Oncorhynchus nerka*: Life history and genetic analysis of Redfish Lake *Oncorhynchus nerka*. Project no. 199009300. Bonneville Power Administration, Completion Report. Portland, Oregon.
- Burgner, R. L. 1991. Life History of Sockeye Salmon. Pages 3-117 in C. Groot, and L. Margolis, editors. Pacific Salmon Life Histories. University of British Columbia Press. Vancouver, British Columbia.
- Chapman, D. W., W. S. Platts, D. Park, and M. Hill. 1990. Status of Snake River sockeye salmon. Final report for Pacific Northwest Utilities Conference Committee. Don Chapman Consultants Inc. Boise, Idaho.
- Cummings, S. A., E. L. Brannon, K. J. Adams, and G. H. Thorgaard. 1997. Genetic analyses to establish captive breeding priorities for endangered Snake River sockeye salmon. Conservation Biology 11:662-669.
- Evermann, B. W. 1895. A preliminary report upon salmon investigations in Idaho in 1894. Bulletin of the United States Fish Commission 15:253-285.
- Evermann, B. W. 1896. A report upon salmon investigations in the headwaters of the Columbia River, in the state of Idaho, in 1895. U.S. Fish Commission Bulletin 16:151-202.
- Faler, J. C., and M. S. Powell. 2003. Genetic analysis of Snake River sockeye salmon (*Oncorhynchus nerka*). Bonneville Power Administration Annual Report. Portland, Oregon.

- Federal Register. 1991. Endangered and threatened species; endangered status for Snake River sockeye salmon-910379-1256. 91. Department of Commerce, National Oceanic and Atmospheric Administration, 50 CFR Part 222.
- Flagg, T. A. 1993. Redfish Lake sockeye salmon captive broodstock rearing and research, 1991-1992. Project no. 199204000. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Flagg, T. A., and W. C. McAuley. 1994. Redfish Lake sockeye salmon captive broodstock rearing and research, 1991-1993. Project no. 199204000. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Flagg, T. A., W. C. McAuley, M. R. Wastel, D. A. Frost, and C. V. W. Mahnken. 1996. Redfish Lake sockeye salmon captive broodstock rearing and research, 1994. Project no. 199204000. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Flagg, T. A., W. C. McAuley, D. A. Frost, M. R. Wastel, W. T. Fairgrieve, and C. V. W. Mahnken. 2001. Redfish Lake sockeye salmon captive broodstock rearing and research, 1995-2000. Project no. 199204000. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Frost, D. A., W. C. McAuley, D. J. Maynard, and T. A. Flagg. 2002. Redfish Lake sockeye salmon captive broodstock rearing and research, 2001. Project no. 199204000. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Griswold, R., A. Kohler, and D. Taki. 2000. Salmon River sockeye salmon habitat and limnological research: 1999 annual progress report. Project no. 199107100. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Hebdon, J. L., M. Elmer, and P. Kline. 2000. Snake River sockeye salmon captive broodstock program, research element, 1999. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Hebdon, J. L., J. Castillo, and P. Kline. 2002. Snake River sockeye salmon captive broodstock program, research element, 2000. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Hebdon, J. L., J. Castillo, C. Willard, and P. Kline. 2003. Snake River sockeye salmon captive broodstock program, research element, 2001. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Hebdon, J. L., P. A. Kline, D. Taki, and T. A. Flagg. 2004. Evaluating reintroduction strategies for Redfish Lake sockeye salmon captive broodstock progeny. American Fisheries Society Symposium 44: 401-413.
- Johnson, K. 1993. Research and recovery of Snake River sockeye salmon, 1992. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.

- Johnson, K., and J. Pravecek. 1995. Research and recovery of Snake River sockeye salmon, 1993. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Johnson, K., and J. Pravecek. 1996. Research and recovery of Snake River sockeye salmon, 1994-1995. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P. 1994. Research and recovery of Snake River sockeye salmon, 1993. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P., and J. Younk. 1995. Research and recovery of Snake River sockeye salmon, 1994. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P., and J. A. Lamansky. 1997. Research and recovery of Snake River sockeye salmon, 1995. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P., and J. Heindel. 1999. Snake River sockeye salmon captive broodstock program, hatchery element, 1998. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P., and C. Willard. 2001. Snake River sockeye salmon captive broodstock program, hatchery element, 2000. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P., J. Heindel, and C. Willard. 2003a. Snake River sockeye salmon captive broodstock program, hatchery element, 1997. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P., C. Willard, and D. Baker. 2003b. Snake River sockeye salmon captive broodstock program, hatchery element, 2001. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kohler A., B. Griswold, and D. Taki. 2001. Snake River sockeye salmon habitat and limnological research: 2000 annual progress report. Project no. 199107100. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kohler A., D. Taki, and B. Griswold. 2002. Snake River sockeye salmon habitat and limnological research: 2001 annual progress report. Project no. 199107100. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Lewis, B., D. Taki, and R. Griswold. 2000. Snake River sockeye salmon habitat and limnological research: 2000 annual progress report. Project no. 199107100. Bonneville Power Administration, Annual Report. Portland, Oregon.
- National Oceanic and Atmospheric Administration (NOAA). 2005. Endangered Species Act status review and listing information. www.nwr.noaa.gov/1salmon/salmesa/index.htm.

- NMFS. 2002. Interim Abundance Targets. Seattle, Washington, National Oceanic and Atmospheric Administration.
- Pravecek, J., and K. Johnson. 1997. Research and recovery of Snake River sockeye salmon, 1995. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Pravecek, J., and P. Kline. 1998. Research and recovery of Snake River sockeye salmon, 1996. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Taki, D., and A. Mikkelsen. 1997. Salmon River sockeye salmon habitat and limnological research: 1996 annual progress report. Project no. 199107100. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Taki, D., B. Lewis, and R. Griswold. 1999. Salmon River sockeye salmon habitat and limnological research: 1997 annual progress report. Project no. 199107100. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Teuscher, D., and D. Taki. 1995. Salmon River sockeye salmon habitat and limnological research, In Teuscher, D., and D. Taki (ed). Snake River sockeye salmon habitat and limnological research: 1994 annual progress report. Project no. 199107100. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Teuscher, D., and D. Taki. 1996. Salmon River sockeye salmon habitat and limnological research, In Teuscher, D., and D. Taki (ed). Snake River sockeye salmon habitat and limnological research: 1995 annual progress report. Project no. 199107100. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Waples, R. S. 1991. Definition of a "species" under the Endangered Species Act: Application to Pacific Salmon. Seattle, Washington, U. S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum NMFS-F/NWC 194.
- Waples, R. S., O. W. Johnson, and R. P. Jones Jr. 1991. Status review for Snake River sockeye salmon. Seattle, Washington, U. S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum NMFS-F/NWC 195.
- Waples, R. S., P. B. Aebersold, and G. A. Winans. 1997. Population genetic structure and life history variability in *Oncorhynchus nerka* from the Snake River Basin. Project no. 93-068. Portland, Oregon, Bonneville Power Administration Annual Report.
- Willard, C., D. Baker, J. Heindel, J. Redding, and P. Kline. 2003a. Snake River sockeye salmon captive broodstock program, hatchery element, 2002. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Willard, C., J. L. Hebdon, J. Castillo, J. Gable, and P. Kline. 2003b. Snake River sockeye salmon captive broodstock program, research element, 2002. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.

- Willard, C., K. Plaster, J. Castillo, and P. Kline. 2005. Snake River sockeye salmon captive broodstock program, research element 2003. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Winans, G. A., P. A. Aebersold, and R. S. Waples. 1996. Allozyme variability in selected populations of *Oncorhynchus nerka* in the Pacific Northwest, with special consideration of populations of Redfish Lake, Idaho. Transactions of the American Fisheries Society 205:645-663.

PART 2—ONCORHYNCHUS NERKA POPULATION MONITORING AND REDFISH LAKE SPORT FISHERY INVESTIGATIONS

INTRODUCTION

Understanding the dynamics of *O. nerka* populations in the Sawtooth Valley lakes is a vital part of sockeye salmon restoration efforts. Knowledge of *O. nerka* abundance coupled with limnology data (collected and reported by the Shoshone-Bannock Tribes [SBT]) is necessary for making responsible decisions regarding the reintroduction of sockeye salmon juveniles from the captive broodstock program. Utilizing multiple release strategies at various life stages allows program managers to design reintroduction plans that take advantage of the nursery lakes' current carrying capacities, which are estimated by trawling, hydroacoustic surveys, and limnological surveys (hydroacoustic and limnological surveys conducted and reported by the SBT). Productivity in the lakes varies annually and the presence of kokanee in the nursery lakes creates increased competition for limited food sources between kokanee and sockeye salmon. During years in which a lake is experiencing low productivity and/or high kokanee abundance, the program limits reintroduction efforts of individuals that would spend more time in nursery lakes and diverts individuals to more productive lakes.

The kokanee fishery on Redfish Lake was closed in 1993 due to the presence of ESA listed residual sockeye salmon but was reopened in 1995 (NOAA Permit 1481). The kokanee fishery was reopened based on the recommendation of the SBSTOC to reduce kokanee competition with sockeye salmon by removing spawning age kokanee through angler harvest. Permit 1481 (NOAA) requires IDFG to monitor angler harvest of listed sockeye salmon in Redfish Lake during the kokanee fishing season. The kokanee season on Redfish Lake opens on January 1 and closes on August 7, when mature kokanee initiate spawning in Fishhook Creek, while residual sockeye salmon remain in the lake.

The roving creel survey conducted on Redfish Lake was designed to estimate total kokanee harvest and to collect tissue samples for genetic analysis from angler-harvested kokanee. The tissue samples were collected for genetic analysis to estimate the number of unmarked sockeye salmon harvested incidental to the kokanee fishery.

METHODS

Oncorhynchus Nerka Population Monitoring

To estimate *O. nerka* (kokanee and sockeye salmon) abundance, density, and biomass in Sawtooth Valley lakes, midwater trawling was conducted at night during the dark (new) phase of the moon in September. Spawning-age kokanee (>250 mm fork length) in Redfish and Alturas lakes migrate to tributaries to spawn in August; therefore, trawling is conducted in September to prevent the collection of biased trawl catch data and because juvenile *O. nerka* that remain in valley lakes are tightly stratified during this time of the year. Redfish, Pettit, and Alturas lakes were sampled September 10-12, 2007. Trawling was performed in a stepped-oblique fashion as described by Rieman (1992) and Kline (1994). A minimum of six trawl transects were conducted per lake. Total *O. nerka* abundance, density, and biomass were estimated using a program developed by Rieman (1992). Abundance estimates generated by this program are extrapolations of actual trawl catch data to the total area of the lake mid-depth in the observed *O. nerka* stratum. Density and biomass estimates are expressed in relation to

lake surface area. Whenever possible, we estimated abundance, density, and biomass by individual age class (assuming representation in the trawl). We calculated confidence intervals using the following formula of Scheaffer, Mendenhall, and Ott (1990):

$$\bar{x} \pm t \sqrt{\frac{s^2}{n}}$$

Fork length (1 mm) and weight (0.1 g) were recorded for all trawl-captured *O. nerka*. Sagittal otoliths and scales were removed from a subsample of *O. nerka* (a minimum of five fish from every 10 mm length group) and returned to the laboratory. Three program technicians aged scales and otoliths to determine length ranges for age classification. Scales were mounted between microscope slides before aging and otoliths were read “dry.” Complete fish carcasses were collected and sent to the University of Idaho’s Center for Salmonid and Freshwater Species at Risk for proximate analysis (proximate analysis results are presented under a separate report cover produced by the University of Idaho). Stomachs were removed and preserved for diet analysis by SBT biologists. Heads were removed and submitted to IDFG’s Eagle Fish Health Lab for whirling disease testing (results are covered under a separate cover; see Baker et al. *in review*) because the Myxosporean parasite, *Myxobolus cerebralis*, which can cause salmonid whirling disease, is present in the upper Salmon River. Fin clips were stored in 100% ethanol and delivered to the IDFG Genetics Laboratory for DNA analysis.

Redfish Lake Sport Fishery Investigations

A roving creel survey was conducted from May 28 through August 7, 2007 (kokanee harvest closes on August 7 to protect residual sockeye salmon). The creel census was stratified by 14-day intervals, broken into weekday and weekend day types and morning (0800 to 1400) and evening (1500 to 2000) instantaneous count periods. Angler counts were conducted two weekdays, two weekend days, and any holiday during each 14-day interval. On each angler count day, the number of boats and bank anglers were counted from a boat for each day period (morning and evening strata). Angler count dates and times were selected randomly. Angler interviews were conducted following the completion of each instantaneous count. Anglers were asked how many fish they had harvested and/or released by species, how many hours they had fished, what their preferred target species was, whether or not they were aware of the Redfish Lake kokanee fishery, and the type of gear they used. All responses were recorded by creel personnel. Fin clips were taken from harvested kokanee that were checked by creel survey personnel. Fin clips were stored in 100% ethanol and delivered to IDFG Genetics Laboratory personnel for DNA analysis. Creel data were analyzed using the Creel Census System computer program developed by McArthur (1992) and used to estimate angler effort, catch rates, and harvest.

RESULTS AND DISCUSSION

Oncorhynchus Nerka Population Monitoring

Redfish Lake

September trawl catch (six transects, Appendix A) included 53 natural origin *O. nerka* and zero hatchery origin sockeye salmon. *Oncorhynchus nerka* abundance was estimated at

73,702 fish (95% CI $\pm$ 24,195). The *O. nerka* population decreased 11% within Redfish Lake over the estimated abundance in 2006 of 82,796 fish; however, abundances were 24% greater than the estimate for 2005 of 56,220 *O. nerka* (Table 6).

Density and biomass were estimated at 119.84 fish/ha and 0.84 kg/ha, respectively (Table 6). This represented decreases of 11% for densities and a 65% decrease in biomass from estimated levels in 2006 (Peterson et al. 2007). Age-0, age-1, and age-2 *O. nerka* were represented in the trawl sample from Redfish Lake (age-3 fish were missing from the trawl, which may explain the decrease in biomass from 2006). Age-0 fish had the highest density (96.83 fish/ha) and age-2 had the highest biomass (0.72 kg/ha) (Table 7).

Alturas Lake

September trawl catch on Alturas Lake (six transects, Appendix A) included 176 natural origin *O. nerka* and zero hatchery origin sockeye salmon. We estimated *O. nerka* abundance, density, and biomass at 124,073 fish (95% CI $\pm$ 23,327), 367.1 fish/ha, and 3.43 kg/ha, respectively (Table 6). Age-0, age-1, age-2, and age-3 *O. nerka* were represented in the trawl sample. Age-1 fish had the highest density (177.14 fish/ha) and contributed 40.2% of the biomass (Table 7). The estimates for abundance and density represent a 17.3% increase over 2006 estimates, while the biomass estimate of 3.43 kg/ha represents a decrease of 2.3% from 2006 (Table 6). Data collected from the SBT indicates we may have overharvested zooplankton within the lake and we may need to decrease sockeye releases within Alturas Lake (Robert Griswold, personal communication).

Kokanee control within the three nursery lakes has proven to be difficult because of many factors such as weir failure, otter damage to weirs and traps, and the lack of a permanent sill structure (Kohler, personal communication). Kokanee control should remain as a priority for the program to control zooplankton abundance and minimize *O. nerka* population fluctuations.

Pettit Lake

September trawl catch on Pettit Lake (six transects, Appendix A) included 20 natural origin *O. nerka* and one hatchery origin (adipose fin-clipped) sockeye salmon. We estimated *O. nerka* abundance, density, and biomass at 14,746 fish (95% CI $\pm$ 7,099), 92.16 fish/ha, and 3.84 kg/suitable ha, respectively (Table 6). Age-1 and age-2 fish were represented in the trawl sample. Age-2 fish had the highest density (64.94 fish/ha) and contributed 82.5% of the biomass (Table 7). The estimates for abundance, density, and biomass represent decreases over 2006 estimates of 55.6%, 55.6%, and 48.1%, respectively. This is likely due to sampling bias and missing the age-0 class of fish completely. Age-0 size fish were seen in hydroacoustic sampling conducted by the tribe during the fall of 2007 (conducted by the SBT, reported under a separate cover).

The capture of one hatchery origin sockeye salmon during trawl sampling along with the capture of another hatchery origin sockeye salmon during a hook-and-line sampling trip may suggest that the number of residual sockeye salmon within Pettit Lake is on the rise. After genetic analysis of trawl captured fish from 2003 and 2006, we identified that the proportion of sockeye sampled from within the lake has increased from approximately 8% in 2003 to 58% in 2006. We recommend that SBSTOC develop and discuss options to evaluate and monitor whether this population is increasing. The SBT conducts night snorkeling on Redfish Lake to

monitor the residual population within the lake, which may prove to be a feasible way to monitor the population on Pettit Lake.

With the additional knowledge we have gained by using the new trawling gear for two years, it appears we have also increased our ability to capture fish from the older age classes. Because these older age classes were missing from earlier datasets, it was difficult to accurately monitor a specific age class of kokanee through maturation. We are able to trawl 50% faster with the new gear, which may increase our capture efficiencies in the older age classes. We believe using the new trawling gear may provide us with more precise estimates of the older age classes of *O. nerka* found within Redfish, Alturas, and Pettit lakes. The estimates of older age (age-2 and older) *O. nerka* are likely the kokanee component of the population found within Redfish and Alturas and may be residual production within Pettit Lake.

It is possible that *O. nerka* populations within the basin lakes are expanding in response to fertilization (conducted by the SBT, reported under a separate cover), and sampling bias is not the cause of these rapid population fluctuations. Zooplankton abundance levels have remained high enough for *O. nerka* growth to occur within each lake (Robert Griswold, personal communication). We have also experienced high levels of adult escapement in both Fishhook Creek and Alturas Lake Creek over the past couple of years (Table 8), which may have led to more recruitment into the systems.

Whether population fluctuations between 2006 and 2007 were a sampling response or a biological response, the estimates of these fish will help us to develop escapement goals, which are necessary to minimize kokanee and zooplankton population fluctuations. These goals should also provide a stable nursery habitat environment for sockeye salmon by reducing competition between kokanee and sockeye. Escapement goals will be established during the summer of 2008 (before kokanee enter the tributaries to spawn) and presented to the SBSTOC for discussion and modification.

Genetic samples collected from trawl-captured fish in 2007 have not been analyzed for any of the three basin lakes as of the completion of this report.

Redfish Lake Sport Fishery Investigations

In 2007, we contacted 53 angler parties (102 individual anglers) on Redfish Lake. Idaho residents made up 84% of those interviewed. Most anglers used lures (91%), followed by bait (6%). Total angler effort was estimated at 1,922 hours (95% CI $\pm$ 139) (Table 9). This effort estimate represents a 27% decrease in fishing pressure between 2006 and 2007 (Table 10). Boat anglers expended more effort (67%) than bank anglers (31.8%), which was similar to results from 2006 (Peterson et al. 2007).

The season catch rate for all fish (harvested and released) was 0.34 fish/hour. Catch rates decreased 53% over the estimate from 2006 (Peterson et al. 2007). Kokanee catch rates (harvested and released) averaged 0.14 kokanee/hour for weekdays and 0.09 kokanee/hour for weekends (0.12 kokanee/hour for the season, Table 11). These estimates were also 61% lower than the estimate of 0.31 kokanee/hour for the season in 2006. Bull trout catch and release rates averaged 0.18 bull trout/hour for weekdays and 0.04 bull trout/hour for weekends (IDFG regulations prohibit harvesting bull trout) (Table 11). We also estimated that 11 bull trout were harvested from Redfish Lake during 2007. Only two bull trout were actually observed as harvested by creel technicians. The season catch rate for bull trout is slightly higher than catch

rates observed in 2005. We have seen a trend of overall increase in bull trout catch rates since the harvest closure in 1994 (Figure 2). This trend will be discussed further in Part 4 of this report under Predator Surveys. Other fish (cutthroat, northern pikeminnow, redbside shiner, sucker *catostomus spp.*) accounted for a combined catch rate (harvested and released) of 0.09 fish/hour for the season.

The total number of fish caught (harvested and released) in Redfish Lake was estimated at 627 fish (95% CI $\pm$ 173.7). This was an overall decrease of 72% from the 2006 estimate. The majority (84%) of all caught fish were released. Kokanee harvest was estimated at 56 fish (95% CI $\pm$ 18) and the number of kokanee released was estimated at 174 (95% CI $\pm$ 37) (Table 12). The harvest of kokanee declined an estimated 75%; however, the number harvested versus released was proportionately similar to 2006 harvest and release numbers. In 2007, we observed the lowest harvest rate for kokanee since 2002 and the third lowest rate since the re-establishment of the kokanee fishery in 1995 (Table 10). Because of the low harvest rate and increased rate of released kokanee by anglers, we initiated an angler education program in the basin lakes to explain the importance of harvesting kokanee to reduce competition between the two forms of *O. nerka* found in the lakes (Peterson et al. 2007). This program began in the spring of 2007 and will continue in 2008. We hope to determine if any additional fishing pressure is applied to Redfish Lake as a result of the program.

No hatchery-produced sockeye salmon (adipose-clipped) were observed by creel personnel during 2007. The direct impact of the kokanee fishery on residual sockeye salmon (through incidental harvest) is evaluated annually using genetic analysis of tissue samples collected from kokanee in the creel. Genetic analysis of fin clips obtained from harvested kokanee (n = 8) has not been completed for 2007.

Table 6. Estimated *O. nerka* population, density, and biomass for Redfish, Alturas, and Pettit lakes, 1990 to 2007.

Year	Population ($\pm$ 95% CI)	Density (fish/ha)	Biomass (kg/ha)
<u>Redfish Lake (615 surface hectares)</u>			
2007	73,702 (24,195)	119.8	0.84
2006	82,796 (47,407)	134.6	2.37
2005	56,220 (4,192)	91.4	0.3
2004	82,258 (3,486)	133.0	0.3
2003	81,727 (2,763)	132.9	1.6
2002	50,204 (4,085)	81.6	1.0
2001	12,980 (2,959)	21.1	<0.1
2000	10,268 (1,605)	16.7	<0.1
1999	42,916 (1,795)	69.7	0.9
1998	31,486 (1,716)	51.2	1.8
1997	55,762 (1,590)	90.7	2.5
1996	56,213 (3,526)	91.4	2.8
1995	61,646 (2,078)	100.2	4.4
1994	51,529 (4,902)	83.8	1.4
1993	49,628 ^a	80.7	1.6
1992	39,481 (2,498)	64.2	1.0
1990	24,431 (11,000)	39.7	0.8
<u>Alturas Lake (338 surface hectares)</u>			
2007	124,073 (23,327)	367.1	3.43
2006	105,779 (50,702)	313.0	3.51
2005	20,956 (2,136)	98.8	0.3
2004	36,206 (2,579)	107.1	1.9
2003	46,234 (5,183)	136.8	5.5
2002	24,374 (2,328)	72.1	2.2
2001	70,159 (1,696)	207.6	2.4
2000	125,462 (1,572)	371.0	2.1
1999	56,675 (4,476)	167.7	0.4
1998	65,468 (2,860)	193.7	1.4
1997	9,761 (933)	28.9	2.1
1996	13,012 (691)	38.5	1.4
1995	23,061 (1,202)	68.2	1.7
1994	5,785 (1,957)	17.1	0.4
1993	49,037 (1,443)	145.1	2.6
1992	47,237 (3,782)	139.8	2.4
1991	125,045 (1,881)	370.0	3.9
1990	126,644 (1,690)	374.7	3.3
<u>Pettit Lake (160 surface hectares)</u>			
2007	14,746 (7,099)	92.2	3.84
2006	33,246 (12,416)	207.8	7.4
2005	23,970 (2,136)	149.8	2.2
2004	46,065 (3,288)	287.9	9.8
2003	11,961 (626)	136.8	5.5
2002	18,328 (384)	114.5	12.1
2001	16,931 (1,311)	105.8	6.1
2000	40,559 (1,317)	253.5	10.2
1999	31,422 (2,515)	196.4	6.3
1998	27,654 (862)	172.8	9.7
1997	21,730 (1,462)	135.8	5.1
1996	71,654 (911)	447.8	15.3
1995	59,002 (1,653)	368.8	14.7
1994	14,743 (1,966)	92.1	3.1
1993	10,511 (640)	65.7	0.8
1992	3,009 (539)	18.8	2.5

^a Confidence limits not calculated—single transect estimate.

Table 7. Estimated 2007 *O. nerka* abundance, density (fish/ha), and biomass (kg/ha) by age class in Redfish, Alturas, and Pettit lakes.

	Age-0	Age-1	Age-2	Age-3	Age-4	Total
Redfish Lake (615 surface ha)						
No. captured	41	7	3	0	NA	53
Mean length (mm) (± 95 CI)	54.9	99.9	183.7	NA	NA	NA
Mean weight (g) (± 95 CI)	1.63	12.04	71.00	NA	NA	NA
Abundance	59,556	9,957	4,188	NA	NA	73,702
95% CI High	76,745	18,626	11,612	NA	NA	97,897
95% CI Low	42,367	1,288	0	NA	NA	49,507
Density (fish/ha)	97	16	7	NA	NA	119.84
Biomass (kg/ha)	0.25	0.29	0.72	NA	NA	0.84
Alturas Lake (338 surface ha)						
No. captured	110	41	21	4	0	176
Mean length (mm) (± 95 CI)	63.9	94.9	149.8	183.7	NA	NA
Mean weight (g) (± 95 CI)	2.30	8.26	37.19	65.75	NA	NA
Abundance	52,056	59,874	9,301	2,843	NA	124,073
95% CI High	83,488	80,952	17,090	5,154	NA	147,400
95% CI Low	20,624	38,796	1,512	532	NA	100,746
Density (fish/ha)	154	177	28	8	NA	367.08
Biomass (kg/ha)	0.23	1.38	1.23	0.59	NA	3.43
Pettit Lake (160 surface ha)						
No. captured	0	2	18	0	0	20
Mean length (mm) (± 95 CI)	NA	120	155.7	NA	NA	NA
Mean weight (g) (± 95 CI)	NA	22.37	44.3	NA	NA	NA
Abundance	NA	4,355	10,390	NA	NA	14,746
95% CI High	NA	10,266	12,695	NA	NA	21,845
95% CI Low	NA	0	8,085	NA	NA	7,647
Density (fish/ha)	NA	27.21	64.94	NA	NA	92.16
Biomass (kg/ha)	NA	0.67	3.17	NA	NA	3.84

Table 8. Estimated kokanee escapement to Fishhook Creek 1991 to 2007 and Alturas Lake Creek 1992 to 2007. Data obtained from the Shoshone-Bannock Tribes.

Year	Fishhook Creek	Alturas Lake Creek
1991	7,200	No survey data
1992	9,600	60
1993	10,800	200
1994	9,200	3,200
1995	7,000	1,600
1996	10,662	744
1997	8,572	8,492
1998	6,149	15,237
1999	2,336	8,334
2000	60	827
2001	5,853	145
2002	8,626	99
2003	9,679	48
2004	1,508	7,101
2005	4,375	11,652
2006	14,021	2,276
2007	11,235	519

Table 9. Estimated angler effort on Redfish Lake for the 2007 fishing season.

Redfish Lake	Boat	Bank	Tube	Total
Hours fished	1,296	613	13	1,922
±95% CI	159	120	8	139

Table 10. Historical kokanee catch rates, kokanee harvest estimates, bull trout catch rates, and angler effort for the Redfish Lake fishery.

	Kokanee catch rates (fish/hour)		Kokanee harvested	Bull trout catch rate (fish/hr)	Angler parties interviewed	Estimated hours fished/season
	Harvested (fish/hour)	Released (fish/hour)				
1996	0.19	0.08	844	0.09	107	3,351
1997	0.19	0.37	466	0.08	117	2,874
1998	0.13	0.17	1,362	0.08	205	7,963
1999	0.38	0.15	1,187	0.28	227	3,951
2000	0.02	0.06	67	0.08	63	3,063
2001	0.00	0.06	0	0.27	88	2,391
2002	0.09	0.16	129	0.16	100	2,127
2003	0.10	0.05	424	0.24	98	2,477
2004	0.13	0.26	621	0.31	96	2,791
2005	0.21	0.09	637	0.09	85	3,620
2006	0.07	0.24	222	0.35	131	2,635
2007	0.03	0.09	56	0.12	53	1,922

Table 11. Catch rates (fish/hour) for summer 2007 on Redfish Lake categorized by day type and species.

Day Type	Kokanee		Cutthroat Trout		Bull Trout		All Fish	
	Kept	Released	Kept	Released	Kept	Released	Kept	Released
Weekday	0.03	0.11	0.04	0.10	0.01	0.18	0.08	0.39
Weekend Day	0.03	0.06	0.00	0.01	0.00	0.04	0.06	0.12
Season Avg.	0.03	0.09	0.02	0.06	0.01	0.12	0.07	0.27

Table 12. Estimated number of fish harvested and released on Redfish Lake during the summer 2007.

Redfish Lake	Harvested				All Fish ± 95% CI
	Kokanee	Cutthroat Trout	Bull Trout	Other	
Harvested	56	20	11	12	99 ± 33.44
Released	174	89	266	0	529 ± 161.09

Figure 2. Estimated bull trout catch rates from 1996-2007 within Redfish Lake. The trend line is used to show the population increase within the lake. The equation for the trend line is $y = 0.0117x + 0.103$ with an R^2 of 0.17.

LITERATURE CITED

- Baker, D., J. Heindel, D. Green, and P. Kline. *In review*. Snake River sockeye salmon captive broodstock program, hatchery element 2005. IDFG Report No. 07-25. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P. A. 1994. Research and recovery of Snake River sockeye salmon. Idaho Department of Fish and Game. Annual Report to U.S. DOE, Bonneville Power Administration, Division of Fish and Wildlife. Project No. 91-72, Contract No. DE-BI79-91BP21065. Portland, Oregon.
- McArthur, T. 1992. Statewide angler opinion and harvest surveys. Idaho Department of Fish and Game, Project No. F-71-R-14, Subproject I, Study I. Boise, Idaho.
- Peterson, M., B. Moore, K. Plaster and P. Kline. 2007. Snake River sockeye salmon captive broodstock program, research element 2006. IDFG Report no. 07-28. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Rieman, B. E. 1992. Kokanee salmon population dynamics—kokanee salmon monitoring guidelines. Idaho Department of Fish and Game, Project No. F-73-R-14, Subproject II, Study II. Boise, Idaho.
- Scheaffer, Mendenhall, and Ott. Elementary Survey Sampling. 1990. Duxbury Press.

PART 3—SCKEYE SALMON JUVENILE OUT-MIGRANT AND ADULT SPAWNING MONITORING AND EVALUATION

INTRODUCTION

SCKEYE SALMON JUVENILE OUT-MIGRANT MONITORING AND EVALUATION

Monitoring overwinter survival and out-migration of sockeye salmon smolts plays an important role in restoration efforts. Trapping conducted on the lake outlet streams provides information on timing of out-migration and smolt sizes. Out-migrant monitoring provides an opportunity to monitor natural production of sockeye salmon in the lakes and to evaluate the success of different release strategies. This information allows us to make informed decisions regarding the placement of future captive broodstock progeny.

Out-migrant trapping also provides overwinter survival information for presmolts (fall direct-release) released into the nursery lakes in October the year prior to out-migrant trapping. Hatchery origin sockeye salmon smolts captured at lake out-migrant traps originated primarily from the October 2006 release of adipose fin-clipped presmolts numbering 61,804 sockeye, 26,994 sockeye, and 18,494 sockeye to Redfish, Alturas, and Pettit lakes, respectively. All presmolts released in 2006 were reared at the Sawtooth Fish Hatchery (SFH).

ADULT SPAWNING MONITORING AND EVALUATION

Releasing mature adult sockeye salmon into Sawtooth Valley lakes has been an important part of the "spread-the-risk" philosophy of the SBSTOC. Prespawn sockeye salmon adults were first released back to the wild in 1993. Adult sockeye salmon raised to maturity in the hatchery and released to valley lakes to spawn provide a "natural" or unmarked smolt component that is subject to natural selection. Beginning in 1999, hatchery origin anadromous sockeye salmon have been released into valley lakes (when available) along with adult sockeye salmon that were raised to maturity in a hatchery. Currently, prespawn adults are released only to Redfish Lake.

Success of releasing hatchery-produced adults to spawn naturally is evaluated by determining if there is a corresponding increase in production in the number of unmarked smolts out-migrating and observed egg-to-smolt survival. Unmarked out-migrants can be progeny of residual sockeye salmon adults that spawn in basin lakes; hatchery origin prespawn adults released to the lakes for natural spawning; anadromous adults released to the lakes for natural spawning; eyed-egg releases; or misclipped hatchery origin smolts. Juvenile kokanee (nonanadromous) could also "fall out" of nursery lakes and contribute to trap counts. The weirs on Redfish Lake Creek and Pettit Lake Creek and the screw trap on Alturas Lake Creek enable us to monitor and estimate unmarked out-migrating smolts and obtain genetic samples.

Current evaluations of adult sockeye salmon releases focus on the number of redds produced and estimations of unmarked juvenile out-migrants and collection of genetic material for future DNA parental exclusion analysis.

METHODS

SOCKEYE SALMON JUVENILE OUT-MIGRANT MONITORING AND EVALUATION

Redfish Lake Creek Trap

The out-migrant trap on Redfish Lake Creek (RLCTRP) is located 1.4 km downstream from the lake outlet at a permanent weir site. The trap functions as a juvenile trap for out-migrating fish, and with only minor modifications, as a trap for returning adults (Craddock 1958; Bjornn et al. 1968; Kline 1994; Kline and Younk 1995; Kline and Lamansky 1997; Hebdon et al. 2000, 2002, 2003; Willard et al. 2004, 2005). The trap was operated from April 14 through June 10, 2007 and contains nine bays, five of which were fitted with incline bar traps. IDFG staff checked the trap at least twice daily in 2007. The trap is fished until high water forces us to remove it, until fish stop emigrating from the lake, or until June 15 (contract deadline for removal of trap and the placement of the adult trap).

Each fishing bay was fitted with an adjustable 1.70 m wide by 1.74 m long aluminum trap box on a winch and pulley system. The trap boxes were constructed of 3 mm aluminum sheeting and framework and 1.9 cm diameter hollow aluminum bars. The 30.5 cm x 169.5 cm x 30.5 cm live wells were also constructed of 3 mm aluminum with 5 mm holes drilled for aeration and water exchange in the live well. Bar spacing (19 mm) allowed debris and large fish to pass downstream, while low velocity water swept *O. nerka* smolts across the bars and into the live well for holding until personnel were able to empty the trap (Kline 1994).

All sockeye salmon smolts captured at RLCTRP were anesthetized in buffered tricaine methanesulfonate (MS-222), measured for fork length (1 mm), weight (0.1 g), and scanned for PIT tags. Scales were removed from a subsample of natural origin and adipose fin-clipped hatchery reared *O. nerka* (five fish from each 5 mm length group) and returned to the laboratory for aging. In the lab, scales were pressed between microscope slides, two program employees individually aged the scales, and a third person aged the discrepancies. The proportions of age-1 and age-2 out-migrants were determined by using the MIX computer program developed by MacDonald and Green (1988). MIX software uses known values (the scale ages in this case) and fits mixture distributions to grouped data by utilizing a maximum likelihood estimator. All captured nontarget species were counted and released immediately. Fin-clip samples were taken from all wild fish for future genetic analyses (Redfish Lake = 841, Alturas Lake = 94, Pettit Lake = 90). Tissue samples and complete fish carcasses were collected and sent to the University of Idaho's Center for Salmonid and Freshwater Species at Risk for proximate analysis (N = 48). These fish were measured for fork length (1 mm) and weighed to the nearest 0.1 g. Proximate sampled fish were sacrificed or were trap mortalities.

To estimate trapping efficiency, up to 20 natural origin sockeye salmon smolts (determined by presence of an adipose fin) and 20 hatchery origin sockeye salmon smolts were PIT tagged daily and released approximately 250 m upstream of the weir one-half hour after sunset. All remaining fish were identified (natural origin and hatchery origin), counted, and released 15 m below the weir one-half hour after sunset. Flow-through live boxes with locking lids were used to hold fish until the evening release. Trapping efficiencies were calculated for natural origin sockeye salmon smolts and fall direct-released sockeye salmon smolts. Intervals were selected based on stream discharge similarities and the number of PIT tagged smolts released upstream of the weir that were available for recapture (trap efficiencies). Natural origin fish typically out-migrate earlier in the season than fall direct-release fish (Figure 3). Stream velocity was measured below the trap weekly. Out-migrant run size was derived using a

modified Bailey estimator and 95% bootstrap confidence intervals using methods described by Steinhorst et al. (2004). Smolt out-migration estimates were calculated separately for natural origin and fall direct-released hatchery origin sockeye salmon smolts. During the spring of 2007, we experienced normal flow conditions throughout the trapping season.

Alturas Lake Creek Trap

Sockeye salmon out-migrant trapping and PIT tagging on Alturas Lake Creek was conducted by the SBT. The Alturas Lake Creek screw trap is located 13 km downstream from the Alturas Lake outlet and was operated from April 19 to May 23, 2007. Hatchery-produced sockeye salmon smolts captured at the trap originated primarily from 26,994 adipose fin-clipped presmolts (reared at SFH) released into the lake in October 2006. The Alturas Lake out-migrant population estimate was derived using the same estimator described above (Steinhorst et al. 2004) except under extreme flow conditions when other methods are used to estimate out-migrant numbers, which would be presented under a separate cover provided by the SBT.

Pettit Lake Creek Trap

Sockeye salmon out-migrant trapping and PIT tagging on Pettit Lake Creek was conducted by the SBT. The Pettit Lake Creek trap is located 1 km downstream from the Pettit Lake outlet at a permanent weir site and was operated from April 18 to May 29, 2007. Hatchery-produced sockeye salmon smolts captured at the trap originated primarily from the 18,494 adipose fin-clipped presmolts (reared at SFH) released into the lake in October 2006. The Pettit Lake Creek weir traps at 100% efficiency under low spring flow conditions (personal communication, Doug Taki, SBT); therefore, out-migration run size for Pettit Lake is based on the actual number of smolts trapped. However, during normal to high flow years, the trap must be removed and other means are used to estimate the number of out-migrants, which would be presented under a separate cover provided by the SBT.

Salmon River and Redfish Lake Creek Smolt Groups

Forty-seven thousand ninety-four (47,094) BY05 smolts were released near the river water intake of SFH. This release group included smolts reared at SFH (46,765) and Eagle Fish Hatchery (EFH) (329). All fish released were adipose fin-clipped and coded-wire tagged with a representative group (n = 1,354) PIT tagged.

A total of 54,582 BY05 smolts were released into Redfish Lake Creek below the out-migrant weir. This release included smolts reared at Oregon Department of Fish and Wildlife Oxbow Fish Hatchery (OFH). All fish were adipose/right ventral fin-clipped and coded-wire tagged with a representative group (n = 1,019) PIT tagged.

Mainstem Snake and Columbia River Dams

In 2007, the majority of sockeye salmon smolts were transported and released below Bonneville Dam; this included PIT tagged fish. As a result, mainstem survival evaluations were only conducted to Lower Granite Dam (LGR). Sockeye salmon smolt survival to LGR was evaluated using PIT tag interrogation data collected at PIT tag detection facilities throughout the Snake and Columbia rivers. Interrogation data were retrieved from the PIT tag information system (PTAGIS) maintained by the Pacific States Marine Fisheries Commission (Portland,

Oregon). This data was used to develop SURPH survival estimates to LGR. The SURPH model uses PIT tag detections at various dams to develop a Cormack/Jolly-Seber estimate of survival to LGR. Total natural origin and hatchery origin smolt out-migration (for each lake and different release strategy) to LGR was estimated using the SURPH survival estimate (for each respective release group) multiplied by the out-migration estimate of each release group at the out-migration trap sites found in the Stanley basin (Table 14).

ADULT SPAWNING MONITORING AND EVALUATION

Sockeye Salmon Spawning Investigations

On September 8, 2007, 494 adult sockeye salmon were released to Redfish Lake (Table 16). The hatchery origin adults were reared at NOAA's Manchester Marine Laboratory and NOAA's Burley Creek Hatchery. Sex was determined by ultrasound, and efforts were made to release fish of equal sex ratios. No anadromous adults were released in 2007.

In order to assist in identifying spawning locations, three male and three female hatchery-produced sockeye salmon were fitted with gastric implant radio transmitters prior to release. Telemetry investigations of adult locations began September 17, 2007 and continued weekly through November 7, 2007. Fish locations were recorded weekly by boat tracking.

Redd counts were also conducted once a week (coinciding with radio telemetry events) beginning on October 29, 2007 and continuing until November 7, 2007. Suspected redds were generally enumerated by two observers on each count date. On the final count date, two observers were used. Areas of excavation (possible redds) were generally 3 m x 3 m in size and likely represented spawning events by multiple parents. As such, we do not know how many parents contributed to the production of natural progeny in spawn year 2007. During redd count surveys, any observed carcasses that could be retrieved were collected to facilitate the collection of biological information (e.g., fish sex and spawning status). We also performed redd counts on Little Redfish Lake because two anadromous fish were observed there in 2004.

Unmarked Juvenile Out-migrant Monitoring

In 2007, unmarked out-migrants produced from program fish releases to Redfish Lake included age-1 out-migrants produced from 464 hatchery-origin adults released for volitional spawning in 2006 and age-2 out-migrants produced from 173 hatchery-origin adults released in 2005. No releases occurred in Alturas Lake that would have produced age-1 or age-2 unmarked out-migrants in 2007. Any age-1 unmarked out-migrants would have been produced by residual sockeye or kokanee found within Alturas Lake. Unmarked out-migrants produced from program fish releases to Pettit Lake included age-2 out-migrants produced from 51,239 eyed-eggs released in 2005. The proportions of age-1 and age-2 unmarked out-migration were determined for 2007 Redfish, Alturas, and Pettit Lake out-migrants by aging scales. The methods are discussed for this analysis above.

Adult Trapping on Redfish Lake Creek

The adult weir on Redfish Lake Creek captures all upstream migrating sockeye salmon and bull trout. Trapping in Redfish Lake Creek for adult migrants started on July 7 and continued until September 22, 2007. During the operation of the adult sockeye salmon weir on

Redfish Lake Creek in 2007, one adult anadromous sockeye and twenty-six adult bull trout were captured.

RESULTS AND DISCUSSION

SOCKEYE SALMON JUVENILE OUT-MIGRANT MONITORING AND EVALUATION

Redfish Lake Creek Trap

A total of 4,535 sockeye salmon smolts (1,331 natural origin and 3,204 hatchery origin) were trapped during the 2007 out-migration season (Figure 3). Fork length of natural origin and hatchery origin sockeye salmon smolts captured averaged 114 mm (range 85 mm to 168 mm; Figure 4) and 100 mm (range 58 mm to 126 mm; Figure 5), respectively. Sockeye salmon smolt lengths were slightly longer in 2007 than during the 2006 out-migration season (Plaster et al. 2007).

Based on observed trapping efficiencies and discharge during out-migration monitoring, we determined that using a single season trapping interval for the natural origin smolts was appropriate for estimating total natural origin sockeye salmon smolt out-migration. Of the 1,331 natural origin smolts handled in 2007, 462 were marked and released upstream of the weir to estimate trapping efficiency.

Trap efficiency decreased from a five-year average (2001-2005) of 42% to 24% during 2007. This was likely due to predation at the trap by small mammals, piscivorous birds, and bull trout and probably not due to the high flows observed. Several personnel visually observed hooded mergansers and bull trout harassing juvenile sockeye at the trap. The artificial decrease in trap efficiency would have caused us to overestimate the number of natural out-migrants. During 2007, we began a capture and haul program to remove bull trout from the juvenile trap site so we can develop better out-migration estimates at the Redfish Lake Creek trap.

The 2007 total natural origin sockeye smolt out-migration was estimated at 5,280 fish (95% CI 4,504—6,352, Table 14). The proportion of age-1 natural origin smolts was estimated at 90.5%, which equals 4,778 smolts; the proportion of age-2 natural origin smolts was estimated at 9.5%, which equals 502 smolts. As referenced in the Methods section of this report, age proportions were estimated using the MIX software application. The above proportions were similar to out-migration proportions we saw last year (Peterson et al. 2007).

Of the 3,204 fall direct-released smolts handled in 2007, 613 were marked and released upstream of the weir to estimate trap efficiency. Total fall direct-released smolt out-migration was estimated at 14,256 fish (95% CI 12,215—16,679 Table 14). Overwinter survival and out-migration for this group was 23% of the number of presmolts planted in 2006 (Table 15). The proportion of age-1, adipose fin-clipped, hatchery-reared smolts was estimated, using the MIX software application, at 99.2% (95% CI $\pm$ 0.7%), which equals 14,142 smolts; the proportion of age-2, adipose fin-clipped, hatchery-reared smolts was estimated at 0.8% (95% CI $\pm$ 0.7%) which equals 114 smolts. As mentioned above, these proportions are similar to those observed from past years.

Alturas Lake Creek Trap

One-thousand six hundred thirteen sockeye salmon smolts (860 natural origin and 753 hatchery origin) were trapped during the 2007 out-migration season. Numbers trapped and handled were probably affected by the spring runoff conditions, as they are slightly higher than the previous year (Peterson et al. 2007). Fork length of natural origin and hatchery origin sockeye salmon smolts captured averaged 81 mm (range 68 mm to 129 mm) and 90 mm (range 76 mm to 134 mm), respectively. The average size of out-migrating sockeye smolts was larger than in 2006 (Peterson et al. 2007).

Total natural origin sockeye smolt out-migration for Alturas Lake was estimated at 8,994 fish (95% CI 6,985—12,064 see SBT annual report to BPA for methods and calculations) and hatchery origin smolt out-migration was estimated at 6,897 fish (95% CI 5,487—8,620 see SBT annual report for methods and calculations) (Table 14). Genetic samples were collected from out-migrating sockeye smolts and could possibly be used to identify the source of out-migrants (whether these fish were produced by kokanee or residual sockeye). Overwinter survival and estimated out-migration for the fall direct-release group was 25.6% of the number of presmolts planted in 2006. Scales for aging were collected from natural origin smolts at the Alturas Lake Creek trap in 2007. The proportion of age-1, natural origin smolts was estimated, using the MIX software application, at 95.8% (95% CI 93.3%-98.3%), which equals 8,616 smolts; the proportion of age-2, natural origin smolts was estimated at 4.2% (95% CI 1.7%-6.7%), which equals 378 smolts. We will continue to collect scales from these fish and monitor proportions for this population.

Pettit Lake Creek Trap

Total natural origin sockeye smolt out-migration for Pettit Lake in 2007 was estimated at 1,749 fish (95% CI not calculated; see SBT annual report to BPA for methods and calculations) and hatchery origin smolt out-migration was estimated at 4,695 fish (95% CI not calculated; see SBT annual report for methods and calculations) (Table 14). Overwinter survival and out-migration for fall 2006 direct-released presmolts was 25.4%. Fork length of natural origin and fall direct-released sockeye salmon smolts captured averaged 139 mm (range 95 mm to 189 mm) and 106 mm (range 80 mm to 135 mm), respectively. Scales for aging were collected from natural origin smolts at the Pettit Lake Creek trap in 2007. The proportion of age-1, natural origin smolts was estimated, using the MIX software application, at 11.2% (95% CI 6.8%-15.6%), which equals 196 smolts; the proportion of age-2, natural origin smolts was estimated at 88.8% (95% CI 84.4%-93.2%), which equals 1,553 smolts. Age class proportions were similar for Pettit Lake as seen in 2005 and 2006 (Peterson et al. 2007).

Salmon River and Redfish Lake Creek Smolt Groups

We released 47,094 BY05 smolts to the Salmon River above the Sawtooth FH river water intake upstream of the weir on May 8, 2007. This release group consisted of smolts reared at Sawtooth FH (46,765) and Eagle FH (329). All fish released were adipose fin-clipped and coded-wire tagged with a representative group (n = 1,354) PIT tagged. Fork length of the Salmon River smolt group averaged 125.6 mm (range 98 mm to 166 mm), and these fish had an average weight of 21.7 g/fish.

A total of 54,582 BY05 smolts were released into Redfish Lake Creek below the out-migrant weir on May 8, 2007. This release included smolts reared at Oregon Department of Fish

and Wildlife Oxbow FH. All fish were adipose/right ventral fin-clipped and coded-wire tagged with a representative group (n = 1019) PIT tagged. The average length and weight of the Redfish Lake Creek smolt group was 158.4 mm (range 115 mm to 195 mm) and 51.6 g/fish (range 15.9 to 79.8 g/fish), respectively. Oxbow FH smolts are larger than Sawtooth FH smolts because of the warmer rearing water temperatures at Oxbow FH (spring water) and temperatures vary throughout the year at Sawtooth FH (Salmon River water).

Mainstem Snake and Columbia River Dams

We estimated smolt out-migration success to LGR for natural origin and hatchery origin sockeye salmon smolt groups using PIT tag interrogation data (Table 14; Appendix B). Estimates reflect numbers of smolts that arrived at LGR based on results from data analyses using the SURPH model (Table 14). Survival among release groups ranged from 31% (Redfish Lake natural origin smolts) to 78% (hatchery origin Salmon River smolt release and Alturas Lake fall direct release presmolt groups). The average survival rate for all the release groups was 55% to LGR (down from 59% survival in 2006). An estimated 1,637 (31% survival) and 5,275 (37% survival) natural origin and fall direct-release sockeye smolts, respectively, survived to LGR from the Redfish Lake Creek trap. Survival from the Alturas Lake Creek trap to LGR was estimated at 4,137 (46% survival) for natural origin smolts and 5,380 (78% survival) for fall direct-release smolts. Survival from the Pettit Lake Creek trap to LGR was estimated at 997 (57% survival) for natural origin smolts and 2,582 (55% survival) for fall direct-release smolts. An estimated 36,733 (78% survival) sockeye smolts survived to LGR from the Salmon River smolt release group and 31,658 (58% survival) sockeye smolts survived from the Redfish Lake Creek smolt release. The total estimate for sockeye smolts that out-migrated from Stanley Basin waters and survived to LGR for 2007 was 88,399 (Table 14).

Since the initiation of the smolt release program we have seen survival increases within each release group. We have also identified that sockeye smolts appear to survive slightly better than chinook smolts released at the same location within the valley. The increase in survival as well as the appearance of better survival for sockeye over chinook smolts will be investigated further during the upcoming year.

This was the second year we used the SURPH model to estimate survival and numbers of smolts to LGR. In the past, we had used daily collection efficiencies estimated for Chinook salmon smolts (Sandford and Smith 2002) and expanded PIT tag interrogation data to develop these estimates. The old methodology did not allow us to develop confidence intervals around the estimates, and annual out-migration estimates were generally not comparable due to variability in systems operations. We will continue to utilize the SURPH model to develop survival and total estimates to LGR and will run past program data through the model to enable us to make comparisons between years.

SOCKEYE SALMON SPAWNING INVESTIGATIONS AND UNMARKED JUVENILE OUT-MIGRANT MONITORING

Sockeye Salmon Spawning Investigations

The first redd was observed near the transfer camp at the south end of Redfish Lake on October 17, 2007. Redd counts were finalized with two observers in a boat on November 7, 2007. This was the first year that we observed adults utilizing the area near Point Campground

as a spawning location since the initiation of the project. Redd construction timing was similar to observations made in the past (Willard et al. 2006, Plaster et al. 2007). During the final counts, 195 redds (areas of excavation) were identified (Table 17). Eighty-four redds were located near the Transfer Camp, 58 redds were located at the south beach area, 42 were located at Sockeye Beach, and 11 were counted on the West side of Redfish Lake near Point Campground (Table 17; Figure 6). This is the highest redd count since the project began and is likely the result of the largest adult release to date. Areas of excavation (possible redds) are typically large (~3 m x 3 m) and may represent multiple spawning events by multiple parents; therefore, we do not know how many parents contributed to potential natural production in 2007. Radio-tagged fish moved to areas where redds were observed; however, we were unable to collect any adult carcasses in 2007 to verify these fish had spawned.

No completed redds or areas of excavation were located in Little Redfish Lake in 2007.

Unmarked Juvenile Out-migrant Monitoring

In 2007, 5,280 unmarked smolts (95% CI 4,504 to 6,352) were estimated to have out-migrated from Redfish Lake (Table 13), 8,994 unmarked smolts (95% CI 6,985 to 12,064) were estimated to have out-migrated from Alturas Lake, and 1,749 unmarked smolts were enumerated to have out-migrated from Pettit Lake (Table 14, Figure 7). The SBSTOC believes it is very important to utilize reintroduction strategies that produce natural origin smolts (e.g., prespawn adult releases and eyed-egg introductions). It is very difficult to evaluate the effectiveness of these release strategies while we have kokanee populations and potentially increasing residual populations within the basin lakes.

Since 2002, we have been working with the University of Idaho's Center for Salmonid and Freshwater Species at Risk and the Eagle Fish Genetics Laboratory to utilize DNA microsatellite methods to identify individual parental contribution to unmarked smolt production through parental exclusion analysis. Parental exclusion analysis allows assignment of an individual smolt to parents of a release strategy, allowing comparisons between the relative individual contribution of adults released to spawn volitionally and survival comparisons to certain life history stages (e.g., green-egg to smolt and/or fry to smolt survival) for various release strategies. Results would allow program managers to emphasize the release strategy with the highest reproduction potential. Additionally, parental exclusion analysis will allow for evaluation of the reproductive contribution of residuals and estimation of the number of kokanee that fall out. We are currently testing the assignments of juveniles from known adult crosses to see how well they assign and then we should be able to employ these techniques to samples collected in the field (Christine Kozfkay, personal communication).

Table 13. Out-migration estimate for natural origin sockeye salmon smolts captured at the Redfish Lake Creek trap from April 14 to June 13, 2007.

Natural Origin Smolts (single interval estimate)	
	Total
Dates	4/14-6/13
Trap efficiency	0.27
Marked	462
Recaptured	116
Total handled	1,331
Estimated total (single interval estimate)	5,280
95% CI upper bound	6,352
95% CI lower bound	4,504
Hatchery Origin Smolts (single interval estimate)	
	Total
Dates	4/14-6/13
Trap efficiency	0.22
Marked	613
Recaptured	137
Total handled	3,204
Estimated total (single interval estimate)	14,256
95% CI upper bound	16,679
95% CI lower bound	12,215

Table 14. Summary of 2007 sockeye salmon smolt out-migration information (by release strategy) at trap locations and at Lower Granite Dam (LGR). Sawtooth Fish Hatchery (SFH) was the rearing location for the fall-direct released (FDR) presmolts and Salmon River smolt release group. Oxbow Fish Hatchery (OFH) was the rearing location for the Redfish Lake Creek smolt release.

Release strategy (rearing location)	Total released ^a	Number tagged prior to release	PIT tags detected at trap	Smolt out- migration estimate	Number tagged at trap	Estimated SURPH survival at LGR	SURPH 95% CI (±)	Estimated no. at LGR
<u>Redfish Lake</u>								
Natural origin smolt	NA	NA	NA	5,280	462	31.00%	7.80%	1,637
FDR presmolt (SFH)	61,804	1,021	53	14,256	613	37.00%	7.80%	5,275
<u>Alturas Lake^b</u>								
Natural origin smolt	NA	NA	NA	8,994	592	46.00%	9.80%	4,137
FDR presmolt (SFH)	26,994	1,016	23	6,897	195	78.00%	33.00%	5,380
<u>Pettit Lake^b</u>								
Natural origin smolt	NA	NA	NA	1,749	198	57.00%	99.90%	997
FDR presmolt (SFH)	18,494	1,021	256	4,695	252	55.00%	17.60%	2,582
<u>Salmon River</u>								
Hatchery origin smolt (SFH)	47,094	1,354	NA	47,094	NA	78.00%	27.40%	36,733
<u>Redfish Lake Creek</u>								
Hatchery origin smolt (OFH)	54,582	1,019	NA	54,582	NA	58.00%	15.70%	31,658

^a Total released from hatchery.

^b Data from Alturas and Pettit lake trap obtained from Shoshone-Bannock Tribes biologists.

^c 95% CIs are two standard errors.

Table 15. Estimated overwinter survival until out-migration for Sawtooth Fish Hatchery-reared presmolts released in the fall to Redfish, Alturas, and Pettit lakes.

<u>Out-migration Year</u>	<u>Redfish Lake</u>	<u>Alturas Lake</u>	<u>Pettit Lake</u>
2000	29%	34%	46%
2001	20%	75%	29%
2002	40%	30%	29%
2003	15%	NA	59%
2004	27%	54%	35%
2005	35%	82%	56%
2006	43%	38%	64%
2007	23%	26%	25%

Table 16. Redfish Lake Sockeye Salmon Captive Broodstock Program prespawn adult release history.

<u>Lake</u>	<u>Rearing origin</u>	<u>Date released</u>	<u>Number released</u>	<u>Number of suspected redds</u>
Redfish	Full-term hatchery	1993	20	Unknown
Redfish	Full-term hatchery	1994	65	One behavioral observation
Redfish	Full-term hatchery	1996	120	30 suspected redds
Redfish	Full-term hatchery	1997	80	30 suspected redds
Pettit	Full-term hatchery	1997	20	1 suspected redd
Alturas	Full-term hatchery	1997	20	Test digs only
Redfish	Full-term hatchery	1999	18	
Redfish	Hatchery-produced anadromous	1999	3	8 suspected redds
Redfish	Full-term hatchery	2000	36	
Redfish	Hatchery-produced anadromous	2000	120	20 to 30 suspected redds
Pettit	Hatchery-produced anadromous	2000	28	none confirmed
Alturas	Full-term hatchery	2000	25	
Alturas	Hatchery-produced anadromous	2000	52	14 to 19 suspected redds
Redfish	Hatchery-produced anadromous	2001	14	12 to 15 suspected redds
Redfish	Full-term hatchery	2001	65	
Redfish	Hatchery-produced anadromous	2002	12	10 to 12 suspected redds
Redfish	Full-term hatchery	2002	178	
Redfish	Full-term hatchery	2003	312	42 suspected redds
Redfish	Full-term hatchery	2004	241	127 suspected redds
Redfish	Full-term hatchery	2005	173	78 suspected redds
Redfish	Full-term hatchery	2006	464	172 suspected redds
Redfish	Full-term hatchery	2007	494	195 suspected redds
Total			2,560	

Table 17. Summary of sockeye salmon redd observations in Redfish Lake in 2007.

	Sockeye Beach	South Beach	Transfer Camp	West Side	Total
Final Count on 11/7/2007	42	58	84	11	195

Figure 3. Daily capture of natural origin and hatchery origin sockeye salmon smolts (unexpanded) at the Redfish Lake Creek trap during the 2007 out-migration.

Figure 4. Length frequency of natural origin sockeye salmon smolts (n = 491) collected at Redfish Lake Creek trap in 2007.

Figure 5. Length frequency of fall-direct released sockeye salmon smolts (n = 609) collected at Redfish Lake Creek trap in 2007.

Figure 7. Natural origin sockeye salmon smolt out-migration estimated at Redfish Lake Creek, Alturas Lake Creek, and Pettit Lake Creek traps from 1991 to 2007 (juvenile out-migrant traps on Pettit Lake Creek were not operated every year).

LITERATURE CITED

- Bjornn, T. C., D. R. Craddock, and D. R. Corley. 1968. Migration and survival of Redfish Lake, Idaho sockeye salmon *Oncorhynchus nerka*. Transactions of the American Fisheries Society 97:360-373.
- Craddock, D. R. 1958. Construction of a two-way weir for the enumeration of salmon migrants. The Progressive Fish-Culturist 20:33-37.
- Hebdon, J. L., M. Elmer, and P. Kline. 2000. Snake River sockeye salmon captive broodstock program, research element, 1999. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Hebdon, J. L., J. Castillo, and P. Kline. 2002. Snake River sockeye salmon captive broodstock program, research element, 2000. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Hebdon, J. L., J. Castillo, C. Willard, and P. Kline. 2003. Snake River sockeye salmon captive broodstock program, research element, 2001. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P. 1994. Research and recovery of Snake River sockeye salmon, 1993. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P., and J. Younk. 1995. Research and recovery of Snake River sockeye salmon, 1994. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Kline, P., and J. A. Lamansky. 1997. Research and recovery of Snake River sockeye salmon, 1995. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- MacDonald, P. D. M., and P. E. J. Green. 1988. User's Guide to Program MIX: an interactive program for fitting mixtures of distributions. *Release 2.3, January 1988*. Ichthus Data Systems, Hamilton, Ontario. iv+60 pp. ISBN 0-9692305-1-6.
- Plaster, K., M. Peterson, D. Baker, J. Heindel, J. Redding, C. Willard and P. Kline. 2007. Snake River sockeye salmon captive broodstock program, research element 2005. IDFG Report no. 06-36. Project no. 199107200. Bonneville Power Administration, Annual Report. Portland, Oregon.
- Sandford, B. P., and S. G. Smith. 2002. Estimation of smolt-to-adult return percentages for Snake River basin anadromous salmonids, 1990-1997. Journal of Agricultural, Biological and Environmental Statistics.
- Steinhorst, K., Y. Wu, B. Dennis, and P. Kline. 2004. Confidence intervals for fish out-migration estimates using stratified trap efficiency methods. Journal of Agricultural, Biological, and Environmental Statistics 9:284-299.

- Willard, C., J. L. Hebdon, J. Castillo, J. Gable, and P. Kline. 2004. Snake River sockeye salmon captive broodstock program, research element, 2002. Report to Bonneville Power Administration, Contract 5342.
- Willard, C., K. Plaster, J. Castillo, and P. Kline. 2005. Snake River sockeye salmon captive broodstock program, research element 2003. Report to Bonneville Power Administration, Contract 5342.

PART 4—PREDATOR SURVEYS

INTRODUCTION

Declines in bull trout populations throughout the Pacific Northwest led to their listing as threatened under the Endangered Species Act in 1998. Prior to listing, IDFG implemented no-harvest fishing regulations to help protect the remaining populations in the State of Idaho. Because bull trout readily consume kokanee and other salmonids (Bjornn 1961; Beauchamp and Van Tassell 2001), a large increase in the number of adult bull trout in Stanley Basin lakes could affect the sockeye salmon and kokanee populations in the lakes.

Bull trout spawner investigations were initiated in 1995 to track population response to no-harvest fishing regulations implemented by IDFG in 1994. Trend data of this nature have been successfully used to measure population response to fishing regulation changes implemented for adfluvial bull trout populations in Oregon and British Columbia (Ratliff 1992; Stelfox and Egan 1995). Index sections were established on Fishhook and Alpine creeks in 1998. Information collected in 2007 represented the tenth year data were collected in these index reaches. Bull trout generally mature at five to six years of age (Leary et al. 1993); therefore, we would expect to see effects of no-harvest regulations become apparent in surveys conducted after 2000 as these cohorts represent those protected from potential harvest by anglers.

METHODS

Monitoring of spawning bull trout was initiated in 1995 to measure bull trout population response to no-harvest fishing regulations implemented by IDFG in 1994. In 2007, we surveyed index reaches, which were established in 1998, on principal tributary streams of Redfish and Alturas lakes to enumerate bull trout spawners and redds. Surveys were conducted on Fishhook Creek (Redfish Lake drainage) and Alpine Creek (Alturas Lake drainage) August 28 and 29 and on September 12 and 13, 2007. These dates typically correspond with the initiation of spawning (first survey, we are typically able to count adult spawners) and the completion of spawning activities (second survey, we typically develop final redd count numbers). No suitable tributary streams feed Pettit Lake and, as such, bull trout spawner surveys were not conducted on this system. Index sections were established with global positioning satellite (GPS) equipment. Two observers walked from the lower boundary of the index section upstream and recorded visual observations of bull trout and known or suspected bull trout redds. Coordinates of redd locations were recorded with a handheld GPS unit. In order to avoid omission of completed redds during the final count, completed redds identified during the first count were flagged. Flagging prevented omitting redds from the final count that were obscured over time.

In 2007, an additional area was surveyed for bull trout redds in Fishhook Creek. The new section includes the lower portion of Fishhook Creek upstream of the first gradient gain above Redfish Lake and ends at the wilderness boundary (between GPS waypoints 44° 08'36.37"N 114°57'29.97"W, and 44°08'53.69" N 114°55'39.13"W).

Adult Bull Trout Trapping Redfish Lake Creek

The adult weir on Redfish Lake Creek captures all upstream migrating sockeye salmon and bull trout. Trapping in Redfish Lake Creek for adult migrants started on July 7 and

continued until September 22, 2007. During the operation of the adult sockeye salmon weir on Redfish Lake Creek in 2007, one adult anadromous sockeye salmon and twenty-six adult bull trout were captured. Bull trout that were captured and passed received PIT tags ($n = 13$). Thirteen bull trout mortalities were recorded in 2007 (five associated with trapping, eight attributed to either angling or post spawn events).

RESULTS AND DISCUSSION

Fishhook Creek

We observed 41 adult bull trout (23 males, 18 females) and 21 developing redds on August 29, 2007. During our second survey on September 13, we observed two adult bull trout and 22 complete redds (Figure 8). Redd counts on the second date included developing redds observed on the previous count (Table 18).

From the ten years of data, it appears that the no-harvest fishing regulation is positively affecting the bull trout population in Fishhook Creek. Redd counts in Fishhook Creek had been holding constant between 1998 and 2004, but recent increases in redds counted (varying between 11 and 30 redds counted each year) shows a dramatic increase in bull trout reproduction. Because bull trout may spawn in alternating or consecutive years (Fraley and Shepard 1989), year-to-year variation would be expected. Our findings are consistent with results from statewide monitoring efforts, which indicate that bull trout are increasing or at least stable across most of their range in Idaho (High et al. 2005).

Alpine Creek

We observed 15 adult bull trout (seven males, six females, and two of unknown gender) and 17 developing redds during the August 28, 2007 survey. On September 12, we observed three adult male bull trout and 18 completed redds (Figure 9). Redd counts on the second date included developing redds and completed redds observed during the previous count (Table 19). The Alpine Creek population has increased steadily since 1998; one redd was observed in 1998, and the number of redds has ranged from nine to 18 between 2000 and 2007. Monitoring bull trout populations with redd counts is advantageous, because they are low cost and cause little disturbance to spawning bull trout. Documentation of significant population changes have been identified from redd count data (Rieman and Meyers 1997).

We believe that our counts of redds in the trend sections were an accurate reflection of the numbers of redds present. The streams in our surveys were much smaller than those used by Dunham et al. (2001) which indicated that redds could be missed in larger systems. For example, in the systems studied by Dunham et al. (2001), deepwater cover was defined as water greater than 1 m deep. In Fishhook and Alpine creeks, water depth rarely approached 1 m deep.

Bull Trout Capture, Mark, and Haul Operations Redfish Lake Creek

As mentioned previously in this report, during the out-migration season of 2007 trap efficiency at the juvenile out-migrant sockeye salmon trap on Redfish Lake Creek dropped from a five-year season average of 41% to 25%. The marked decrease in trap efficiency was attributed to avian, mammalian, and fish predation observed by trap tenders. Mammalian

predation was controlled (to some extent) by setting traps along the fish trap location and lethal removal of mink occurred as they were captured ($n = 2$). Fish predation was attributed primarily to bull trout keying in on the release of marked sockeye salmon used to measure the trap efficiency. In an attempt to develop accurate juvenile sockeye out-migration estimates with reliable trap efficiencies, IDFG collected, PIT tagged, and transported bull trout away from the juvenile sockeye trap located on Redfish Lake Creek.

Thirty adult bull trout were captured at the Redfish Lake Creek trap by hook and line, PIT tagged, and transported to a release site above Little Redfish Lake (Table 20). Six bull trout were subsequently recaptured at the trap site following their release. Six bull trout were found to have PIT tags from *O. nerka* tagging files detected in their stomachs (tag codes read through abdominal wall of adult bull trout).

Work done by Schoby (2006) suggests that bull trout in the upper Salmon River migrate from spawning tributaries in early October and that some individuals migrate to Redfish Lake to overwinter. Over the past few years, numbers of bull trout captured at the East Fork Salmon River weir have increased (from 169 bull trout in 2004 to 227 bull trout in 2007, Vidergar, personal communication). Due to the appearance of an expanding bull trout population on the East Fork of the Salmon River and the fact that most of these bull trout are handled at the weir, we are planning to continue PIT tagging bull trout at the East Fork Salmon River (EFSR) weir (Sawtooth FH satellite weir) to determine if bull trout from this system migrate upriver to Redfish Lake to overwinter.

The collection of these data should provide us with a more complete picture of how bull trout populations have responded to “no harvest regulations” in the upper Salmon River drainage and whether we have predation issues within Redfish Lake that are not accounted for using our redd count trend sites. If increasing numbers of bull trout are migrating into the system to overwinter, we may see predation on *O. nerka* increase.

Table 18. Bull trout adult fish counts and redd counts in index sections of Fishhook Creek from 1998 to 2007.

Year	Dates	Number of bull trout observed	Number of redds
1998	8/22	40	5
	9/10	2	11
1999	8/22	40	0
	8/26	33	15
2000	8/31	16	12
	9/14	2	18
2001	8/28	31	15
	9/11	3	11
2002	9/04	23	6
	9/11	5	17
2003	8/27	40	6
	9/08	15	17
2004	8/30	31	7
	9/9	8	11
2005	8/30	24	12
	9/12	2	23
2006	8/29	32	16
	9/13	0	25
2007	8/29	41	21
	9/13	2	22

Table 19. Bull trout adult fish counts and redd counts in index sections of Alpine Creek from 1998 to 2007.

Year	Dates	Number of bull trout observed	Number of redds
1998	8/23	6	0
	9/11	6	1
1999 ^a	8/26	13	3
	8/30	18	6
2000	9/15	5	9
	8/28	8	11
2001	9/11	1	15
	8/30	20	8
2002	9/12	0	14
	8/27	27	11
2003	9/08	0	14
	8/31	16	6
2004	9/9	0	9
	8/30	16	9
2005	9/12	0	13
	8/28	15	6
2006	9/12	1	13
	8/28	15	17
2007	9/12	3	18

^a Only one count completed.

Table 20. Bull trout relocation effort juvenile out-migrant trapping at Redfish Lake Creek, 2007.

Year	Bull trout captured, PIT tagged, relocated	Bull trout recaptured at trap site	Bull trout with <i>O. nerka</i> PIT tags detected in stomach	Bull trout mortalities associated with transport activities
2007	30	6	6	1

Figure 8. Location bull trout redd index sections in Fishhook Creek in 2007.

Figure 9. Location of bull trout redd index sections in Alpine Creek in 2007.

LITERATURE CITED

- Beauchamp, D. A., and J. J. Van Tassell. 2001. Modeling seasonal trophic Interactions of adfluvial bull trout in Lake Billy Chinook, Oregon. Transactions of the American Fisheries Society 130:204-216.
- Bjornn, T. C. 1961. Harvest, age, and growth of game fish populations from Priest to Upper Priest Lakes. Transactions of the American Fisheries Society 118:597-607.
- Dunham, J., B. Rieman, and K. Davis. 2001. Sources and magnitude of sampling error in redd counts for bull trout. North American Journal of Fish Management 21:343-352.
- Fraley, J. J., and B. Shepard. 1989. Life history, ecology and population status of migratory bull trout *Salvelinus confluentus* in the Flathead lake and river systems, Montana. Northwest Science 63:133-142.
- High, B., K. A. Meyer, D. J. Schill, and E. R. J. Mamer. 2005. Wild trout investigations. Job performance report 2004. Grant F-73-R-26. Report no. 05-24. Idaho Department of Fish and Game, Boise, Idaho.
- Leary, R. F., F. W. Allendorf, S. H. Forbes. 1993. Conservation genetics of bull trout in Columbia and Klamath River drainages. Conservation Biology 7:856-865.
- Ratliff, D. E. 1992. Bull trout investigations in the Metolius River-Lake Billy Chinook system. *In* Howell, P. J. and D. V. Buchanan, editors. Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society. Corvallis, Oregon.
- Rieman, B. E., and D. L. Meyers. 1997. Use of redd counts to detect trends in bull trout *Salvelinus confluentus* populations. Conservation Biology 11:1015-1018.
- Schoby, G. 2006. Home range analysis of bull trout (*Salvelinus confluentus*) and westslope cutthroat trout (*Oncorhynchus clarki lewisi*) in the Upper Salmon River Basin, Idaho. Master's Thesis. Idaho State University.
- Stelfox, J. D., and K. L. Egan. 1995. Bull trout investigations in the Smith-Dorien Creek/Lower Kananaskis Lake system. Alberta Environmental Protection, Fisheries Management Division, and Golder Associates Limited, Calgary, Canada.

ACKNOWLEDGEMENTS

Special thanks to the Eagle and Sawtooth fish hatchery staffs for assisting with coordination and field activities. Special thanks also to Cheryl Leben for formatting the final document and Debbie Myers for creating the GIS maps. We would like to thank the three reviewers: Jeff Heindel, Carl Stiefel, and Chris Kozfkay.

APPENDICES

Appendix A. Fork length, weight (g) and age of *O. nerka* captured during midwater trawls conducted during September 2007 on Redfish, Pettit, and Alturas lakes.

Transect	Length (mm)	Weight (g)	Age
<u>Redfish Lake</u>			
1	38	0.4	0
1	47	0.9	0
1	54	1.6	0
2	45	0.6	0
2	48	0.9	0
2	52	1.2	0
2	53	1.3	0
2	55	1.4	0
2	58	2.0	0
2	59	2.0	0
2	60	1.9	0
2	60	1.8	0
2	176	66.9	2
2	190	76.5	2
3	35	0.4	0
3	42	0.8	0
3	50	1.0	0
3	53	1.2	0
3	58	1.6	0
3	61	2.1	0
3	63	1.7	0
3	65	2.8	0
3	78	4.5	0
3	80	5.2	1
4	47	0.9	0
4	48	1.0	0
4	53	1.2	0
4	63	2.6	0
4	63	2.7	0
4	130	21.5	1
5	44	0.7	0
5	47	1.0	0
5	53	1.4	0
5	53	1.3	0
5	56	1.7	0
5	58	1.4	0
5	74	4.5	0
5	83	5.3	1
5	91	6.9	1
6	43	0.8	0
6	48	1.2	0
6	50	1.0	0
6	53	1.2	0
6	57	1.4	0
6	57	1.6	0
6	65	2.6	0
6	66	3.5	0
6	70	2.7	0
6	75	4.0	0
6	77	5.5	0
6	102	10.8	1
6	135	30.1	1
6	185	69.6	2

Appendix A. Continued.

Transect	Length (mm)	Weight (g)	Age
<u>Alturas Lake</u>			
1	48	0.88	0
1	51	1.15	0
1	70	3	1
1	70	2	1
1	70	3	1
1	72	3	1
1	72	3	1
1	73	2	1
1	73	3	1
1	73	3	1
1	75	3	1
1	76	4	1
1	83	4	1
1	84	5	1
1	85	5	1
1	86	5	1
1	93	7	1
1	95	7	1
1	95	8	1
1	98	8	1
1	100	8	1
1	107	13	1
1	108	14	1
1	127	22	1
1	171	54	2
1	194	73	3
2	53	1.31	0
2	57	1	0
2	58	1.58	0
2	59	1	0
2	60	1	0
2	60	1	0
2	61	1	1
2	68	2	1
2	69	2	1
2	70	3	1
2	75	4	1
2	76	3	1
2	77	2	1
2	78	4	1
2	80	4	1
2	83	4	1
2	83	4	1
2	90	6	1
2	93	8	1
2	93	7	1
2	95	7	1
2	96	7	1
2	98	8	1
2	98	8	1
2	116	17	1
2	119	17	1
2	125	23	1
2	130	27	2
2	130	24	2

Appendix A. Continued.

Transect	Length (mm)	Weight (g)	Age
2	153	40	2
2	164	49	2
2	166	25	2
2	182	57	3
3	43	0.71	0
3	45	0.68	0
3	45	1.08	0
3	53	1.29	0
3	55	0.77	0
3	55	1.36	0
3	56	1	0
3	65	2	1
3	70	3	1
3	72	3	1
3	75	3	1
3	76	4	1
3	80	5	1
3	83	4	1
3	90	6	1
3	140	30	2
3	153	39	2
3	160	43	2
3	162	44	2
3	173	60	2
3	203	82	3
4	41	.65	0
4	42	1.00	0
4	43	.82	0
4	43	.94	0
4	44	.84	0
4	45	.78	0
4	45	.77	0
4	46	.73	0
4	46	.82	0
4	47	.71	0
4	48	.53	0
4	48	.82	0
4	50	.85	0
4	50	1.34	0
4	52	1.11	0
4	56	1	0
4	58	1	0
4	58	1	0
4	58	1.14	0
4	58	2	0
4	63	1	1
4	65	3	1
4	68	2	1
4	69	3	1
4	70	4	1
4	72	2	1
4	75	3	1
4	78	4	1
4	83	6	1
4	83	5	1
4	84	6	1
4	88	5	1

Appendix A. Continued.

Transect	Length (mm)	Weight (g)	Age
4	88	5	1
4	89	5	1
4	95	8	1
4	100	10	1
4	106	10	1
5	45	.98	0
5	46	.92	0
5	48	.84	0
5	48	.91	0
5	72	3	1
5	75	3	1
5	76	3	1
5	77	4	1
5	77	3	1
5	80	4	1
5	81	4	1
5	84	4	1
5	85	5	1
5	85	5	1
5	90	7	1
5	90	7	1
5	91	6	1
5	92	7	1
5	92	6	1
5	96	8	1
5	100	10	1
5	105	11	1
5	105	11	1
5	130	19	1
5	150	40	2
5	151	41	2
5	158	43	2
5	170	52	2
5	189	69	3
6	87	6	1
6	91	7	1
6	94	8	1
6	95	8	1
6	96	9	1
6	98	9	1
6	101	11	1
6	110	14	1
6	133	25	2
6	160	51	2
<u>Pettit Lake</u>			
1	160	43.0	2
1	172	61.9	2
1	177	58.9	2
2	120	19.0	1
2	132	28.7	1
2	165	47.3	2
2	165 (hatchery)	48.2	2
3	150	38.1	2
3	155	41.5	2
3	158	43.6	2
4	122	23.3	1

Appendix A. Continued.

Transect	Length (mm)	Weight (g)	Age
4	128	24.8	1
4	135	25.7	1
4	147	39.7	2
4	152	45.7	2
4	193	78.7	2
5	153	41.7	2
5	175	58.5	2
6	132	27.1	1
6	154	41.4	2
6	156	43.6	2

Appendix B. Arrival dates at Lower Granite Dam for PIT-tagged sockeye salmon smolts during the 2007 migration year.

Date	Redfish Lake						Pettit Lake			Alturas Lake		
	Natural Origin	Fall Direct	Oxbow Smolts	Sawtooth Smolts	Eagle Smolts	Oxbow Smolts	Natural Origin	Fall Direct	Hatchery Smolts	Natural Origin	Fall Direct	Hatchery Smolts
4/30/2007	1											
5/1/2007	2									1		
5/2/2007										1		
5/3/2007	3									2		
5/4/2007	2	1	4							11	1	
5/5/2007	4		2							14	2	1
5/6/2007	5		2							5	2	2
5/7/2007	2	1	2							3	1	1
5/8/2007	3		2							2	4	1
5/9/2007	4		5							2	5	1
5/10/2007	2	6	5							2	5	3
5/11/2007	1	2	3						1	3	2	
5/12/2007	1	5	2					2		2	5	3
5/13/2007	4	9	8	6			1		7	2	9	
5/14/2007	3	1	4	23				1	3		2	2
5/15/2007	4		2	18	1	8				1	2	
5/16/2007	2	2	6	9	4	20	2	2	2		1	1
5/17/2007	3	1	1	11	6	27		6	1	1		3
5/18/2007	3	7	2	2	7	28	1	3		1	2	2
5/19/2007		4	2	7	2	18	1	6	2			4
5/20/2007	1	2	8	2	3	5	1		2		3	
5/21/2007		2	1			1		1	2	1	4	3
5/22/2007		1				1	2	2	1			1
5/23/2007		2	2						1			
5/24/2007			2				1	2	4			
5/25/2007		3					1	1	2		1	
5/26/2007								1	1			
5/27/2007							1	1		1		2
5/28/2007		2	1					3	3		3	
5/29/2007								2				
5/30/2007									1			
5/31/2007												1
6/1/2007		2								1		
Total	50	53	66	78	23	108	11	32	33	56	54	31

^a Ad-clipped hatchery smolts captured and tagged at trap locations.

^b Sawtooth smolts released into the main Salmon River upstream of the Sawtooth Fish Hatchery weir.

^c Oxbow smolts released downstream of the Redfish Lake Creek juvenile trap site.

Prepared by:

Mike Peterson
Fisheries Research Biologist

Kurtis Plaster
Senior Fisheries Technician

Laura Redfield
Senior Fisheries Technician

Jeff Heindel
Conservation Hatchery Supervisor

Paul Kline
Principal Fisheries Research Biologist

Approved by:

IDAHO DEPARTMENT OF FISH AND GAME

Edward B. Schriever, Chief
Bureau of Fisheries

Daniel J. Schill
Fisheries Research Manager