UCRL 10
cy 85/A
C.2

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. División, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-10
C.2

**Special Review of
Declassified Reports**

Authorized by USDOE JK Bratton

Unclassified TWX P182206Z May 79

REPORT PROPERLY DECLASSIFIED

AUG 15 1979

Date

Authorized Derivative Classification

Lawrence Radiation Laboratory Library
University of California, Berkeley

~~RESTRICTED~~

UCRL - 10

HEALTH AND BIOLOGY GENERAL
CLASSIFICATION CONTROLLED BY AUTHORITY
OF THE DISTRICT ENGINEER
BY THE DECLASSIFICATION COMMITTEE

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-Eng.-48

PHOSPHORUS TURNOVER AND PHOTOSYNTHESIS

By

Sam Aronoff and Melvin Calvin

20 November 1947

Special Review of Declassified Reports
Authorized by USDOE JK Bratton
Unclassified TWX P182206Z May 79

REPORT PROPERLY DECLASSIFIED

Authorized Derivative Classifications	Date
8-15-79	8-20-79
By	

This document contains restricted data within the meaning of the Atomic Energy Act of 1946 and/or information affecting the national defense of the United States within the meaning of the Espionage Act U.S.C. 31 & 32, as amended. Its transmission or the revelation of its contents in any manner to an unauthorized person is prohibited and may result in severe criminal penalty.

Berkeley, California

UCRL-10
~~RESTRICTED~~
 HEALTH AND BIOLOGY-GENERAL

Standard Distribution

CLASSIFIED BY AUTHORITY
 No. Copies
 BY THE DECLASSIFICATION COMMITTEE

Argonne National Laboratory	10
Atomic Energy Commission, Washington	3
Battelle Memorial Institute	1
Brookhaven National Laboratories	8
Carbide & Carbon Chemicals Corp. (K-25 Area)	2
Carbide & Carbon Chemicals Corp. (Y-12 Area)	2
Clinton Laboratories	8
General Electric Company	4
Hanford Engineer Works	2
Iowa State College	1
Los Alamos	3
Madison Square Area	1
Massachusetts Institute of Technology	1
Monsanto Chemical Company, Dayton	1
National Bureau of Standards	1
Patent Advisor	1
Research Division (for NEPA), Oak Ridge	1
Research Division, Oak Ridge	15
University of Rochester	2
Declassification Procedure	10
J. H. Lawrence	1
J. G. Hamilton	1
R. S. Stone	1
M. Calvin	2
University of California, Radiation Laboratory	3

Total 85

Radiation Laboratory
 Information Division
 University of California
 Berkeley, California

PHOSPHORUS TURNOVER AND PHOTOSYNTHESIS

BY

Sam Aronoff and Melvin Calvin

From the Radiation Laboratory and
Department of Chemistry,
University of California, Berkeley*

20 November 1947

The participation of phosphorus in biological oxidation-reduction reactions of the type found in glycolysis $ADP + PO_4H^- + 3\text{-phosphoglyceraldehyde} + DPN^+ = 3\text{-phosphoglycerate}^- + 2H^+ + DPNH + ATP$ has suggested theories in which similar reactions are proposed for photosynthesis (Lipmann, (6), Ruben (8), Emerson, Stauffer, and Umbreit (2). In these theories the reducing power of photosynthesis is utilized not only for reduction of carbon dioxide but also, by means of coupled oxidations, for the generation of high-energy phosphate bonds, or in the last reference directly for the generation of high-energy phosphate. Since in these theories acyl phosphate is formed from inorganic phosphate, they are amenable to proof without isolation of particular intermediates, by means of radioactive phosphorus. It would be expected that the rate of conversion of inorganic phosphate to organic phosphate would be greater in light than in the dark. We have investigated this possibility under a variety of conditions and are unable to substantiate the theories. The following experiments are reported:

a) with grana

The use of grana, which are capable of accomplishing the photochemical reduction of a variety of substrates, e.g. of quinones (Warburg (9)

(*) This paper is based on work performed under contract No. W-7405-Eng.-48 with the Atomic Energy Commission in connection with the Radiation Laboratory, University of California, Berkeley.

Fan, et.al, (3), Aronoff, (1), permits a test of the theory of Emerson, et.al. It is to be expected that the direct utilization of light should result in the formation of organic phosphate.

b) with leaves

There is some evidence to indicate that the light reaction involving the fixed carbon dioxide occurs within the chloroplast (Frenkel (4)). This suggests that the chloroplasts, isolated from lighted and darkened halves of a leaf will show differences in rate of formation of organic phosphate.

c) with algae

Diverse aspects were investigated; light vs. dark, the presence and absence of carbon dioxide and of oxygen.

Method:

The P^* obtained had a specific activity of approximately $35 \mu\text{C}/\text{g}$, the concentration of the P^* unfortunately varying from sample to sample and commonly being as high as $10 \text{ g}/\text{ml}$. Experimental total counts of P^* were of the order of $10^6 \rightarrow 10^7$, involving aliquots of $0.1 \rightarrow 0.2$ of the P^* solution. The amount of absorption of P^* into e.g. 100 c.m.m. algae during a normal run of ca. 15 minutes is of the order of $0.1 \rightarrow 5\%$ depending upon particular conditions. All experiments with algae were performed with the usual manometric equipment, P^* being tipped in after equilibration. The determination of the activity in the algae was accomplished, after quantitative removal from the reaction vessels, by centrifugation (complete within 2.5', $20-30^\circ \text{C}$), and washing twice with distilled water, and finally being made to volume 2.5 or 10 ml. (depending on activity of P^* or number of cells) with cold $\frac{M}{10} \text{KH}_2\text{PO}_4$. An aliquot of this suspension was pipetted onto a counting plate, made by glueing filter paper onto an aluminum disc with rubber cement. (In preparation

the use of the rubber cement should not be minimized, and pressure should be applied during drying). With our activities and cell numbers, counts on a plate usually were of the order of 100---1000/minute.

The method is based on the observation that two washings of cells with distilled water removes external radio phosphate completely, and further washings do not remove additional phosphate from the cells. The suspension of the cells in $\frac{M}{10}$ $H_2PO_4^-$, for a period of 8-10 minutes, results in a semi-quantitative exchange between the relatively concentrated external inactive phosphate and the intracellular inorganic radio phosphate. Different cultures appeared to require varying pH's, up to 6.5 for quantitative removal of iP^* . Centrifugation and resuspension of the cells in $H_2PO_4^-$ does not cause appreciable additional removal of phosphate. However, periods longer than the 8-10 minutes diminish the apparent organic P^* (oP^*) since there is continual interchange ($oP^* \rightleftharpoons iP^*$) between external P (eP) and iP^* . The extension of the suspension period, e.g. from 10' $\rightarrow$ 30' results in an apparent diminution of oP^* from 80% $\rightarrow$ 65% (see Fig. 1). This is complicated by the $iP \rightarrow oP$ change, but the use of cold solutions diminishes the rate of these processes. Results within a particular experiment are believed to be valid to within 10%.

The total P^* (tP^*) having thus been obtained, the inorganic P^* (iP^*) may be determined from the supernatant after the suspension has equilibrated with the cold $H_2PO_4^-$ for $\sim 10'$. Determination of oP from the cells is preferable, however, since the "geometry" of the counting plate is more comparable to that for the total.

In either theory under consideration, an increased rate of conversion to organic acyl-phosphate compounds is expected. Were it not for considerations of permeability, one could measure the effect merely by determination of total P^* uptake, since this would be increased

under the influence of light. Actually, by far the slower of the two processes is that of the initial entrance of P into the cell. Considerations discussed below will show that although it is desirable to be aware of both rates, a) the rate of entry of external radio-phosphorus (eP^*) into the cell, becoming internal inorganic radio-phosphorus (iP^*), and b) the rate of conversion of iP^* to internal organic radio-phosphorus (oP^*), the critical quantity is the rate of conversion of iP^* to oP^* .

EXPERIMENTS

Grana

Grana were prepared from spinach leaves, purchased locally, via a method similar to that used for the chloroplasts, except for the further centrifugation in a high-speed centrifuge (20,000 g. 10', T = 2.5° C). (see Aronoff (1)).

Almost all of the P^* was removed by washing with KH_2PO_4 . Thus no conversion to oP had occurred, and the equality of P^* uptake indicated no effect of light. The theory of Emerson, et.al. (2) is thus not confirmed. (see Table I),

Leaves

Tobacco leaves, used for chloroplast preparation, were obtained from plants grown in a green house in nutrient solution, during summer months. Chloroplasts were prepared by pressing leaves with a pestle in a mortar containing ice-cold M/20 $KHCO_3$, and differential centrifugation, and washing with distilled water.

An excised tobacco leaf was permitted to absorb P^* in the dark, followed by an equal period of distilled water absorption, including three changes of water. Half the leaf was then illuminated, the stem being in distilled water.

From Table II we note that the leaf punches show slightly more

activity in the illuminated than in the dark side, but the chloroplasts, isolated from each half (2° C, as simultaneously as possible) possessed not only approximately equal activity, but they indicate no preferential conversion of ${}_iP^* \longrightarrow {}_oP^*$.

If one accepts the percent of total volume occupied by the chloroplasts of a tobacco leaf as equal to a spinach leaf (17%) (Mencke (7), then the specific activities of the chloroplasts were only ca. 1/10 that of the cytoplasm.

If the leaf is half-illuminated directly while absorbing P^* , a much higher P^* (~ 2.5 times) is found on the illuminated than the dark side, although the ratio of chloroplast P^* /total P^* is equal for each half. This effect is presumably due to transpirational differences.

It is often stated, even among physiologists, that the use of algae are preferable to leaves, since they are more uniform, statistically than different leaves. The comparison is not valid, since the chloroplasts within a single leaf are probably as uniform as the algae within a culture, and it is doubtful whether two algal cultures are more identical than opposite leaves. Furthermore, algae are complicated by a completeness and complexity of physiology not present within a more specialized organ as a leaf. Thus, although a variety of experiments have been done with algae, differences in behaviour of different cultures have made comparisons in experiments difficult, and the data offered are to be considered only as preliminary.

Algae

The primary comparison to be made is that of light vs. dark. Furthermore, since respiration is known to involve considerable phosphorylation, while photosynthesis requires only minute amounts of oxygen, most experiments were anaerobic.

Table I

Effect of light on P^* uptake during photochemical reduction by grana (*).

substrate	c/m P^*	gas exchanges
xs quinone, light	3620	+ 86 cmm.
xs quinone, dark	3860	- 4 cmm.
no substrate, light	3660	+ 12 cmm.
no substrate, dark	3660	+ 6 cmm.

(*). For grana preparation, see text. Grana washed once and resuspended in distilled H_2O .

Table II

Effect of light on P^* uptake in tobacco chloroplasts (*).

material	light	dark
leaf punches	373,000	340,000
total P^* in chloroplasts	5,000	6,200
oP^* in chloroplasts	570	620

(*). Leaf used $\frac{1}{2}$ hour after cutting. Area = 300 cm^2 . Stem immersed in 5 cc distilled H_2O containing 9.0×10^6 "counts" P^* . In dark 45', after which P^* solution is replaced with 20 cc distilled H_2O , let stand 15' and twice repeated. One half covered with Al foil. Leaf illuminated 30' with 300 W reflector spot lamp. Illumination intensity at leaf ca. 10,000 lux. FS 311 infra-red absorbing filter between source and leaf.

Chlorella vulgaris, grown in Emerson's solution over two 100 W white fluorescent lights (10 cm., fan circulation of air) were generally 3-4 day old cultures, which had been stored after growth at 2-3° C until use. Because of variabilities in growth and duration of storage, the algae proved to be rather unpredictable both as to length of induction period (for photosynthesis) and, to a lesser extent, respiration. This variability appeared to affect both rate and extent of P^* pickup so that experiments were directly comparable only if algae had been aliquoted from the same stock. Further work undoubtedly requires algae cultured and used under identical conditions.

A typical experiment testing the effect of light is summarized in Table III. It is apparent that no effect of light is found.

Table III

0.120 cc algae. $P^* = 0.6 \%$, $c/m = 6.0 \cdot 10^6$. $t = 23'$. Gas phases $N_2 + 4\% CO_2$. Medium 0.1 M K_4CO_3 .

	Total	Organic P	% ${}_oP$
light	101,000	38,000	38%
dark	102,000	38,000	38%

Furthermore, even in the presence of oxygen, which could conceivably be specifically coupled with the phosphorylations, a short run correlation could not be found, as is evident from Table IV.

Identical experiments with cultures in the presence of oxygen (or for protracted periods (60') in its absence) indicated that 20 minutes was generally required for equilibration between ${}_iP + {}_oP$ (i.e., in algae, ${}_iP/{}_oP = \frac{1}{4}$, so that when the % ratio is 20/80, complete

Table IV

0.100 cc algae. $eP = 0.6 \text{ } \mu\text{Ci}$, $c/m = 3 \times 10^6$, $t = 15'$. Gas phases,
air + 4% CO_2 .

	Total	dP^*	% dP^*
light	16,300	8,200	50%
dark	17,300	8,700	49%

equilibration has occurred and the continuation of the experiment beyond this time is of no further value, (e.g., see Fig. 1).

It is however, apparent from Table III that the permeability is the limiting factor, since less than 2% of the original activity had entered the cell. To calculate the distribution between cells and medium at equilibrium, assume that P enters only by exchange. Equilibrium will occur when the specific activity of $eP^* = iP^* + oP^*$. The average composition of algae may be assumed to be ~0.1% of the wet weight, so that 120 mg. of algae contains 120 μCi of P. The external radioactive solution contained 0.6 μCi P, so that at equilibrium, the "count" in the external solution should be $\left(\frac{0.6}{0.6 + 120}\right)(6.0 \times 10^6) = 5.0 \times 10^4$, while the cells should contain $\left(\frac{120}{120.6}\right)(6.0 \times 10^7) = 5.9(5) \times 10^7$. It is obvious that the cells are far from equilibrium with the external solution and that the permeability is by far the slower rate.

There is a very definite effect of oxygen in the dark, as shown in Table V. The data for oP^* appear to be somewhat high and indicate in this instance incomplete removal of iP^* .

It is to be noted that not only does the presence of oxygen accelerate the rate of conversion of $iP^* \rightarrow oP^*$, but also the permeability

of the cell, i.e., $eP \longrightarrow iP$. Although anaerobiosis does not always affect the rate of absorption of ions through cell membranes, there are numerous instances (e.g. Hoagland (5) as this, where it is believed to occur.

An attempt was therefore made to determine the relation between O_2 assimilation and P fixation. Duplicate flasks were used, one for the determination of respiratory pickup of P, another for fermentative, i.e., in the presence of argon. Respiratory rate was determined in a third vessel. Aliquots were made from the flask containing air + P^* at 5' intervals and oP^* determined. At the conclusion of the experiment, a similar analysis was made for the vessel containing argon.

Table V

0.150 cc algae. $eP = 6\%$, $c/m = 3.0 \times 10^6$. $t = 30'$. Gas Phases, $N_2 + 4\% CO_2$; argon, air- CO_2 .

	Total P	oP	$\%oP$
$N_2 + 4\% CO_2$	23,900	11,000	46%
argon	26,800	12,000	45%
air - CO_2	142,000	127,000	89%

One may calculate the relation between O_2 consumption + P conversion either from the rate formula as follows:

It has been shown earlier that the rate of entry of P^* from the external medium into the cells is very much slower than its rate of conversion within the cell to organic P^* . Therefore, neglecting $eP^* \longrightarrow iP^*$, we may write the equation for the rate of formation of oP^* from iP^* as:

$$\frac{d_oP^*}{dt} = k\left(\frac{iP^*}{iP} - \frac{oP^*}{oP}\right)$$

where, at $t = 0$, $iP^* = oP^* = 0$; and $iP = iP^* + oP^*$

The solution of the differential equation is

$$oP^* = \frac{oPiP}{oP + iP} \left[1 - e^{-k\left(\frac{oP + iP}{iPoP}\right)t} \right]$$

which may be rewritten as

$$-\ln(1 - u_oP^*) = ukt. \quad \text{where } u = \frac{oP + iP}{oPiP}$$

From the curve (Fig. 1) it is seen that half of the conversion of $iP^* \longrightarrow oP^*$ (i.e., $oP^* = 40\%$ occurs at 7.5 minutes. Substitution in the above, after utilization of the relation, $iP = iP^* + oP^*$, results in the evaluation of $k = 14.5\%/min$.

The 290 cmm. of cells are estimated to contain 290 δ P of which 80% or 232 δ , are believed to be organic (Emerson, et. al. (2)). This is equivalent to $232/290 = 0.80 \delta_oP/cmm.$ cells or 0.0258 micromoles/cmm. cells. The fraction formed is $14.5\% = 0.0038$ micromoles/min/cmm. cells. The rate of O_2 assimilation is 0.015 cmm. $O_2/min/cmm.$ cells. This is equivalent to $0.015/22.4 = 0.0067$ micromoles/min/cmm. cells. The ratio of O_2P is thus $67/380$ or approximately one atom $O/3$ atoms P.

DISCUSSION

It is evident from the experimental data presented that there is no direct connection between light and the gross formation of organic phosphorus compounds. Thus the scheme of Ruben-Lipmann is not applicable in its largest interpretation. There still remain two possible interpretations by which phosphorus might still play a direct part in photosynthesis.

Fig. 1

If an organic phosphorus fraction constituting only 1-10% of the total organic phosphorus were involved in the photochemical interchange with inorganic phosphorus, the present method would not have been able to detect an acceleration of this interchange by light. The second possibility is that the light accomplishes transformation of organic phosphorus from one form to another without passing through the inorganic state.

SUMMARY

Using radioactive phosphorus, no direct connection between gross formation of organic compounds and photosynthesis or photochemical reductions has been found to occur.

The cooperation of Dr. Victoria A. Haas in a portion of this work is gratefully acknowledged.

Bibliography

1. Aronoff, S., Photochemical reduction of chloroplast grana, *Plant Physiology*, 21, 393-409 (1946).
2. Emerson, R. L., Stauffer, J. F., and Umbreit, W. M., Relationships between phosphorylation and photosynthesis in chlorella, *Am. J. Bot.*, 31, 107-120 (1944).
3. Fan, C. S., Stauffer, J. F., and Umbreit, W. M., An experimental Separation of oxygen liberation from carbon dioxide fixation in photosynthesis by chlorella, *J. Gen. Physiol.*, 27, 15-28 (1943).
4. Frenkel, A. W., Photosynthesis with radioactive carbon and the distribution of the intermediate products in the plant cell, *Plant Physiol.*, 16, 654-55 (1941).
5. Hoagland, D. R., Lectures in the inorganic nutrition of plants, Chronica Botanica Co., 1944.
6. Lipmann, F., and Tuttle, L. C., The condensation of acetyl phosphate with formate of CO₂ in bacterial extracts, *J. Biol. Chem.*, 158, 505-519 (1945).
7. Mencke, W., quoted in Rabinowitch, E. I., *Photosynthesis*, I, Interscience Publishers, Inc., 1945.
8. Ruben, S., Photosynthesis and phosphorylation, *J. Am. Chem. Soc.*, 65, 279-282 (1943).
9. Warburg, O., and Lüttgens, W., Carbon dioxide assimilation, *Naturwissenschaften*, 32, 301 (1944); *ibid* 32, 161 (1944).