

Special Review of  
Declassified Reports

Authorized by USDOE JK Bratton  
Unclassified TWX P182206Z May 79

*UCRL-50 1948*

**REPORT PROPERLY DECLASSIFIED**

*3-25-80*

Date

UCRL 50  
*cy. 1*


Authorized Reviewer  
Date  
CLASSIFICATION  
Y OF  
NIA

*Laboratory*

**FOR REFERENCE**

NOT TO BE TAKEN FROM THIS ROOM

BERKELEY, CALIFORNIA



UNIVERSITY OF  
ALABAMA

LIBRARY

UNIVERSITY OF CALIFORNIA  
Radiation Laboratory  
Contract No. W-7405-eng-48

ARTIFICIAL ALPHA-ACTIVE BISMUTH ISOTOPES

by

D. H. Templeton and I. Perlman

Berkeley, California

~~RESTRICTED~~  
CLASSIFIED BY THE DEPARTMENT OF ENERGY  
BY THE DEPARTMENT OF ENERGY AUTHORITY  
COMMITTEE

## ARTIFICIAL ALPHA-ACTIVE BISMUTH ISOTOPES

D. H. Templeton and I. Perlman

Radiation Laboratory and Department of Chemistry  
University of California, Berkeley, California

## ABSTRACT

Four new alpha-emitters of half lives 2 min., 9 min., 27 min., and 1-2 hr. have been found from the bombardment of lead with high energy deuterons. The three of longest half-life have been identified as isotopes of bismuth. It is believed that the mass numbers of all of the isotopes are less than 203.

Contract No. W-7405-eng-48

To be published as a letter to the Editor of Physical Review

~~RESTRICTED~~  
CLASSIFICATION CONTINUED BY AUTHORITY  
OF THE SECRETARY OF DEFENSE  
BY THE DECLASSIFICATION COMMITTEE

## ARTIFICIAL ALPHA-ACTIVE BISMUTH ISOTOPES

D. H. Templeton and I. Perlman

Radiation Laboratory and Department of Chemistry  
University of California, Berkeley, California

When lead was bombarded with approximately 200-Mev deuterons in the 184-inch cyclotron a great variety of nuclear reactions were noted. Fission has been demonstrated by the identification of radioactive fission products<sup>(1)</sup> and a mixture of isotopes from Bi (element 83) to Au (element 79) have been partially identified. These were produced by reactions analogous to those reported from other irradiations with high energy particles<sup>(2)</sup>.

This report has to do with some new short-lived emitters of alpha-particles produced from high energy deuterons on lead. The alpha-activity was in very low abundance compared with the Geiger counter activity and was first observed in the unseparated target material. The decay curves could be resolved into 2 min., 9 min., 27 min., and 1-2 hr. components. The longest period was too weak for accurate half-life determination. The three longest periods were chemically identified with the bismuth fraction but the chemical separation was too slow to permit observation of the 2-min. period. Lead which was bombarded with 100 Mev deuterons showed the 1-2 hr. and 27 min. periods but the shorter periods could not be detected. Therefore, the 2-min. and 9-min. activities are probably of lower mass number than the longer-lived ones. All of these isotopes of bismuth are believed to be of lower mass number than 203 since lead, enriched in  $\text{Pb}^{204}$ , did not show any alpha-activity when bombarded with 20-Mev deuterons<sup>(3)</sup>.

The alpha-particle energies of the three longer periods were all in the range of  $5.5 \pm 0.3$ -Mev as measured by an alpha-particle pulse analyzer<sup>(4)</sup>. The lack of precision was caused by the high background of electrons, and by the inability to prepare extremely thin samples. Because these energies seem too low for alpha-particle half-lives of the order of minutes and since the bismuth fraction showed a great preponderance of electrons over alpha-particles it is

concluded that the alpha-emission represents but a small branching in the predominantly orbital electron capture decay. For example, a 30-min. period and a 100-min. period could be resolved out of the Geiger-counter decay curve and under the assumption that these are the same as the 27-min. and 1-2 hr. alpha periods, alpha-particle to electron ratios of  $7 \times 10^{-4}$  and  $6 \times 10^{-5}$ , respectively, were calculated. The half-lives for alpha-decay for the 27-min. and 1-2 hr. activities then become about 1-month and 3-years if it is assumed that there is almost one electron emitted per orbital electron capture disintegration.

These data show that alpha-instability as reflected by the alpha-particle half-lives follows the same trend in the case of bismuth isotopes as that noted for the isotopes of polonium. In the case of polonium it has long been noted that there is a minimum half-life at ThC' ( $\text{Po}^{212}$ ) and that the half-lives increase both toward lower and higher mass numbers. This curve has been extended<sup>(5)</sup> to show that the half-life for  $\text{Po}^{208}$  is even longer than for  $\text{Po}^{210}$  but that the partial alpha half-life then decreases again at lower mass numbers as exemplified by  $\text{Po}^{206}$ . In the case of bismuth, the earlier noted half-lives showed a decrease from heavier masses toward AcC ( $\text{Bi}^{211}$ ) which has a 3-min. half-life. Recently Broda and Feather<sup>(6)</sup> reported that RaE( $\text{Bi}^{210}$ ) undergoes rare alpha-branching corresponding to a partial alpha half-life of about  $10^4$ - $10^5$  yrs. Below RaE the half-life increases still further as shown by the failure to detect alpha-activity in  $\text{Bi}^{209}$ . However, the present studies show that very light isotopes of bismuth ( $\text{Bi}^{203}$ ) again have measurable alpha half-lives.

We wish to acknowledge the assistance and cooperation of Dr. John Woodyard, Dr. D. C. Sewell and all those whose operation of the 184-inch cyclotron made these irradiations possible.

This paper is based on work performed under Contract Number W-7405-eng-48 with the Atomic Energy Commission in connection with the Radiation Laboratory of the University of California, Berkeley, California.

## References

- (1) Perlman, I., R. H. Goeckermann, D. H. Templeton and J. J. Howland, Phys. Rev. 72, 352 (1947).
- (2) Cunningham, B. B., H. H. Hopkins, M. Lindner, D. R. Miller, P. R. O'Connor, I. Perlman, G. T. Seaborg and R. C. Thompson, Phys. Rev. 72, 739 (1947).
- (3) Templeton, D. H., J. J. Howland and I. Perlman, Phys. Rev. 72, 766 (1947).
- (4) Ghiorso, A., A. H. Jaffey, H. P. Robinson and B. Weissbourd, "An Alpha Pulse Analyzer Apparatus" (Plutonium Project Record, Vol. 1413, 17.3 (1946)) (to be issued).
- (5) Templeton, D. H., J. J. Howland and I. Perlman, Phys. Rev. 72, 758 (1947).
- (6) Broda, E. and N. Feather, Proc. Roy. Soc., London A190, 20 (1947).